

NOTIZEN

The Mössbauer-Effect of the 104 keV Transition in ^{180}W

K. ZIOUTAS and B. WOLBECK

Institut für Strahlen- und Kernphysik, Universität Bonn

(Z. Naturforsch. 26 a, 1088 [1971]; received 22 April 1971)

The Mössbauer effect of the $2+ (104 \text{ keV}) 0+$ transition in ^{180}W has been observed. The source was ^{180}Ta which was produced by irradiation of a foil of natural tantalum by fast neutrons: $^{181}\text{Ta}(n, 2n)^{180}\text{Ta}$. The width of the resonance absorption line was $(3.1 \pm 0.8) \text{ mm/sec}$ which corresponds to the linewidth derived from lifetime measurements.

Whereas Mössbauer measurements have been performed in the tungsten isotopes $^{182}, ^{183}, ^{184}, ^{186}\text{W}$, the ME in ^{180}W has not been observed previously. The expected linewidth of about 2.2 mm/sec of the 104 keV transition should open the possibility to study the electric and magnetic hyperfine interaction of this nucleus in various surroundings.

In order to obtain an unsplit source we used W in tantalum metal which has cubic symmetry. A Ta foil was irradiated by fast neutrons of about 15 MeV which were produced by bombarding beryllium with 30 MeV deuterons in the Karlsruhe-Cyclotron. Neutrons with an energy of 15 MeV have a cross-section of 2 barn for the desired process $^{181}\text{Ta}(n, 2n)^{180}\text{Ta}$. All other possible reactions are much weaker. The $2+$ state in ^{180}W is populated only to 3% in the decay of ^{180}Ta ($T_{1/2} = 8.1 \text{ h}$), while a 27% branch leads to the $93 \text{ keV } 2+$ state in ^{180}Hf .

For the absorber we used 50 mg/cm^2 W-metal powder (^{180}W enriched to 6.9%) which was obtained by reducing WO_3 in a hydrogen atmosphere at 900°C .

For the detection of the $104 \text{ keV } \gamma$ -line we used a Ge(Li)-detector with 1 keV resolution at 100 keV . In this way any background from other lines could be avoided. In particular the contribution of the $100 \text{ keV } \gamma$ -rays of the ^{182}W produced by the weak (n, γ) reaction could be reduced to less than 0.1% .

Our experiment was performed by use of a standard Mössbauer transmission apparatus at a temperature of 20°K .

Fig. 1 shows the observed single line spectrum of the $2+ \rightarrow 0+$ transition in ^{180}W with a least squares fitted Lorentzian. The total time for the measurement was two days and three different sources were used.

Reprints request to Dipl.-Phys. K. ZIOUTAS, Institut für Strahlen- und Kernphysik der Universität Bonn, D-5300 Bonn, Nußallee 14–16.

¹ B. PERSSON, H. BLUMBERG, and D. AGRESTI, Phys. Lett. **24B**, 522 [1967].

² D. AGRESTI, E. KANKELEIT, and B. PERSSON, Phys. Rev. **155**, 1342 [1967].

³ G. W. SCHWEIMER, Gesellschaft für Kernforschung m.b.H., Karlsruhe KFK 504 [1966].

The observed linewidth was $(3.1 \pm 0.8) \text{ mm/sec}$. After correction for the finite absorber thickness one obtains a lower limit for the half-life of the $2+$ state of $T_{1/2} > (0.91 \pm 0.35) \text{ ns}$. This value is in agreement with lifetime measurements⁵.




Fig. 1. Mössbauer spectrum of the 104 keV transition of ^{180}W at 20°K .

The Mössbauer effect enables an accurate determination of the magnetic moment and the quadrupole moment of the $2+$ level in ^{180}W . This would extend the systematics for the even tungsten isotopes towards the strongly deformed region which is of great theoretical interest^{6–8}.

Acknowledgement

We thank Prof. E. BODENSTEDT for his interest in this experiment. Particularly we are indebted to Dr. W. G. SCHWEIMER in Karlsruhe for very useful discussions and the preparation of the sources. — This work was financially supported by the Bundesministerium für Bildung und Wissenschaft.

⁴ I. S. BRZOSKO, E. GIERLIK, A. SOLTAN, JR., and Z. WILHELM, Nucl. Phys. A **123**, 603 [1969].

⁵ W. M. CURRIE, Nucl. Phys. **47**, 551 [1963].

⁶ O. PRIOR, F. BOEHM, and S. G. NILSSON, Nucl. Phys. A **110**, 257 [1968].

⁷ K. KUMAR and M. BARANGER, Nucl. Phys. A **122**, 273 [1968].

⁸ Y. W. CHOW et al., Phys. Lett. **30b**, 171 [1969].



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.