NMR Study of 3,5-Dichloropyridine in the Nematic Phase of the Liquid Crystal

M. ŽAUCER and A. AŽMAN

Chemical Institute Boris Kidrič and Department of Chemistry University of Ljubljana, Ljubljana, Yugoslavia

(Z. Naturforsch. 27 a, 1535 [1972]; received 14 August 1972)

The importance of NMR studies on oriented molecules dissolved in the nematic phase of liquid crystals is well recognized ¹. The aim of this note is to report such a study of 3,5-dichloropyridine in the nematic phase of N-(p-Methoxy-benzyliden)-p-n-butylanilin (MBBA). The observed spectrum is that of an AB₂ symmetry with nine detected lines (Fig. 1). The analysis ¹ of this spectrum is straightforward and the values for the direct (D) coupling constants, the indirect (J) coupling constants and the elements of the orientation matrix S are:

$$\begin{split} D_{\rm AB} = - & 51 \; {\rm Hz} \; , & S_{11} = 0.003 \; , \\ D_{\rm BB} = - & 241 \; {\rm Hz} \; , & S_{22} = 0.139 \; . \\ J_{\rm AB} = & \pm & 2 \; {\rm Hz} \; , \end{split}$$

J was supposed to have the same value as in the isotropic phase and elements of the S matrix were obtained by using the geometry of pyridine 2 . The values of D suggest that the molecule is preferentially oriented with the axes B-B along the magnetic field. The anisotropies of the chemical shifts are $\sigma_a\!=\!0.46$ ppm for protons A and $\sigma_a\!=\!0.06$ ppm for protons B. These shifts are towards higher fields and support the mentioned conclusion about the orientation of the molecule. We could not explain in a simple manner why the chemical shifts are so different for A and B protons.

Acknowledgement

This work was partially supported by the Boris Kidrič Fund.




Fig. 1. The spectrum of an AB2 symmetry with nine detected lines.

- P. DIEHL and C. L. KHETRAPAL, NMR Basic Principles and Progress, Vol. 1, ed. P. DIEHL, E. FLUCK, and R. KOSFELD, Springer-Verlag, Berlin 1969.
- ² Tables of Interatomic Distances and Configuration in Molecules and Ions, ed. L. E. SUTTON, Special Publication No. 18, Chemical Society, London.