

Orbital Valence Force Field Constants of Tetrachlorides, Tetrabromides and Tetraiodides of Titanium, Zirconium and Hafnium

A. N. PANDEY, D. K. SHARMA, and H. P. MITAL

Department of Physics, Meerut College, Meerut (U.P.), India

(Z. Naturforsch. 27 a, 1211—1212 [1972]; received 2 June 1972)

The orbital valence force field has been employed to calculate the force constants for tetrachlorides, tetrabromides, and tetraiodides of titanium, zirconium, and hafnium making use of recent vapor-phase Raman spectral data. The results are compared with the Urey-Bradley force constants, and it has been found that these are well comparable. The applicability of the Lennard-Jones potential has been examined to understand the nature of non-bonded interactions. The force constants have also been utilized to study the relative strength of the chemical bonds in tetrahalides and hexahalide anions

Very recently CLARK et al.^{1, 2} have studied the Raman spectra of the tetrachlorides, tetrabromides and tetraiodides of titanium, zirconium and hafnium in the vapor-phase. Complete assignments of all the fundamentals, on the basis of tetrahedral symmetry, have been given. They have also reported the thermodynamic functions and force constants for the molecules employing the modified valence force field (MVFF) as well as the Urey-Bradley force field (UBFF). In the present communication it is aimed (i) to apply the orbital valence force field (OVFF) to compute the force constants, (ii) to compare these force constants with the corresponding UBFF force constants, (iii) to discuss the relative stability of the chemical bonds in different molecules, and (iv) to examine the applicability of the Lennard-Jones potential to understand the nature of non-bonded forces.

WILSON's³ FG matrix method has been used to evaluate the force constants employing OVFF. The expressions for the *F* and *G* matrices are taken from KREBS and MÜLLER⁴. For the detail procedure one may refer to⁵.

First, all the four OVFF constants; *k*₁, the bond stretching force constant; *k*_α', the angle distortion constant; *A* and *B/R*, the interaction constants representing forces between nonbonded atom pairs, were evaluated with the help of four force field equations⁴. The OVFF constants, thus calculated, are presented in Table 1 together with the corresponding UBFF constants¹. The fundamental frequencies used in these computations are listed in Table 2. It is apparent from Table 1 that the OVFF

constants, *k*₁, *k*_α', and *A* are in good agreement with the corresponding UBFF constants, *K*, 3 *H*, and *F*/2. As shown in this table, the present calculations lead to a negative value of *B/R* for most of the molecules. From this it follows that the OVFF model,

Table 1. OVFF constants (in mdyne/Å) for the group IV b tetrahalides. (The values in parentheses indicate UBFF constants *K*, 3 *H*, *F*/2 and $-F'$.)

Halide	<i>k</i> ₁ , <i>K</i>	<i>k</i> _α ', 3 <i>H</i>	<i>A</i> , <i>F</i> /2	<i>B/R</i> , $-F'$
TiCl ₄	2.5095	0.1185	0.0813	— 0.0098
	(2.509)	(0.105)	(0.0815)	(+ 0.005)
	2.4355 *	0.0762 *	0.0906 *	+ 0.0139 *
TiBr ₄	2.1597	0.1453	0.0454	— 0.0152
	(2.150)	(0.126)	(0.0465)	(+ 0.003)
	2.0761 *	0.1005 *	0.0558 *	+ 0.0085 *
TiI ₄	1.6030	0.1116	0.0448	— 0.0069
	(1.619)	(0.108)	(0.043)	(+ 0.002)
	1.548 *	0.0830 *	0.0517 *	+ 0.0079 *
ZrCl ₄	2.2961	0.0257	0.0840	+ 0.0067
	(2.296)	(0.024)	(0.084)	(+ 0.010)
	2.2786 *	0.0148 *	0.0862 *	+ 0.0132 *
ZrBr ₄	2.0953	0.1064	0.0373	— 0.0116
	(2.093)	(0.093)	(0.037)	(+ 0.002)
	2.0361 *	0.0731 *	0.0447 *	+ 0.0068 *
ZrI ₄	1.6325	0.0882	0.0292	— 0.0085
	(1.627)	(0.075)	(0.030)	(+ 0.003)
	1.5846 *	0.0623 *	0.0352 *	+ 0.0054 *
HfCl ₄	2.369	0.0327	0.0848	+ 0.0128
	(2.373)	(0.030)	(0.084)	(+ 0.017)
	2.3687 *	0.0323 *	0.0848 *	+ 0.0130 *
HfBr ₄	2.1146	0.0664	0.0620	— 0.0036
	(2.117)	(0.060)	(0.0615)	(+ 0.004)
	2.0757 *	0.0427 *	0.0668 *	+ 0.0102 *
HfI ₄	1.8430	0.2489	0.0029	— 0.0286
	(1.844)	(0.219)	(0.0025)	(+ 0.002)
	1.7477 *	0.1942 *	0.0148 *	+ 0.0022 *

* OVFF constants using Lennard-Jones potential function (*A* = 6.5 *B/R*).

Reprint requests to A. N. PANDEY, Department of Physics, Meerut College, Meerut (U.P.), Indien.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

making use of a single term in B/R , cannot represent satisfactorily the interaction between non-bonded atom pairs. Negative values of B/R have also been reported in ^{4, 5}.

The force constants are a convenient measure of the strength of a chemical bond and, therefore, it is interesting to study the trend of the force constants among the tetrahalides (except tetrafluoride). The only force constant sufficiently large to warrant discussion is the bond stretching force constant (k_1). A comparison of the stretching force constant k_1 , from Table 1, shows that it decreases in the order $MCl > MBr > MI$ ($M = Ti, Zr$ or Hf). The relative strength of the chemical bond must vary in the same order. The interaction constant, A , follows the same pattern and supports the above conclusion. The variation of the interaction constant is corroborated by the fact that for smaller non-bonded distances its magnitude should be larger, as it happens here¹. The angle bending force constant, k_a' , does not show a regular variation.

The OVFF bond stretching force constants of hexachloride and hexabromide anions⁶ with those

Table 2. Comparison of the observed and calculated fundamental frequencies (in cm^{-1}) for the group IV b tetrahalides using OVFF constants from Table 1 with the approximation $A = 6.5 B/R$.

Halide		$\nu_1(A)$	$\nu_2(E)$	$\nu_3(F_2)$	$\nu_4(F_2)$
TiCl ₄	Obs.	389	114	498	136
	$A = 6.5 B/R$	389	114	493	152.3
TiBr ₄	Obs.	231.5	68.5	393	88
	$A = 6.5 B/R$	231.5	68.5	389	89.4
TiI ₄	Obs.	162	51	323	67
	$A = 6.5 B/R$	162	51	321.6	72.9
ZrCl ₄	Obs.	377	98	418	113
	$A = 6.5 B/R$	377	98	417	117.6
ZrBr ₄	Obs.	225.5	60	315	72
	$A = 6.5 B/R$	225.5	60	312.8	82
ZrI ₄	Obs.	158	43	254	55
	$A = 6.5 B/R$	158	43	253.4	61.5
HfCl ₄	Obs.	382	101.5	390	112
	$A = 6.5 B/R$	382	101.5	406.5	105.4
HfBr ₄	Obs.	235.5	63	273	71
	$A = 6.5 B/R$	235.5	63	271	78.4
HfI ₄	Obs.	158	55	224	63
	$A = 6.5 B/R$	158	55	220.3	74

of the tetrachlorides and tetrabromides of titanium, zirconium, and hafnium, which possess the same oxidation state⁴, are compared in Table 3. From this comparison it is inferred that the chemical bond in the tetrahedral species is stronger than in the octahedral species.

Table 3. Comparison of the bond stretching force constants (in $\text{mdyne}/\text{\AA}$) of the tetrachlorides and tetrabromides with the hexachloride and hexabromide anions of Ti, Zr, and Hf.

Halide	k_1	Halide anion	K
TiCl ₄	2.5095	TiCl ₆ ²⁻	0.68
ZrCl ₄	2.2961	ZrCl ₆ ²⁻	1.03
HfCl ₄	2.3690	HfCl ₆ ²⁻	1.12
TiBr ₄	2.1597	TiBr ₆ ²⁻	0.60
ZrBr ₄	2.0953	ZrBr ₆ ²⁻	0.74
HfBr ₄	2.1146	HfBr ₆ ²⁻	0.78

In order to understand the nature of the forces acting between the non-bonded atom pairs, a second set of OVFF constants has been calculated by approximating the interaction by a Lennard-Jones potential which includes the dispersion forces. This potential function establishes a relation $A = 6.5 B/R$ between the non-bonded interaction constants. The set of force constants thus evaluated is included in Table 1. For testing the validity of this approximation these constants, in turn, are employed to calculate the fundamental frequencies. The calculated and observed frequencies are compared in Table 2. It is found that, except TiBr₄, the frequencies are not well reproduced. Therefore, it is concluded that for titanium tetrabromide the Lennard-Jones potential is adequate to explain the interaction between non-bond atom pairs while in all other cases it is not a reasonable approximation. To account for these discrepancies Coulomb forces besides dispersion forces should also be taken into consideration as suggested by KREBS and MÜLLER⁴.

Acknowledgements

The authors are grateful to Prof. Dr. A. MÜLLER, University of Dortmund, West Germany, for his continued interest and encouragement.

¹ R. J. H. CLARK, B. K. HUNTER, and D. M. RIPPON, *Inorg. Chem.* **11**, 56 [1972].

² R. J. H. CLARK and C. J. WILLIS, *J. Chem. Soc. A*, 838 [1971].

³ E. B. WILSON, JR., J. C. DECIUS, and D. C. CROSS, *Molecular Vibrations*, McGraw-Hill, New York 1955.

⁴ B. KREBS and A. MÜLLER, *J. Mol. Spectroscopy* **22**, 290 [1967].

⁵ B. B. SRIVASTAVA, A. K. DUBLISH, A. N. PANDEY, and A. K. MITHAL, *Z. Naturforsch.* **27a**, 1213 [1972].

⁶ M. N. AVASTHI and M. L. MEHTA, *Z. Naturforsch.* **26a**, 1137 [1971].