

Notizen

⁶⁷Zn NMR Anomalous Solvent Isotope Effect in Aqueous SolutionsB. W. Epperlein, H. Krüger, O. Lutz,
and A. Schwenk

Physikalisches Institut der Universität Tübingen, Germany

(Z. Naturforsch. **29 a**, 660–661 [1974];
received February 15, 1974)

For ⁶⁷Zn NMR lines of solutions of ZnCl₂, ZnBr₂, and ZnI₂ in H₂O and D₂O an anomalous solvent isotope effect is reported. In D₂O solutions the lines are shifted to higher frequencies. The difference between the shieldings in H₂O and D₂O is e. g. $\sigma(\text{H}_2\text{O}) - \sigma(\text{D}_2\text{O}) = (13.1 \pm 0.7)$ ppm for a concentration of 0.02 moles zinc bromide per mole solvent.

The frequency of the NMR signal (Larmor frequency) of a nucleus in an ion in aqueous solution can depend on the isotopic composition of the solvent. Such a behaviour has been found for alkali^{1–4}, halide^{1, 5, 6} and several other nuclei^{7–11} in aqueous salt solutions. The largest effect was observed for ²⁰⁷Pb in aqueous lead salt solutions⁷. The difference between the shieldings in H₂O and D₂O is $\sigma(\text{H}_2\text{O}) - \sigma(\text{D}_2\text{O}) = -(31 \pm 3)$ ppm for a concentration of 0.004 moles lead nitrate per mole solvent. These shielding differences are presumably due to the different vibronic states of the H₂O and D₂O molecules surrounding the ions⁵.

For ¹¹¹Cd an indication for a shift to higher frequency was found in a cadmium chloride solution¹¹, whereas in perchlorate and nitrate solutions the usual solvent isotope effect was observed.

During a NMR investigation of IIb elements, ⁶⁷Zn NMR signals were observed in aqueous solutions for a determination of the shielding constant¹². The ⁶⁷Zn studies were done at 4.841 MHz in a field of 1.807 Tesla with a Fourier Transform Spectrometer basing on a Bruker pulse spectrometer KR 322 s. For experimental details see for example².

The dependence of the Larmor frequency of ⁶⁷Zn on the concentration of aqueous zinc chloride, bromide, and iodide solutions was measured. The observed linewidths are in the range between 20 Hz for low and 170 Hz for high concentrations. For all these salts an anomalous shift to higher frequencies

was found for the solutions in D₂O, compared with those in H₂O. Results are given in the Table 1.

In other zinc salts, like perchlorate, nitrate, and sulfate, contrarily to cadmium salts, no solvent isotope effect was observed within an experimental error of ± 1 ppm. But there is also no detectable

Table 1. ⁶⁷Zn chemical shifts in aqueous zinc salt solutions at (300 ± 2) K using cylindrical samples; no bulk susceptibility correction was made, since this correction is smaller than the given errors (standard deviation).

Salt	Concentration moles salt moles solvent	⁶⁷ Zn NMR line shift (to higher frequency in Hz *		⁶⁷ Zn shielding $\sigma(\text{H}_2\text{O}) - \sigma(\text{D}_2\text{O})$ in ppm **
		H ₂ O	D ₂ O	
ZnCl ₂	0.02	341	363	4.6
	0.04	518	531	2.7
	0.06	596	605	1.9
ZnBr ₂	0.02	125	188	13.1
	0.04	314	356	8.7
	0.06	421	446	5.2
ZnI ₂	0.02	48	61	2.7
	0.04	109	125	3.3
	0.06	136	148	2.5

* Relative to vanishing concentration of zinc salts in H₂O.

** The error is smaller than ± 1 ppm.

dependence of the Larmor frequency on the concentration of these salts¹².

The reason for the difference between the normal and anomalous solvent isotope effect must be looked for in the strong halide complexes which are found in zinc and cadmium solutions. For ⁵⁵Mn in the MnO₄[−]-ion, the normal isotope effect is reported¹⁰.

The magnetic shielding of zinc ions by water was evaluated in Ref.¹²: $\sigma^* = -(6.9 \pm 0.1) \cdot 10^{-4}$. With this number the given differences of Table 1 must be compared. The relative solvent isotope effect is therefore for the first bromide solution:

$$\frac{\sigma(\text{H}_2\text{O}) - \sigma(\text{D}_2\text{O})}{\sigma^*(\text{H}_2\text{O})} = -1.9 \cdot 10^{-2}.$$

For comparison the following values of ¹³³Cs and ⁸⁷Rb are reported⁶: $0.35 \cdot 10^{-2}$ and $0.14 \cdot 10^{-2}$.

Acknowledgment. We thank the Deutsche Forschungsgemeinschaft for the financial support.

¹ A. Loewenstein, M. Shporer, P. C. Lauterbur, and J. E. Ramirez, Chem. Comm. **1968**, 214.

² O. Lutz and A. Nolle, Z. Naturforsch. **27 a**, 1577 [1972].

³ J. Halliday, H. D. W. Hill, and R. E. Richards, Chem. Comm. **1969**, 219.

Reprint requests to Dr. O. Lutz, Physikalisches Institut der Universität Tübingen, D-7400 Tübingen, Morgenstelle 14, Germany.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

- ⁴ J. D. Halliday, R. E. Richards, F. R. S. and R. R. Sharp, Proc. Roy. Soc. Lond. A **313**, 45 [1969].
⁵ C. Deverell, K. Schaumburg, and H. J. Bernstein, J. Chem. Phys. **49**, 1276 [1968].
⁶ J. Blaser, O. Lutz, and W. Steinkilberg, Z. Naturforsch. **27 a**, 72 [1972].
⁷ O. Lutz and G. Stricker, Phys. Letters **35 A** 397 [1971].
⁸ B. W. Epperlein and O. Lutz, Z. Naturforsch. **23 a**, 1413 [1968].

- ⁹ O. Lutz, Phys. Letters **29 A**, 58 [1969].
¹⁰ O. Lutz and W. Steinkilberg, Phys. Letters **30 A**, 183 [1969].
¹¹ H. Krüger, O. Lutz, A. Schwenk, and G. Stricker, Z. Physik **266**, 233 [1974].
¹² B. W. Epperlein, H. Krüger, O. Lutz, and A. Schwenk, Phys. Letters **45 A**, 255 [1973].

Cross Sections for the Destruction of an Alignment in the Metastable 6^3P_2 -State of Hg by Collisions with H_2 , N_2 and CO_2

M. Baumann, E. Jacobson and W. Koch

Physikalisches Institut der Universität Tübingen

(Z. Naturforsch. **29 a**, 661–662 [1974];
received February 2, 1974)

The cross sections for destruction of an alignment in the metastable 6^3P_2 -state of mercury atoms by collisions with H_2 , N_2 and CO_2 molecules have been determined from the observation of transient signals. The results were (in units of 10^{-16} cm^2) H_2 : 63(7); N_2 : 104(13); CO_2 : 663(68).

The cross sections for depolarization of aligned Hg atoms in the metastable 6^3P_2 -state by collisions with noble gases have been studied by several authors^{1–4}. It seemed to be of some interest to extend these measurements to collisions with other molecules. In this paper we report on an investigation of relaxation processes caused by collisions with H_2 , N_2 and CO_2 molecules.

The experimental arrangement was as follows: The Hg atoms (even isotopes) were excited to the metastable 6^3P_2 -state in a vhf. discharge ($\nu = 215 \text{ MHz}$) which was driven by the electric field (about 150 V/cm) between the two plates of a capacitor outside the resonance vessel which contained the mercury vapour and the foreign gas. The direction of the electric field vector was chosen parallel to an external static magnetic field H_0 ($\approx 2.6 \cdot 10^{-4} \text{ Tesla}$). With these conditions a longitudinal component of an alignment in the excited state could be produced⁵. It was necessary to work with a continuously pumped system instead of the usual sealed-off resonance cell in order to avoid variations of the foreign gas pressure by clean-up effects in the gas discharge⁶. By stimulating rf transitions $\Delta m = \pm 1$ between the Zeeman sublevels of the metastable state by a magnetic rf field H_1 perpendicular to H_0 the alignment will be disturbed. The change of the alignment can be monitored by the absorption of linearly polarized radiation at $\lambda = 5461 \text{ Å}$ corre-

sponding to the transition $6^3P_2 \rightarrow 7^3S_1$. After switching off the disturbing rf field the alignment will be restored according to an exponential time law, which leads to transient signals in the absorption with a time constant τ_2 . τ_2 is the relaxation time for collisional destruction of the alignment in the metastable state⁷. The transient signals have been detected by means of a sampling technique which was described elsewhere^{7, 8}.

Figure 1 and Fig. 2 show the pressure dependence of the relaxation rate $1/\tau_2$ for H_2 , N_2 and CO_2 .




Fig. 1. Pressure dependence of the relaxation rate $1/\tau_2$ for H_2 and N_2 .




Fig. 2. Pressure dependence of the relaxation rate $1/\tau_2$ for CO_2 .

Reprint requests to Prof. Dr. M. Baumann, Physikalisches Institut der Universität Tübingen, D-7400 Tübingen, Morgenstelle 14.