

A Simple Variational Method for Quantitative Assessment of "Through-space" and "Through-bond" Interactions

Hartmut Schmidt and Armin Schweig

Fachbereich Physikalische Chemie der Universität
3550 Marburg (Lahn), Germany

(Z. Naturforsch. 31 a, 215–216 [1976];
received December 30, 1975)

A simple method is described for separating "through-space" from "through-bond" interactions. The method is applied to 1,4-cyclohexadiene, 2,5-dihydrofuran, 2,5-dihydrothiophene, and 2,5-dihydropyrrole.

It was recently shown^{1–9} that the π orbital "localization" procedure of Baird^{10, *} and a variant^{9, 11, *} of it are very useful for a quantitative treatment of several fundamental problems (as e.g. orbital interactions, aromaticity, inductive and conjugative effects or conformational forces) in theoretical organic chemistry. In this communication we proceed to study the potentiality of the method for a quantitative assessment¹² of "through-space" and "through-bond" interactions^{13, 14, **}.




Fig. 1. Calculated interaction diagram for 1,4-cyclohexadiene using the CNDO/S method. The basis orbital energy, the "through-space" (\longleftrightarrow), "through-bond" (\rightleftharpoons) and total (\longleftrightarrow) effects are evaluated on the base of the two individual π decoupled models (1) and (2) and the fully coupled case (3). For details see text.

To this end, we choose the well-known 1,4-cyclohexadiene case^{12, 15–17} (Figure 1). Here, three models, (1) to (3), are considered: (1) means complete π conjugative decoupling of one double bond from the rest of the molecule, (2) means π isolation of both double bonds from the methylene groups, and (3) refers to a usual full π coupled calculation. With (1) we interrupt both "through-space" and "through-bond" interactions, with (2) we cut off the "through-bond" but retain the "through-space" process, and with (3) both events remain intact.

In the actual calculation the planar geometry^{18–21} of 1,4-cyclohexadiene was taken and the CNDO/S (MINDO/2) method used. We obtain a "through-space" splitting of 0.82 (0.59) eV, a "through-bond" destabilization of the π_+ orbital of 2.05 (1.84) eV so that the predicted orbital sequence for 1,4-cyclohexadiene is π_+/π_- and the energy difference between both orbitals 1.21 (1.21) eV⁺. It is gratifying to note that these values agree well with the MINDO/2 results ("through-space" splitting = 0.66 eV, "through-bond" destabilization of the π_+ orbital = 1.86 eV, energy difference between π_+ and π_- = 1.20 eV) of Heilbronner and Schmelzer¹². These authors obtained their results on the base of a rather different and seemingly more complicated perturbational approach. The present variational method is simple and easy to apply.

To check the consistency of the method we considered an additional model* (4) which admits "through-bond" but excludes "through-space" interaction. In this case the π_+/CH orbital is destabilized by 0.31 eV and the π_- orbital stabilized by 0.41 eV relative to case (3) in harmony with what is expected from (2). This result of near additivity of effects corroborates the method used.

The method was further applied to the question of "through-space" interaction in 2,5-dihydrofuran²², 2,5-dihydrothiophene²³, and 2,5-dihydropyrrole²⁴. Taking the π lone pair orbital on the heteroatom as an indicator we obtained as percentage[†] "through-space" interactions in this series: 19%, 20%, and 83%, respectively. Thus, for 2,5-dihydropyrrole, "through-space" interaction may be relevant.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie. The calculations were carried out using the TR4 computer at the Rechenzentrum der Universität Marburg.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

- * In the Baird method π orbital localization on a selected molecular subunit is accomplished by setting the SCF matrix elements between any π atomic orbital of this subunit and any π atomic orbital of the molecular remainder zero. Instead of this full decoupling our variant method allows us to set individually selected matrix elements zero so that individual interactions (as e.g. here the "through-space" interaction) can be selectively investigated.
- ** Current usage equates "through-space" interactions with "homoconjugative" or "transannular" interactions and "through-bond" interactions with "hyperconjugative" interactions.
- + The measured separation between the two π ionization events is 1.0 eV; see Reference ¹⁵.
- ++ Taken to be planar and the nitrogen atom sp^2 hybridized.
- + Referred to the calculated "through-bond" interaction.
- ¹ H. L. Hase and A. Schweig, *Tetrahedron* **29**, 1759 [1973].
- ² C. Müller, A. Schweig, and H. Vermeer, *Angew. Chem.* **86**, 275 [1974]; *Angew. Chem. Internat. Edn.* **13**, 273 [1974].
- ³ H. Schmidt, A. Schweig, R. W. Hoffmann, and H. Kurz, *Tetrahedron Lett.* **1974**, 1953.
- ⁴ H. Schmidt, A. Schweig, and A. Krebs, *Tetrahedron Lett.* **1974**, 1471.
- ⁵ W. Schäfer, A. Schweig, G. Maier, T. Sayrac, and K. J. Crandall, *Tetrahedron Lett.* **1974**, 1213.
- ⁶ C. Müller, A. Schweig, and H. Vermeer, *J. Amer. Chem. Soc.* **97**, 982 [1975].
- ⁷ H. Schmidt, A. Schweig, F. Mathey, and G. Müller, *Tetrahedron* **31**, 1287 [1975].
- ⁸ W. Schäfer, A. Schweig, and F. Mathey, *J. Amer. Chem. Soc.*, in press.
- ⁹ W. Schäfer, A. Schweig, K. Dimroth, and H. Kanter, *J. Amer. Chem. Soc.*, in press.
- ¹⁰ N. C. Baird, *Theoret. Chim. Acta* **16**, 239 [1970].
- ¹¹ H. L. Hase, C. Müller, and A. Schweig, in preparation.
- ¹² E. Heilbronner and A. Schmelzer, *Helv. Chim. Acta* **58**, 936 [1975].
- ¹³ R. Hoffmann, *Accounts Chem. Res.* **4**, 1 [1971].
- ¹⁴ R. Gleiter, *Angew. Chem.* **86**, 770 [1974].
- ¹⁵ P. Bishof, J. A. Hashmall, E. Heilbronner, and V. Hornung, *Helv. Chim. Acta* **52**, 1745 [1969].
- ¹⁶ R. Hoffmann, E. Heilbronner, and R. Gleiter, *J. Amer. Chem. Soc.* **92**, 706 [1970].
- ¹⁷ E. Heilbronner, *Isr. J. Chem.* **10**, 143 [1972].
- ¹⁸ H. Oberhammer and S. H. Bauer, *J. Amer. Chem. Soc.* **91**, 12 [1969].
- ¹⁹ J. Laane and R. C. Lord, *J. Mol. Spectrosc.* **39**, 340 [1971].
- ²⁰ O. Ermer and S. Lifson, *J. Amer. Chem. Soc.* **95**, 4121 [1973].
- ²¹ G. Ahlgren, B. Åkermarck, and J. E. Bäckvall, *Tetrahedron Lett.* **1975**, 3501.
- ²² H. Schmidt and A. Schweig, *Chem. Ber.* **107**, 725 [1974].
- ²³ H. Schmidt and A. Schweig, *Tetrahedron Lett.* **1973**, 1437.