


Oxygen Isotope Effect on ^{55}Mn Nuclear Magnetic Shielding in Permanganate

A. R. Haase, O. Lutz, M. Müller, and A. Nolle

Physikalisches Institut der Universität Tübingen, Germany

(Z. Naturforsch. **31a**, 1427–1428 [1976];
received October 5, 1976)

By Fourier transform NMR spectroscopy the ^{55}Mn resonance lines of the different permanganate species



have been resolved in aqueous solutions of potassium permanganate. An isotope effect on the Larmor frequency of ^{55}Mn of (0.599 ± 0.015) ppm to lower frequency was found for the substitution of an ^{16}O atom by an ^{18}O atom in the permanganate ion. An oxygen exchange rate in the permanganate-water system is given.

Introduction

Isotopic substitution is the minimal modification of a molecule. This may result in a change of the nuclear magnetic shielding. The study of such an effect is therefore important for elucidating the origin of chemical shift.

Intramolecular isotopic effects on the Larmor-frequency induced by the replacement of protons by deuterons are well known for many nuclei^{1–4}. The order of magnitude of this effect comes up to 1 ppm in some molecules. An isotope effect induced by heavier nuclei has been found as far as we know only in a few favourable cases in the NMR spectra of ^{13}C (Ref. 5), ^{19}F (Ref. 6–8) and ^{59}Co (Ref. 9, 10). A further example is the ^{55}Mn nucleus in the permanganate ion since the linewidth in this ion is small¹¹ and further large chemical shifts have been found for manganese compounds¹².

Experimental

^{55}Mn FT NMR spectra have been measured at 22.31 MHz with a multinuclei Bruker spectrometer¹³ at (299 ± 2) K. A rotating spherical sample of 10 mm diameter was used. Potassium permanganate of natural abundance (^{55}Mn : 100%, ^{16}O : 99.76%, ^{17}O : 0.037%, ^{18}O : 0.204%) was dissolved in H_2^{18}O (99% enriched in ^{18}O). The concentration was 0.27 molal.

The ^{55}Mn NMR spectra have been measured as a function of time after preparation of the sample since the exchange between the water oxygen atoms and the oxygen atoms in the permanganate is very slow at room temperature.

Reprint requests to Prof. Dr. O. Lutz, Physikalisches Institut der Universität Tübingen, Morgenstelle, D-7400 Tübingen.

Results

In Fig. 1 three of the ^{55}Mn spectra of the mentioned sample at different times after preparation are given. Due to the slow exchange of the oxygen atoms the five permanganate species





Fig. 1. ^{55}Mn spectra of a 0.27 molal solution of natural abundant KMnO_4 in H_2^{18}O (99% enriched in ^{18}O) as a function of time: a) 4 hours, b) 5 days, c) 34 days after preparation. The frequency increases from the left to the right. Experimental parameters: Larmor frequency 22.311341 MHz; experimental spectrum width: 1 kHz; plotted spectrum width: 222 Hz; pulse repetition frequency: 1.67 Hz; number of pulses: 100; measuring time: 1 min; rotating spherical sample with 10 mm outer diameter; 1 K data points were accumulated followed by 7 K of zero-filling before Fourier transformation.

have highly different abundances as a function of time. Shortly after preparation only the $^{55}\text{Mn}^{16}\text{O}_4^-$ ion of the solute is present in the solution and as a result only the usual narrow single line¹¹ is observable. The substitution of one ^{16}O -atom by an ^{18}O -atom gives a further ^{55}Mn resonance line the intensity of which is proportional to the occurrence



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

of the species $^{55}\text{Mn}^{16}\text{O}_3^{18}\text{O}^-$. With running time all five possible species are present in the solution with characteristic abundances which can be calculated assuming a time constant T for the exchange process. At the end of the oxygen substitution the $^{55}\text{Mn}^{18}\text{O}_4^-$ ion is present with overwhelming concentration. At a time of $20 \cdot T$ about 97% of the permanganate ions are in this species with $n = 4$.

All five lines which are due to the five species are well resolved in the spectra at different times (see Figure 1). The substitution of an ^{16}O atom by an ^{18}O atom ($n = 0$ to $n = 1$) results in a change of the Larmor frequency of (12.48 ± 0.33) Hz to lower frequency, which is an isotopic effect of $-(0.559 \pm 0.015)$ ppm. The evaluation of the spectra yields an isotopic effect of $-(2.245 \pm 0.032)$ ppm for the two species $^{55}\text{Mn}^{16}\text{O}_4^-$ and $^{55}\text{Mn}^{18}\text{O}_4^-$, this is $-(0.281 \pm 0.004)$ ppm per mass unit. It should be noted, that the lines are equidistant within the limits of error.

For comparison, in the ^{59}Co -spectrum of the $\text{Co}(\text{CN})_6^{3-}$ -ion an isotopic effect of (0.914 ± 0.004)

ppm for the isotopic pair ^{12}C , ^{13}C and of (0.197 ± 0.007) ppm for the isotopic pair ^{14}N , ^{15}N has been found⁹, which are of the same order of magnitude and also to lower frequency¹⁰.

The evaluation of the intensities of the five lines as a function of time yields the time constant for the oxygen exchange in the water-permanganate system. A preliminary value for the time constant in the given sample is $T \approx 7$ days at room temperature.

Using these values, the very complicated ^{55}Mn pattern of a ^{17}O enriched permanganate sample, which contains also ^{18}O and ^{16}O , used in Ref. ¹³ for evaluating the scalar spin-spin coupling between ^{55}Mn and ^{17}O can be explained. A detailed investigation of these isotopic effect phenomena is running.

Acknowledgement

We are indebted to Prof. Dr. H. Krüger for his continuous support of this work and to W. Nepple for the preliminary measurement of an ^{17}O enriched permanganate sample. We thank the Deutsche Forschungsgemeinschaft for financial support.

¹ H. Batiz-Hernandez and R. A. Bernheim, Progress in Nuclear Magnetic Resonance Spectroscopy (J. W. Emsley, J. Feeney, and L. H. Sutcliffe, Eds.), Vol. 3, p. 63 [1967], Oxford and references therein.

² M. Sphor and A. Loewenstein, Mol. Phys. 15, 9 [1968].

³ Yu. K. Grishin, N. M. Sergeyev, and Yu. A. Ustynyuk, Mol. Phys. 22, 711 [1971] and references therein.

⁴ A. A. Borisenko, N. M. Sergeyev, and Yu. A. Ustynyuk, Mol. Phys. 22, 715 [1971].

⁵ S. Aa. Linde and H. J. Jakobsen, J. Mag. Res. 17, 411 [1975].

⁶ R. J. Gillespie and J. W. Quail, J. Chem. Phys. 39, 2555 [1963].

⁷ S. Rodmar, B. Rodmar, M. K. Sharma, S. Gronowitz, H. Christiansen, and U. Rosén, Acta Chem. Scand. 22, 907 [1968].

⁸ T. Birchall, S. L. Crossley, and R. J. Gillespie, J. Chem. Phys. 41, 2760 [1964].

⁹ P. C. Lauterbur, J. Chem. Phys. 42, 799 [1965].

¹⁰ A. Loewenstein and M. Shporer, Mol. Phys. 9, 293 [1965].

¹¹ O. Lutz and W. Steinkilberg, Z. Naturforsch. 29a, 1467 [1974].

¹² F. Calderazzo, E. A. C. Lucken, and D. F. Williams, J. Chem. Soc. (A) 1967, p. 154.

¹³ O. Lutz, W. Nepple, and A. Nolle, Z. Naturforsch. 31a, 1046 [1976].