

# Soret Coefficient and Thermoelectric Power of Thermocells Containing Molten Binary Mixtures

R. Haase

Lehrstuhl für Physikalische Chemie II der Rheinisch-Westfälischen Technischen Hochschule Aachen

(Z. Naturforsch. 31a, 1731–1732 [1976];  
received October 10, 1976)

For an ionic melt consisting of two components and of three ion constituents, we derive the interrelation between the Soret coefficient and the initial and final values of the thermoelectric power of a thermocell containing the melt considered. We only use the general formulas of Thermodynamics of Irreversible Processes and there is no restriction with respect to the valency types of the electrolytes.

For a solution of a single electrolyte in a neutral solvent, the interrelation between the Soret coefficient and the thermoelectric power of a thermocell has been established long ago<sup>1</sup>. For a molten binary mixture, however, such a relation has not yet been derived except in the special case of uni-univalent electrolytes<sup>2</sup>. We will, therefore, develop the general formula<sup>3</sup> for two-component ionic melts with three ion constituents.

The melt consisting of the two components 1 and 2 contains the following ion constituents:  $\alpha$  (occurring in component 1 only),  $\beta$  (occurring in component 2 only), and  $\gamma$  (common to both components). For transport quantities such as the heats of transport  $^*Q_\alpha$  and  $^*Q_\beta$  and the transported entropies  $^*S_\alpha$  and  $^*S_\beta$  the reference substance is the ion constituent  $\gamma$ .

The state of the melt is described by the thermodynamic temperature  $T$ , the pressure  $P$ , and the mole fraction  $x$  of component 2. The Faraday constant is denoted by  $F$ . The chemical potentials of the two components are denoted by  $\mu_1$  and  $\mu_2$ , those of the three ion constituents by  $\mu_\alpha$ ,  $\mu_\beta$ , and  $\mu_\gamma$ . The symbol  $(\text{grad } \mu_i)_{T,P}$  refers to that part of the gradient of the chemical potential of substance  $i$  which is due to a gradient of  $x$  for uniform values of  $T$  and  $P$ .

Let us consider a melt in which there are gradients of temperature and composition, the pressure being uniform. Then both thermal diffusion (Soret effect) and (ordinary) diffusion (interdiffusion) occur. Eventually a stationary (non-equilibrium) state or steady state will be established where the transport of matter is zero. The general equations governing this steady state in our case (steady Soret

effect in ionic melts with three ion constituents) can be shown to be<sup>5</sup>

$$z_\alpha F \text{ grad } \psi + (\text{grad } \mu_\alpha)_{T,P} + (^*Q_\alpha/T) \text{ grad } T = 0, \quad (1)$$

$$z_\beta F \text{ grad } \psi + (\text{grad } \mu_\beta)_{T,P} + (^*Q_\beta/T) \text{ grad } T = 0 \quad (2)$$

where  $z_i$  is the charge number of the ion constituent  $i$  (positive or negative for cation or anion constituents, respectively) and  $\psi$  is the (inner) electric potential.

We have the relations

$$z_\alpha \nu_\alpha + z_\gamma \nu_\gamma = 0, \quad z_\beta \nu_\beta + z_\gamma \nu_\gamma' = 0, \quad (3)$$

$$\mu_1 = \nu_\alpha \mu_\alpha + \nu_\gamma \mu_\gamma, \quad \mu_2 = \nu_\beta \mu_\beta + \nu_\gamma' \mu_\gamma, \quad (4)$$

$$(1-x) (\text{grad } \mu_1)_{T,P} + x (\text{grad } \mu_2)_{T,P} = 0. \quad (5)$$

Here  $\nu_i$  denotes the dissociation number of the ion constituent  $i$ , the two values for  $\gamma$  ( $\nu_\gamma$  and  $\nu_\gamma'$ ) relating to the two components 1 and 2, respectively.

Combining Eqs. (3) and (4), we find:

$$\mu_\alpha/z_\alpha - \mu_\beta/z_\beta = \mu_1/(z_\alpha \nu_\alpha) - \mu_2/(z_\beta \nu_\beta). \quad (6)$$

We obtain from (5) and (6):

$$(1/z_\alpha) (\text{grad } \mu_\alpha)_{T,P} - (1/z_\beta) (\text{grad } \mu_\beta)_{T,P} = - [\zeta/(1-x)] (\text{grad } \mu_2)_{T,P} \quad (7)$$

with [see Eq. (3)]

$$\zeta \equiv \frac{z_\alpha \nu_\alpha (1-x) + z_\beta \nu_\beta x}{z_\alpha \nu_\alpha z_\beta \nu_\beta} = - \frac{\nu_\gamma' x + \nu_\gamma (1-x)}{z_\gamma \nu_\gamma \nu_\gamma'}. \quad (8)$$

For  $\nu_\gamma' = \nu_\gamma$  the dimensionless quantity  $\zeta$  becomes

$$\zeta = -1/(z_\gamma \nu_\gamma) \quad (9)$$

which for  $z_\gamma = \pm 1$ ,  $\nu_\gamma = 1$  reduces to

$$\zeta = \mp 1. \quad (10)$$

The last relation implies that the two electrolytes (components 1 and 2) are uni-univalent (example: the molten salt mixture  $\text{KNO}_3 + \text{AgNO}_3$  where  $\zeta = 1$ ).

Dividing Eqs. (1) and (2) by  $z_\alpha$  and  $z_\beta$ , respectively, subtracting the resulting equations, and taking account of (7), we find:

$$\zeta \mu_{22} \text{ grad } x = (^*Q/T) (1-x) \text{ grad } T \quad (\text{steady state}) \quad (11)$$

where the relation

$$(\text{grad } \mu_2)_{T,P} = (\partial \mu_2 / \partial x)_{T,P} \text{ grad } x \quad (12)$$

and the abbreviations

$$\mu_{22} \equiv (\partial \mu_2 / \partial x)_{T,P} \quad (13)$$

$$^*Q \equiv ^*Q_\alpha/z_\alpha - ^*Q_\beta/z_\beta \quad (14)$$

Reprint requests to Prof. Dr. R. Haase, Lehrstuhl für Physikalische Chemie der Technischen Hochschule Aachen, Templergraben 59, D-5100 Aachen.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

have been used. Obviously  ${}^*Q$  is a linear combination of the two heats of transport.

We introduce the Soret coefficient  $\sigma$  by

$$\text{grad } x = \sigma x (1-x) \text{ grad } T \text{ (steady state)} . \quad (15)$$

This definition is analogous to that used for binary nonelectrolyte solutions and for solutions of a single electrolyte in a neutral solvent.

It follows from (11) and (15) that

$${}^*Q = \sigma T x \zeta \mu_{22} . \quad (16)$$

Thus the quantity  ${}^*Q$  can be derived from measured values of  $\sigma$ .

The transported entropies  ${}^*S_\alpha$  and  ${}^*S_\beta$  and the heats of transport  ${}^*Q_\alpha$  and  ${}^*Q_\beta$  of the ion constituents  $\alpha$  and  $\beta$  are interrelated by

$${}^*S_\alpha = {}^*Q_\alpha/T + S_\alpha, \quad {}^*S_\beta = {}^*Q_\beta/T + S_\beta \quad (17)$$

where  $S_\alpha$  and  $S_\beta$  are the partial molar entropies of the ion constituents  $\alpha$  and  $\beta$ . The partial molar entropies  $S_1$  and  $S_2$  of the components 1 and 2 are given by relations analogous to the Equations (4). We thus derive in view of (3):

$$S_\alpha/z_\alpha - S_\beta/z_\beta = S_1/(z_\alpha \nu_\alpha) - S_2/(z_\beta \nu_\beta) , \quad (18)$$

an equation corresponding to (6). Combining (14), (17), and (18), we obtain:

$${}^*S_\alpha/z_\alpha - {}^*S_\beta/z_\beta = {}^*Q/T + S_1/(z_\alpha \nu_\alpha) - S_2/(z_\beta \nu_\beta) , \quad (19)$$

another useful relation.

We now consider a thermocell containing the melt and two chemically identical electrodes reversible<sup>6</sup> to the ion constituent  $\beta$  (e.g. two silver electrodes dipping into the molten salt mixture  $\text{KNO}_3 + \text{AgNO}_3$ ). Then the expression for the difference between the final (steady-state) value  $\varepsilon_\infty$  and the initial value  $\varepsilon_0$  of the thermoelectric power of the thermocell can be shown to be<sup>7</sup>

$$F(\varepsilon_\infty - \varepsilon_0) = t_\alpha {}^*Q/T = (1 - t_\beta) {}^*Q/T \quad (20)$$

where Eq. (19) has been used and  $t_\alpha$  or  $t_\beta$  denotes the (internal) transport number of the ion constituent  $\alpha$  or  $\beta$ , respectively, the ion constituent  $\gamma$  being again the reference substance.

We derive our final formula from (16) and (20):

$$F(\varepsilon_\infty - \varepsilon_0) = \sigma t_\alpha \zeta x \mu_{22} = \sigma (1 - t_\beta) \zeta x \mu_{22} . \quad (21)$$

Thus there is a general relation among the three measurable quantities  $\sigma$ ,  $\varepsilon_0$ , and  $\varepsilon_\infty$  referring to non-isothermal transport processes. The quantity  $\zeta$  may be replaced by  $(1-x)/(z_\beta \nu_\beta X_1)$  where  $X_1$  denotes the equivalent fraction of component 1.

Since  $\mu_{22} > 0$  (stability condition) and  $\zeta > 0$  for  $z_\gamma < 0$  or  $\zeta < 0$  for  $z_\gamma > 0$  [see Eq. (8)], it follows from (21) that  $\varepsilon_\infty - \varepsilon_0$  and  $\sigma$  have the same sign if  $\gamma$  is an anion constituent while  $\varepsilon_\infty - \varepsilon_0$  and  $\sigma$  have opposite signs if  $\gamma$  is a cation constituent.

For  $\nu'_\gamma = \nu_\gamma$  Eq. (21), when combined with (3) and (9), reduces to the relation

$$F(\varepsilon_\infty - \varepsilon_0) = \sigma [(1 - t_\beta)/(z_\beta \nu_\beta)] x \mu_{22} . \quad (22)$$

We may compare Eq. (22) to the corresponding formula<sup>8</sup> valid for a solution of a single electrolyte (with two ion constituents) in a neutral solvent:

$$F(\varepsilon_\infty - \varepsilon_0) = \sigma [(1 - t_i)/(z_i \nu_i)] m (\partial \mu_2 / \partial m)_{T,P} . \quad (23)$$

Here  $t_i$ ,  $z_i$ , and  $\nu_i$  are the (Hittorf) transport number, the charge number, and the dissociation number of that ion constituent (i) for which the electrodes are reversible and  $m$  and  $\mu_2$  denote the molality and the chemical potential of the electrolyte.

For any binary ionic melt with three ion constituents, the transported entropies  ${}^*S_\alpha$  and  ${}^*S_\beta$  can be derived from measurements of either  $\varepsilon_0$  and  $\varepsilon_\infty$  or  $\varepsilon_0$  and  $\sigma$  or  $\varepsilon_\infty$  and  $\sigma$ . This follows from Eqs. (16) and (19) and from relations given earlier<sup>7</sup>. So in this respect, too, the two-component melts resemble electrolyte solutions.

<sup>1</sup> R. Haase, Trans. Faraday Soc. **49**, 724 [1953].

<sup>2</sup> U. Prüser, Dissertation Technische Hochschule Aachen, 1976.

<sup>3</sup> Though the general equations given by Schönert and Sinistri<sup>4</sup> implicitly contain the desired formula, we will give a new derivation since we do not want to change the reference system, as do the authors mentioned.

<sup>4</sup> H. Schönert and C. Sinistri, Z. Elektrochem. **66**, 413 [1962].

<sup>5</sup> R. Haase, Thermodynamics of Irreversible Processes, Reading (Mass.) 1969, p. 330, Equation [4-24.24].

<sup>6</sup> The question of whether  $\alpha$  or  $\beta$  is taken to be the "reversible ion"  $r$  is only a matter of choice of the com-

ponents. The case, however, where the electrodes are reversible to the common ion constituent  $\gamma$  ( $r=\gamma$ ) is really different. Here we find that  $t_\alpha$  in (20) and (21) has to be replaced by  $t_\alpha - X_1$ . The formula for the thermoelectric power  $\varepsilon$  at any instant (for any value of the mole fraction difference  $\Delta x$  corresponding to the temperature difference  $\Delta T$  in the melt) results from (21), or from the equation replacing it for  $r=\gamma$ , if one substitutes  $\varepsilon$  for  $\varepsilon_\infty$  and  $\lim_{\Delta T \rightarrow 0} \Delta x / \Delta T$  for  $\sigma x(1-x)$ .

<sup>7</sup> R. Haase and J. Richter, Z. Naturforsch. **24a**, 418 [1969], Eqs. (1), (5) and (6).

<sup>8</sup> R. Haase<sup>5</sup>, p. 391 to 393, Eqs. [4-29.27], [4-29.33], and [4-29.37].