

# Electron Diffraction Investigation of the Molecular Structure of Trifluoromethyl Sulphonyl Chloride

Jon Brunvoll

Department of Physical Chemistry, The University of Trondheim

István Hargittai and Mária Kolonits

Central Research Institute of Chemistry, Hungarian Academy of Sciences

Z. Naturforsch. **33a**, 1236–1238 (1978);  
received May 30, 1978

An electron diffraction investigation yielded the following geometric parameters for trifluoromethyl sulphonyl chloride  $r_g(\text{C}-\text{F})$  1.326(4),  $r_g(\text{S}=\text{O})$  1.416(7),  $r_g(\text{S}-\text{Cl})$  2.016(5),  $r_g(\text{S}-\text{C})$  1.857(6) Å,  $\angle \text{O}=\text{S}-\text{C}$  108.3(7)°,  $\angle \text{C}-\text{S}-\text{Cl}$  98.7(4)°,  $\angle \text{O}=\text{S}=\text{O}$  122.4(10)°,  $\angle \text{O}=\text{S}-\text{Cl}$  108.3(3)°,  $\angle \text{F}-\text{C}-\text{F}$  109.9(3)°. The S—C bond is in  $\text{CF}_3\text{SO}_2\text{Cl}$  by almost 0.1 Å longer than in  $\text{CH}_3\text{SO}_2\text{Cl}$ .

Recently we determined the molecular structure of trichloromethyl sulphonyl chloride [1] as an extension of our studies on simple sulphone molecules [2]. The sulphur bond configuration in  $\text{CCl}_3\text{SO}_2\text{Cl}$  is consistent with the characteristic structural variations observed in the sulphone series. There was also indication of a particularly long C—S bond,  $1.87 \pm 0.03$  Å, as compared with that of methane sulphonyl chloride ( $1.763 \pm 0.005$  Å [3]). This result, however, suffered from very large uncertainty. Trifluoromethyl sulphonyl chloride is more suitable for an electron diffraction determination of the S—C distance. We found it also interesting to examine the influence of the  $\text{CF}_3$  group on the sulphonyl chloride structure and to compare the barriers to internal rotation in the  $\text{CF}_3$  and  $\text{CCl}_3$  derivatives.

The electron diffraction patterns were taken at a nozzle temperature of about 10 °C with the Budapest EG-100 apparatus [4–6]. For references on data treatment and scattering functions, see [3]. The reduced molecular intensities and radial distributions are shown in Figs. 1 and 2.

The experimental radial distribution gave evidence for the staggered conformation, similarly to that of the trichloromethyl derivative [1].  $C_s$  symmetry was assumed for the molecule as a whole. The  $\text{CF}_3$  group was assumed to have  $C_3$  symmetry

Reprint requests to Dr. I. Hargittai, Central Research Institute of Chemistry, Hungarian Academy of Sciences, H-1525 Budapest, P.O. Box 17, Hungary.




Fig. 1. Experimental (E) and theoretical (T) molecular intensities and their differences ( $\Delta$ ). The theoretical distribution was calculated from the parameters of Table 1.





Fig. 2. Experimental (E) and theoretical (T) radial distributions.

with the S—C bond coinciding with the symmetry axis. The numbering of atoms is given as follows:



Mean amplitudes of vibration ( $l$  values) and also perpendicular vibration corrections ( $K$  values) were calculated from an approximate force field compiled and adjusted to the frequencies given by Lindner and Weber [7]. The calculated  $l$  and  $K$  values are listed in Table 1.

Since some geometric and vibrational parameters were strongly correlated, fixed differences corresponding to the spectroscopic calculations were applied for several pairs of  $l$  values in the least-squares refinement. Of the important geometric parameters,  $r(\text{S}=\text{O})$  showed some sensitivity to the assumption for the difference between  $l(\text{S}=\text{O})$  and  $l(\text{C}-\text{F})$ . This was taken into account in the error estimates. For the other parameters

$$\sigma_t = [2\sigma^2 + (0.002r)^2]^{1/2}$$

(for distances and angles) and

$$\sigma_t = [2\sigma^2 + (0.02)^2]^{1/2}$$

(for the  $l$  values) were used, where  $\sigma$  is the standard deviation from the least-squares refinement.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

Table 1. Structural parameters of trifluoromethyl sulphonyl chloride.

	$r_a$ (Å), $\angle$ (°)	$\sigma_t$	$l^{ED}$ (Å)	$\sigma_t$	$l^{SP}$ (Å)	$K^{SP}$ (Å)
Independent parameters						
C—F	1.324	0.004	0.052	0.005	0.049	0.0149
S=O	1.415	0.007	0.042	0.003	0.035	0.0066
S—Cl	2.015	0.005	0.045	0.002	0.049	0.0061
S—C	1.856	0.006	0.046	0.006	0.052	0.0013
O ... O	2.480	0.014	0.066	0.018	0.066	0.0097
Cl ... F4 ( <i>anti</i> )	4.153	0.012	0.080	0.009	0.070	0.0010
C—S=O	108.3	0.7				
S—C—F	109.1	0.4				
Dependent parameters						
Cl ... O	[2.802]		0.070	0.004 <sup>a</sup>	0.072	0.0091
Cl ... F5 ( <i>gauche</i> )	[3.125]		0.173	0.030 <sup>b</sup>	0.175	0.0079
Cl ... C	[2.939]		0.067	<sup>a</sup>	0.069	0.0035
F4 ... O2	[3.089]		0.138	<sup>b</sup>	0.140	0.0107
F5 ... O2	[3.765]		0.069	0.034	0.070	0.0021
F6 ... O2	[2.966]		0.140	<sup>b</sup>	0.143	0.0096
O ... C	[2.664]		0.070	0.016 <sup>c</sup>	0.070	0.0042
F ... F	[2.169]		0.055	0.003	0.059	0.0240
F ... S	[2.608]		0.068	<sup>c</sup>	0.068	0.0072
Cl—S—C	[98.7]	[0.4]				
O=S=O	[122.4]	[1.0]				
Cl—S=O	[108.3]	[0.3]				
F—C—F	[109.9]	[0.3]				

<sup>a</sup>, <sup>b</sup>, <sup>c</sup> mean amplitudes refined in groups.

The long S—C bond is the only striking feature of the molecular geometry of trifluoromethyl sulphonyl chloride as compared with that of methane sulphonyl chloride [3, 8]. The difference in  $r$ (S—C) is larger than would have been expected on the basis of known structural changes by CH<sub>3</sub>/CF<sub>3</sub> substitution [9].

The other geometric parameters of the sulphur bond configuration in trifluoromethyl sulphonyl chloride are consistent with the structural variations in the sulphone series [2]. The S=O bond length agrees within the error limits with the estimates from the correlation relationship between bond lengths and bond stretching frequencies [10]. The C—F bond seems to be of normal length for this compound (cf. the compilation of Yokozeki and Bauer [9]).

The barrier to internal rotation in CF<sub>3</sub>SO<sub>2</sub>Cl was estimated to be between 2.2 and 5.0 kcal mol<sup>-1</sup> (9 and 21 kJ mol<sup>-1</sup>) using Karle's method [11] as shown in Figure 3. These values can be compared




Fig. 3. Estimation of the barrier to internal rotation around the S—C bond according to Ref. [11]. The solid lines correspond to the experimental  $l$ -values and the broken lines to their error limits. The straight vertical lines indicate the calculated framework contributions.

with the interval of 3.9 to 5.6 kcal mol<sup>-1</sup> estimated for CCl<sub>3</sub>SO<sub>2</sub>Cl [1]. The barrier to internal rotation in CH<sub>3</sub>SO<sub>2</sub>Cl was reported to be at least 2 kcal mol<sup>-1</sup> [8].

[1] J. Brunvoll, I. Hargittai and R. Seip, Z. Naturforsch. **33a**, 1178 (1978).

[2] I. Hargittai, Sulphone Molecular Structures, Lecture Notes in Chemistry, Vol. 6, Springer-Verlag, Berlin 1978.

- [3] M. Hargittai and I. Hargittai, *J. Chem. Phys.* **59**, 2513 (1973).
- [4] I. Hargittai, J. Hernádi, M. Kolonits, and Gy. Schultz, *Rev. Sci. Instrum.* **42**, 546 (1971).
- [5] I. Hargittai, J. Hernádi, and M. Kolonits, *Prib. Tekh. Eksp.* 239 (1972).
- [6] J. Tremmel, M. Kolonits, and I. Hargittai, *J. Phys. E. Sci. Instrum.* **10**, 664 (1977).
- [7] E. Lindner and H. Weber, *Z. Naturforsch.* **22b**, 1243 (1967).
- [8] B. P. Van Eijck, A. J. Korthof, and F. C. Mijlhoff, *J. Mol. Struct.* **24**, 222 (1975).
- [9] A. Yokozeki and S. H. Bauer, *Topics in Current Chemistry* **53**, 71, Springer-Verlag, Berlin 1975.
- [10] J. Brunvoll and I. Hargittai, *Spectroscopy Letters* **10**, 815 (1977); *Acta Chim. (Budapest)*, **96**, 337 (1978).
- [11] J. Karle, *J. Chem. Phys.* **45**, 4149 (1966).