

# The Electrical Conductivity of Molten Lead Iodide

A. J. Easteal and P. H. Khoo  
Chemistry Department, University of Auckland,  
New Zealand

Z. Naturforsch. **35a**, 1415–1416 (1980);  
received November 4, 1980

The electrical conductivity of molten  $\text{PbI}_2$  has been determined in the temperature range 798–978 K. The data have been fitted to the equation  
$$\kappa = -0.8167 + 1.5668 \times 10^{-3} T + 3.810 \times 10^{-7} T^2.$$
The data are compared with previously published data for  $\text{PbI}_2$ .

The electrical conductivity of molten lead iodide has recently been critically re-examined by Janz and co-workers [1], with the result that the data of Bogacz and Zuca [2] have been recommended as reference data for the salt. The data base which was selected for the conductivity of  $\text{PbI}_2$  supersedes the previous recommendation [3] which was based on the data published by Karl and Klemm [4]. The purpose of this communication is to present new data for lead iodide, which are in substantially better agreement with the older data [4] than with the currently recommended data, and hence cast some doubt on the reliability of the recommended data.

The  $\text{PbI}_2$  was British Drug Houses Laboratory Reagent grade material, used without additional purification. The conductivity cell\* and associated equipment, and the experimental procedure were essentially as described previously [5], [6]. The overall precision of values of conductivity is estimated to be within  $\pm 0.5\%$ . Temperatures were measured with a calibrated Pt/13% Rh-Pt thermocouple, with estimated accuracy  $\pm 0.5$  K.

The conductivity ( $\kappa$ ) of  $\text{PbI}_2$  was measured in the temperature range 798–978 K. The temperature dependence of  $\kappa$  is described with good precision (r.m.s. deviation of experimental points from the calculated curve 0.005) by the equation

$$\kappa = -0.8167 + 1.5668 \times 10^{-3} T + 3.810 \times 10^{-7} T^2, \tag{1}$$

Reprint requests to Dr. A. J. Easteal, Chemistry Department, University of Auckland, Private Bag, Auckland, New Zealand.

\* An all-quartz construction with platinum electrodes; melt protected by nitrogen.

where the units of  $T$  and  $\kappa$  are K and  $\text{S cm}^{-1}$ . The variation of  $\kappa$  with temperature is shown graphically in Fig. 1, together with previously published data. Our plotted data are values interpolated at 20 K intervals using Eq. (1) plus the values corresponding to the extremes of the experimental temperature range. The raw data of Karl and Klemm are shown, and the data due to Bogacz and Zuca in Fig. 1 are the values tabulated by Janz and co-workers [1].

It is clear from Fig. 1 that the present data and those due to Karl and Klemm show fairly good agreement, particularly at higher temperatures.




Fig. 1. Electrical conductivity of molten lead iodide.  $\Delta$ : Ref. [1];  $\square$ : Ref. [4];  $\bullet$ : this work.

| $T/\text{K}$ | $\kappa/\text{S cm}^{-1}$ | data base | $\delta\kappa$ (%) | Table 1. Comparison of electrical conductivity data for molten lead iodide at selected temperatures.<br><br>a: Ref. [2], cited in Ref. [1];<br>b: Ref. [4];<br>c: this work. |
|--------------|---------------------------|-----------|--------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 723          | 0.600                     | a         | 16.5               |                                                                                                                                                                              |
|              | 0.489                     | b         | 5.0                |                                                                                                                                                                              |
|              | 0.515                     | c         | —                  |                                                                                                                                                                              |
| 863          | 0.903                     | a         | 10.3               |                                                                                                                                                                              |
|              | 0.805                     | b         | 1.7                |                                                                                                                                                                              |
|              | 0.819                     | c         | —                  |                                                                                                                                                                              |
| 973          | 1.108                     | a         | 3.7                |                                                                                                                                                                              |
|              | 1.069                     | c         | —                  |                                                                                                                                                                              |

0340-4811 / 80 / 1200-1415 \$ 01.00/0. — Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

Both data sets deviate considerably from the values of Bogacz and Zuca. A quantitative comparison of the three sets of data is given in Table 1 below. To make comparison, Karl and Klemm's data have been fitted by the equation

$$\alpha = -0.6501 + 1.0054 \times 10^{-3} T + 7.888 \times 10^{-7} T^2. \quad (2)$$

The deviations ( $\delta\alpha$ ) listed in Table 1 are expressed as percentages relative to our values at the tem-

peratures specified. The discrepancies between our data and those of Karl and Klemm are greater than can be accounted for by the probable experimental uncertainties of the measurements and may be due largely to use of  $\text{PbI}_2$  from different sources. The much greater discrepancies between the data of Bogacz and Zuca and the other two sets are, however, too large to have arisen from differing sources of material, and no explanation for the discrepancies is apparent.

- [1] G. J. Janz, R. P. T. Tomkins, C. B. Allen, J. R. Downey, and S. K. Singer, *J. Phys. Chem. Ref. Data* **6**, 409 (1977).
- [2] A. Bogacz and S. Zuca, *Rev. Roum. Chim.* **11**, 183 (1966).
- [3] G. J. Janz, F. W. Dampier, G. R. Lakshminarayanan, G. R. Lorenz, and R. P. T. Tomkins, *Nat. Stand. Ref. Data Ser., N.B.S. (U.S.)* **15**, 84 (1968).
- [4] W. Karl and A. Klemm, *Z. Naturforsch.* **19a**, 1619 (1964).
- [5] A. J. Easteal and I. M. Hodge, *J. Phys. Chem.* **74**, 730 (1970).
- [6] A. J. Easteal and P. H. Khoo, *Aust. J. Chem.* **24**, 1581 (1971).