

Lifetime Measurements of All Excited States of the 4d⁹5p Configuration in the Pd-I Spectrum

M. Baumann, H. Liening, and H. Loos
Physikalisches Institut der Universität Tübingen

Z. Naturforsch. **36a**, 778–779 (1981);
received May 19, 1981

The lifetimes for the 4d⁹5p configuration have been determined by zero field level crossing technique. The results are (in nsec):

$\tau(^3P_1^0) = 7.5$ (4)	$\tau(^1D_2^0) = 7.2$ (5)
$\tau(^3P_2^0) = 6.9$ (8)	$\tau(^3F_2^0) = 9.0$ (6)
$\tau(^1P_1^0) = 5.0$ (4)	$\tau(^3F_3^0) = 7.9$ (5)
$\tau(^3D_1^0) = 4.9$ (4)	$\tau(^3F_4^0) = 7.1$ (5)
$\tau(^3D_2^0) = 8.1$ (5)	$\tau(^1F_3^0) = 8.4$ (6)
$\tau(^3D_3^0) = 7.0$ (5)	

An atomic beam of natural Palladium, produced in an oven of coaxial construction, was irradiated by the light of a hollow cathode lamp [1]. By absorption of the corresponding resonance lines (Fig. 1) the atoms were raised from the ground state 4d¹⁰1S₀ and from the metastable D-states of the configuration 4d⁹5s to the excited states of the 4d⁹5p configuration. (At an oven temperature of about 1850 K the metastable D-states are noticeably populated.)




Fig. 1. Part of the level scheme of the Pd I spectrum (the wavelength are indicated in Table 1).

Reprints requests to Dr. H. Liening, Physikalisches Institut der Universität, Morgenstelle 14, D-7400 Tübingen.

The lifetimes of all excited states except the ³P₀⁰-state were determined by measuring the depolarization of the fluorescence light in a magnetic field (Hanle-effect) [2]. The ³P₀⁰-state gives no Hanle-signal because $J=0$ for the even isotopes; the half width of the Hanle-signal of the odd isotope ¹⁰⁵Pd($I=5/2$) is very large ($g_F=g_I'$), therefore this signal is not detectable.

The $\Delta m=2$ zero field level crossing signals were detected by an experimental arrangement similar to that described in an earlier paper [3]. The spectral lines under study, which partially are close together in wavelength, were selected by a monochromator, hence a considerable decrease of the detectable intensities. A further reduction in the signal-to-noise ratio may be given due to the different possibilities of excitation of the states under study. As may be seen from the Breit-formula [4] the signal amplitudes of the Hanle-effect changes in sign whether the excitation is via a $J \rightarrow J$ transition or a $J \pm 1 \rightarrow J$ transition. Therefore if the excitation is achieved by all corresponding lines in the spectrum of a hollow cathode lamp, the Hanle signal in some special cases nearly cancels. This explains the very poor signal-to-noise ratio of the Hanle effect we measured in some excited states of the 4d⁹5p-configuration. In order to overcome this difficulty signal processing up to ten hours had to be employed.

For the geometry chosen in our experiment the signal curve of the even isotopes of the field dependent part of the observed fluorescence intensity plotted against the magnetic field is lorentzian shaped with a half width ΔB related to the coherence time by $\Delta B=1/\gamma \cdot T$, where γ is the gyromagnetic factor of the excited state. Only for small particle density the coherence time is equal to the lifetime of the excited state [5]. Therefore in all measurements the particle density has been varied over a range of one order of magnitude in order to get rid of the coherence narrowing [5].

But because of the different gyromagnetic factors for the hfs levels of the odd isotope ¹⁰⁵Pd (22%) the resulting signal is a superposition of Lorentzian curves with different half widths. In addition the broad band excitation, which is supposed in the Breit formula is no longer valid as may be seen from relative isotope shifts in the Pd I spectrum

0340-4811 / 81 / 0700-0778 \$ 01.00/0. — Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

Table 1.

Term	$\lambda/\text{\AA}$	$\Delta H_{1/2}$	g_J	τ/nsec without hfs	τ/nsec with hfs	τ/nsec Corliss & Bozman
4d ⁹ 5p ³ P ₁ ⁰	2763	10.9 ± 0.5	1.396	7.5	7.5 ± 0.4	3.8
³ P ₂ ⁰	3635	11.1 ± 1.3	1.482	6.9	6.9 ± 0.8	7.4
¹ P ₁ ⁰	2448	29.7 ± 2.1	0.768	4.9	5.0 ± 0.4	1.9
³ D ₁ ⁰	2476	28.0 ± 2.3	0.831	4.8	4.9 ± 0.4	2.5
³ D ₂ ⁰	3421	14.2 ± 0.8	0.988	7.9	8.1 ± 0.5	6.0
³ D ₃ ⁰	3243	12.8 ± 0.9	1.276	6.9	7.0 ± 0.5	5.3
¹ D ₂ ⁰	3442	14.2 ± 0.9	1.114	7.2	7.2 ± 0.5	2.5
³ F ₂ ⁰	3482	16.8 ± 1.0	0.752	8.8	9.0 ± 0.6	2.9
³ F ₃ ⁰	3610	13.5 ± 0.9	1.063	7.7	7.9 ± 0.5	5.7
³ F ₄ ⁰	3405	12.8 ± 0.9	1.25	7.0	7.1 ± 0.5	7.5
¹ F ₃ ⁰	3553	12.5 ± 0.9	1.081	8.3	8.4 ± 0.6	3.4

recently become available [6]. Taking into account approximately the spectral distribution of the incident light the resulting shapes of the signals were calculated due to the Breit formula and fitted to the measured values with the aid of a mini-computer (PDP 11/45) making use of the experimental g_J -values [7, 8]. The results are given in column 6 of Table 1. The errors include uncertainties made by fitting as well as systematic errors.

If the measured Hanle signals are fitted simply with one Lorentzian curve without regard of the lamp profile we get lifetimes that are at most 2.5% smaller. This seems to be a little surprising in view of the odd-even isotope shift [6] and the large hfs-

splitting in the D-states of the 4d⁹5s configuration [9] and the states of the configuration 4d⁹5p [10]. (Table 1 also includes the results of earlier measurements of lifetimes in the Pd I spectrum [11, 12].) The lifetime of the ³P₁⁰-state has already been measured by Budick [13]. His result was $\tau(^3P_1^0) = 8.7(9)$ nsec. Another measurement was performed by Hese and Weise [14]. In the last column in Table 1 we calculated the τ -values from the transition probabilities of Corliss and Bozman [15]. As can be seen, there is no systematic trend in the deviation.

We like to thank Prof. H. Krüger for his continuous support of this work.

[1] A. D. White, J. Appl. Phys. **30**, 711 (1959).

[2] W. Hanle, Z. Physik **30**, 93 (1924).

[3] M. Baumann, H. Liening, and H. Lindel, Phys. Lett. **68A**, 319 (1978).

[4] P. A. Franken, Phys. Rev. **121**, 508 (1961).

[5] J. P. Barrat, J. Phys. Rad. **20**, 541, 633, 657 (1959).

[6] P. E. G. Baird, Proc. Roy. Soc. London **A351**, 267 (1976).

[7] A. G. Shenstone, Phys. Rev. **36**, 669 (1930).

[8] C. H. Lindsley and N. Rosen, J. Opt. Soc. Amer. **31**, 531 (1941).

[9] K. H. Channappa and J. M. Pendlebury, Proc. Phys. Soc. London **86**, 267 (1965).

[10] A. Steudel, Z. Physik **132**, 429 (1952).

[11] M. Baumann and H. Liening, Phys. Lett. **36A**, 329 (1971).

[12] H. Liening, Z. Physik **266**, 287 (1974).

[13] B. Budick, Phys. Rev. **168**, 89 (1968).

[14] A. Hese and J. Weise, 3rd EGAS-Conference Reading 1971 (unpublished).

[15] C. Corliss and W. Bozman, Experimental Transition Probabilities for Spectral Lines of Seventy Elements, Nat. Bureau of Standards, Monograph **53** (1962).