

## A Second Integral of Motion in a Triaxial Galaxy

N. Caranicolas

Department of Astronomy, University of Thessaloniki, Thessaloniki, Greece

Z. Naturforsch. **39a**, 395–396 (1984);  
received January 5, 1984

We construct a formal second integral of motion in a bisymmetrical potential field describing the motion on the plane of symmetry of a triaxial galaxy. We also check the constancy of the second integral adding higher-order terms which are found using a computer program.

Since the first indication that elliptical galaxies may not be rotationally flattened [1] and the confirmation of this by Illingworth [2] the dynamics of elliptical galaxies have been an active field of research.

In order to describe the motion on the plane of symmetry of a triaxial galaxy we adopt the bisymmetrical potential

$$V = \frac{1}{2} (Px^2 + Qy^2) - \varepsilon (ax^4 + 2bx^2y^2 + cy^4), \quad (1)$$

where  $P, Q, a, b, c$  are positive constants and  $\varepsilon$  is a small parameter. It is also supposed that the ratio  $P^{1/2}/Q^{1/2}$  is irrational. Similar potentials with cubic [3], [4] or quartic [5] terms have been used to describe the motion on the meridian plane of biaxial galaxies.

In the case of the potential (1) the Hamiltonian

$$H = \frac{1}{2} (X^2 + Y^2 + Px^2 + Qy^2) - \varepsilon (ax^4 + 2bx^2y^2 + cy^4) = H_0 + \varepsilon H_1 = h, \quad (2)$$

where  $X = dx/dt$ ,  $Y = dy/dt$  and  $h$  is the numerical value of the function  $H$ , is an exact integral of motion.

In applying the method of Contopoulos [6] we shall try to construct a formal second integral of motion of the form

$$I = I_0 + \varepsilon I_1 + \dots, \quad (3)$$

where

$$I_0 = \frac{1}{2} (X^2 + Px^2) \quad (4)$$

and

$$I_{i+1} = - \int (I_i, H_1) dt + f_{i+1}(I_0, J_0) \quad (5)$$

$$(i = 0, 1, 2, \dots).$$

In (5),  $(I_i, H_1)$  are Poisson brackets,  $J_0 = H_0 - I_0$ ,  $t$  is an auxiliary variable and  $f_{i+1}(I_0, J_0)$  is an arbitrary function. Equation (5) for  $i = 0$  gives

$$I_1 = -4 \int (ax^3X + bx^2y^2X) dt + f_1(I_0, J_0). \quad (6)$$

If we take into account the solutions of the “unperturbed” problem ( $\varepsilon = 0$ )

$$\begin{aligned} x &= (2I_0/P)^{1/2} \sin P^{1/2}(t - t_0), \\ y &= (2J_0/Q)^{1/2} \sin Q^{1/2}t, \\ X &= (2I_0)^{1/2} \cos P^{1/2}(t - t_0), \\ Y &= (2J_0)^{1/2} \cos Q^{1/2}t, \end{aligned} \quad (7)$$

where  $t_0$  is a constant, we find

$$\begin{aligned} I_1 &= \frac{bJ_0}{PQ} (X^2 - Px^2) \\ &\quad - \frac{b}{P-Q} \left[ \frac{1}{2Q} (X^2 - Px^2) (Y^2 - Qy^2) + 2xyXY \right] \\ &\quad - ax^4 + f_1(I_0, J_0), \end{aligned} \quad (8)$$

where after finishing the integration we used the following identities:

$$\begin{aligned} 2I_0 \cos 2P^{1/2}(t - t_0) &= X^2 - Px^2, \\ I_0 \sin 2P^{1/2}(t - t_0) &= P^{1/2}xX, \\ 2J_0 \cos 2Q^{1/2}t &= Y^2 - Qy^2, \\ J_0 \sin 2Q^{1/2}t &= Q^{1/2}yY. \end{aligned} \quad (9)$$

Assuming that  $f_1(I_0, J_0)$  is equal to zero we have the following expression for the first two terms of the second integral:

$$\begin{aligned} I &= I_0 + \varepsilon I_1 = \frac{1}{2} (X^2 + Px^2) \\ &\quad + \varepsilon \left\{ \frac{bJ_0}{PQ} (X^2 - Px^2) - \frac{b}{P-Q} \left[ \frac{1}{2Q} (X^2 - Px^2) \right. \right. \\ &\quad \cdot (Y^2 - Qy^2) + 2xyXY \left. \right] - a_1x^4 \right\}. \end{aligned} \quad (10)$$

The higher order terms of the second integral are given through the recurrent formula (5). However,

Reprint requests to Dr. N. Caranicolas, Department of Astronomy, University of Thessaloniki, Thessaloniki, Greece.

0340-4811 / 84 / 0400-0395 \$ 01.3 0/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

Table 1. Constancy of the second integral.

$I$ (Degree)	max	min	$D$	$2D/(\max + \min)$
$I_0(2)$	0.0100	0.0074	0.0026	0.30
$+ \varepsilon I_1(4)$	0.00862	0.00850	0.00012	0.014
$+ \varepsilon^2 I_2(6)$	0.008611	0.008594	0.000017	0.0020
$+ \varepsilon^3 I_3(8)$	0.0086002	0.0085984	0.0000018	0.00021
$+ \varepsilon^4 I_4(10)$	0.00859982	0.00859967	0.00000015	0.000017
$+ \varepsilon^5 I_5(12)$	0.00859978	0.00859977	0.00000001	0.0000012
$+ \varepsilon^6 I_6(14)$	0.00859977	0.00859977	0.00000000	0.00000000

the actual calculation of these terms becomes soon prohibitively long.

Using a computer program [7], we calculated the higher order terms of the second integral up to the terms of the fourteenth degree in the variables. The number of terms of different degrees are (2) 2; (4) 8; (6) 20; (8) 40; (10) 70; (12) 112; (14) 168. It is interesting to note that as higher order terms are included in the expansion (10) the second integral is better conserved when the energy is small. For this reason the values of the second integral were found for a number of orbits in many points along the orbit. The orbits were calculated by numerical integration of the equations of motion using the Runge-Kutta method in double precision.

The units used were  $kpc$  for  $x, y$  and  $kpc(10^7 yr)^{-1}$  for  $X, Y$ . The step of integration was  $0.02(10^7 yr)$ .

In Table 1 max and min are the maximum and minimum values of the second integral for an orbit with initial conditions  $x_0 = 0.2, y_0 = X_0 = 0$  and  $h = 0.1$ , for a time interval equal to  $5 \times 10^9$  years. The values of the constants are  $P = 0.5, Q = 0.6, a = 1.0, b = 0.8, c = 0.5$  while  $\varepsilon$  is equal to 0.05. The maximum deviation  $D = \max - \min$  and the maximum relative deviation  $2D/(\max + \min)$  are also given in this Table. We see that when the terms of the fourteenth degree are included 8 significant figures of the second integral are constant. The energy constant was conserved up to the twelfth printed figure.

- [1] F. Bertola and M. Cappacioli, *Astrophys. J.* **236**, 439 (1975).
- [2] G. Illingworth, *Astrophys. J. Lett.* **218**, L43 (1977).
- [3] M. Henon and C. Heiles, *Astron. J.* **69**, 73 (1964).
- [4] G. Contopoulos and M. Moutsoulas, *Astron. J.* **70**, 817 (1965).

- [5] A. Saaf, *Astrophys. J.* **154**, 483 (1968).
- [6] G. Contopoulos, *Z. Astrophys.* **49**, 273 (1960).
- [7] A. Giorgilli and L. Galgani, *Cell. Mech.* **17**, 281 (1978).