

A Note on Solutions of the Korteweg-de Vries Equation

K. Murawski

Agricultural Academy, Department of Physics,
Akademicka 13, 20-033 Lublin, Poland

Z. Naturforsch. **40a**, 193–194 (1985);
received October 22, 1984

A new method to construct solutions of the Korteweg-de Vries equation and an example of such a solution are given.

In recent years, a number of interesting papers have appeared on the Korteweg-de Vries (KdV) equation solved by various exact techniques [1–22]. In spite of this, the main purpose of this note is to present a new way of finding solutions of the KdV equation. We write

$$p_t + \beta p p_x + \alpha p_{xxx} = 0, \quad (1)$$

where β and α are nonlinearity and dispersion coefficients, respectively. The subscripts x and t indicate partial differentiations.

An introducing the function w , which has been done for the first time by Wahlquist and Estabrook [1],

$$p \equiv -w_x, \quad (2)$$

and substituting it in (1) we obtain

$$w_t = \frac{\beta}{2} w_x^2 - \alpha w_{xxx} \equiv \frac{\beta}{2} p^2 + \alpha p_{xx}, \quad (3)$$

($F = \text{const}$)

where the partial derivatives of w are chosen to be

$$w_x = g(w^2 - k^2) \quad (4)$$

$$w_t = e(w^2 - f k^2). \quad (5)$$

Keeping $F = 0$ let us transform (4) and (5) into the form

$$w_t - e/g w_x = 0, \quad (7)$$

$$w_x = g(w^2 - k^2). \quad (8)$$

We immediately obtain from (7)

$$w = w(e/g t + x) \equiv w(\eta). \quad (9)$$

Thus (8) takes the form of Riccati equation

$$w_\eta = g(w^2 - k^2). \quad (10)$$

Its solutions are connected via the ladder

$$w_{i+1} = w_i + 1/z_{i+1}, \quad i = 0, 1, 2, \dots, \quad (11)$$

where

$$z_{(i+1)\eta} + 2g w_i z_{i+1} = -g. \quad (12)$$

Choosing $w_0 = k$ we obtain

$$w_1 = k + \frac{2k}{2kC \exp(-2gk\eta) - 1}, \quad C = \text{const}, \quad (13)$$

and then from (2)

$$p_1 = - \frac{2k^3 g C}{\left[\frac{2kC \exp(-kg\eta) - \exp(kg\eta)}{2} \right]^2}. \quad (14)$$

In case when $C = -1/2k$ or $C = 1/2k$ we have soliton,

$$p_1 = g k^2 \text{sech}^2 g k \eta \quad (15)$$

or shock wave [1],

$$p_1 = -g k^2 \text{csch}^2 g k \eta. \quad (16)$$

Taking (12) into account and using (14), from (11) we obtain again

$$w_2 = k + \frac{2k}{2kC \exp(-2gk\eta) - 1} + \frac{4Ck^2 \exp(-2gk\eta)}{[4CC_1 k^2 - \ln |1 - 2kC \exp(-2gk\eta)|] \ln |1 - 2kC \exp(-2gk\eta)|},$$

(17)

$C_1 = \text{const.}$

k is a spectral parameter.

It is easy to note that

$$g = \beta/12\alpha, \quad e = -\beta g^2 k^2/3,$$

$$f = 1 - \frac{1}{e k^2} F. \quad (6)$$

0340-4811 / 85 / 0200-0193 \$ 01.30/0. – Please order a reprint rather than making your own copy.

It is a new solution of the KdV equation.

To end this note we would like to say that we can also obtain from (11) other solutions of the KdV equation having described the function z_{i+1} . Unfortunately they cannot be expressed with the help of analytical functions in general.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

- [1] H. G. Wahlquist and F. B. Estabrook, Phys. Rev. Lett. **31**, 1386 (1973).
- [2] M. S. Marinov, Yad. Fiz. **28**, 251 (1978).
- [3] R. Sasaki, Phys. Lett. **71A**, 390 (1979) and Nucl. Phys. **B154**, 343 (1979).
- [4] A. R. Chowdhury and T. Roy, J. Phys. **A12**, L89 (1979).
- [5] C. Au and P. C. W. Fung, Phys. Rev. **B25**, 25 (1982) and J. Math. Phys. **25**, 1364 (1984).
- [6] T. C. Au-Yeung, C. Au, and P. C. W. Fung, Phys. Rev. **A29**, 2370 (1984).
- [7] P. D. Lax, Pure and Appl. Math. **21**, 467 (1968).
- [8] G. Bowtell, J. Math. Phys. **24**, 969 (1983).
- [9] C. S. Gardner, J. M. Green, M. D. Kruskal, and R. M. Miura, Phys. Rev. Lett. **19**, 1095 (1967).
- [10] F. A. Howes, Stud. Appl. Math. **69**, 75 (1983).
- [11] D. J. Kaup, Phys. **1D**, 391 (1980).
- [12] M. Jaworski, Phys. Lett. **100A**, 321 (1984).
- [13] F. Calogero, Preprint Nr. 70 (1978).
- [14] K. Konno, T. Mitsuhashi, and Y. H. Ichikawa, J. Phys. Soc. Japan **46**, 1907 (1979).
- [15] A. Degasperis, Lett. Nuovo Cim. **33**, 425 (1982).
- [16] Chr. Pöppe, Preprint Nr. 240 (1983).
- [17] M. J. Ablowitz and H. Cornille, Phys. Lett. **72A**, 277 (1979).
- [18] K. Konno and A. Jeffrey, J. Phys. Soc. Japan **51**, 4108 (1982).
- [19] D. Levi, Preprint Nr. 388 (1984).
- [20] R. Hirota, Phys. Rev. Lett. **27**, 1192 (1971).
- [21] R. Hirota, in "Solitons", ed., R. K. Bullough and P. J. Caudrey, Topics in Current Physics 17, Springer-Verlag, Berlin 1980.
- [22] T. Warn, Stud. Appl. Math. **69**, 127 (1983).