

Diffusion in the Liquid System Chloroform + Methyl Acetate

R. Haase and W. Engels

Institut für Physikalische Chemie der Rheinisch-Westfälischen Technischen Hochschule Aachen

Z. Naturforsch. **41a**, 1337–1338 (1986);
received August 23, 1986

We present and discuss the results of measurements of the diffusion coefficient D and the activity coefficients as functions of the mole fraction x of one of the components in the liquid system chloroform + methyl acetate at 10 °C, 30 °C, and 50 °C. The function $D(x)$ exhibits a pronounced maximum at each temperature, while the kinematic diffusion coefficient D^* , considered as function of x , shows a flat minimum, its composition dependence being nearer to a linear relation than that of $D(x)$.

Continuing our work on diffusion in binary liquid systems [1–5], we investigated [6] the non-ideal liquid system chloroform + methyl acetate. The measurements cover the entire range of compo-

sitions at 10 °C, 30 °C, and 50 °C. The diffusion coefficients are derived from interferometric experiments. The limits of reproducibility lie within $\pm 3\%$.

As shown by the isothermal vapour-liquid-equilibrium data [6], the system exhibits negative azeotropy (minimum of vapour pressure at all temperatures). The values of the thermodynamic factor (f_i : activity coefficient, x_i : mole fraction of component i , T : temperature, P : pressure)

$$\beta = 1 + (\partial \ln f_i / \partial \ln x_i)_{T,P}$$

are given in Table 1. We also present the values of the diffusion coefficient D , of the kinematic diffusion coefficient $D^* = D/\beta$ and of the quantities D_0^* and E_D defined by the relation [7]

$$D^* = D_0^* \exp(-E_D/RT).$$

The quantities D_0^* and E_D ("activation energy") depend on the composition only, which we describe by the mole fraction x of methyl acetate.




Fig. 1. Liquid system chloroform + methyl acetate: Diffusion coefficient D as function of the mole fraction x of methyl acetate and of the temperature t_c (see Table 1).

Reprint request to Prof. Dr. R. Haase, Melatener Str. 107, 5100 Aachen.




Fig. 2. Liquid system chloroform + methyl acetate: Kinematic diffusion coefficient D^* as function of the mole fraction x of methyl acetate and of the temperature t_c (see Table 1).

0340-4811 / 86 / 1100-1337 \$ 01.30/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

Table 1. Liquid system chloroform + methyl acetate at 10 °C, 30 °C, and 50 °C. Thermodynamic factor β , diffusion coefficient D , kinetic diffusion coefficient D^* (both in units $10^{-9} \text{ m}^2 \text{ s}^{-1}$) as functions of the mole fraction x of methyl acetate.

$t_c / ^\circ\text{C}$	10			30			50		
x	β	D	D^*	β	D	D^*	β	D	D^*
0.01	1.029	1.88 1.93 1.99	1.88	1.030	2.56	2.49	1.020	3.02 3.20 3.21	3.08
0.10	1.273	2.12 2.23	1.71	1.255	3.04	2.42	1.184	—	—
0.20	1.492	2.70	1.81	1.414	3.36 3.38	2.28	1.326	4.45	3.36
0.30	1.629	2.87	1.76	1.492	3.47	2.33	1.412	4.35 4.39	3.10
0.40	1.670	3.24	1.94	1.504	3.63 3.70	2.44	1.438	4.30 4.55	3.08
0.50	1.616	3.11 3.02	1.90	1.465	3.58	2.44	1.407	4.38 4.54	3.17
0.60	1.484	2.74	1.85	1.389	3.43	2.47	1.327	4.51	3.40
0.70	1.303	—	—	1.290	3.59	2.78	1.218	4.34	3.56
0.80	1.120	—	—	1.183	3.35	2.83	1.103	4.25 4.24	3.85
0.90	0.994	2.77	2.79	1.082	3.49 3.53	3.24	1.017	4.36	4.29
0.99	0.991	2.95	2.98	1.007	3.60 3.33	3.45	0.997	3.94 4.06 4.23	4.09

Table 2. Liquid system chloroform + methyl acetate. The quantities D_0^* and E_D (independent of temperature) as functions of the mole fraction x of methyl acetate.

x	$10^6 D_0^* / (\text{m}^2 \text{ s}^{-1})$	$E_D / (\text{kJ mol}^{-1})$
0.01	0.132	10.0
0.20	0.166	10.6
0.40	0.180	10.9
0.60	0.148	10.2
0.80	0.090	8.6
0.99	0.049	6.7

Figures 1 and 2 show the plots of the functions $D(x)$ and $D^*(x)$ according to our measurements [6].

We see that the graph $D(x)$ exhibits a pronounced maximum, as to be expected for a system with negative deviations from Raoult's Law [5], while the graph $D^*(x)$ shows a flat minimum and is nearer to the straight line than the plot of the function $D(x)$. The numerical values of the activation energy E_D are considerably smaller than in other binary nonelectrolyte solutions [3–5].

For ideal systems ($f_i = 1$, $\beta = 1$, $D = D^*$), such as chlorobenzene + bromobenzene, we have: $E_D = \text{const}$, while D_0^* is a linear function of x [2–4]. In nonideal liquid mixtures there is obviously no general rule with respect to the composition dependence of D^* , D_0^* and E_D .

We are most indebted to Dr. K.-H. Dücker and Dr. H.-J. Jansen for helpful discussions.

- [1] R. Haase und M. Siry, Z. physik. Chem. Neue Folge **57**, 56 (1968).
- [2] R. Haase, R. Kottmann and K.-H. Dücker, Z. Naturforsch. **30a**, 1084 (1975).
- [3] R. Haase, Z. Naturforsch. **31a**, 1025 (1976).
- [4] R. Haase and H.-J. Jansen, Z. Naturforsch. **35a**, 1116 (1980).

- [5] R. Haase and W. Engels, Z. Naturforsch. **38a**, 281 (1983).
- [6] W. Engels, Dissertation, RWTH Aachen, 1986.
- [7] We previously used the symbols B , D^0 and E^* for β , D_0^* and E_D , respectively.