

A Note on the ¹⁴N Electric Field Gradient Tensors in Incommensurate [N(CH₃)₄]₂ZnCl₄

J. Dolinšek and R. Blinc
J. Stefan Institute, E. Kardelj University of Ljubljana,
Ljubljana, Yugoslavia

Z. Naturforsch. 42a, 305–306 (1987);
received December 12, 1986

The ¹⁴N electric field gradient tensors of [N(CH₃)₄]₂ZnCl₄ have been re-determined in the paraelectric phase at 26 °C and in the incommensurate phase at 16 °C. The results in the incommensurate phase show the “non-local” nature of the ¹⁴N EFG tensor interaction.

Tetramethylammonium tetrachlorozincate [N(CH₃)₄]₂ZnCl₄ (TMATC-Zn) belongs to the group of A₂BX₄ crystals. It first transforms with decreasing temperature from the normal (P) to the incommensurate (I) phase and then exhibits at lower temperatures a sequence of commensurate (C) phases. In a recent paper [1] we reported on the ¹⁴N EFG tensors of TMATC-Zn in the paraelectric phase at 26 °C

Reprint requests to J. Dolinšek and Prof. Dr. R. Blinc, J. Stefan Institute, 61000 Ljubljana, Jamova 39, Yugoslavia.

Table 1. ¹⁴N EFG tensors in the crystal fixed frame in paraelectric TMATC-Zn expressed in frequency units (kHz), i.e. multiplied by 3e Q/2h.

$T_0(1,2) =$	19.5	0	0	$;(e^2qQ/h)_{1,2} = 41.4 \text{ kHz}$ $\eta_{1,2} = 0.373$
	0	42.5 ± 4	± 4	
$T_0(3,4) =$	-78	0	0	$;(e^2qQ/h)_{3,4} = 72.4 \text{ kHz}$ $\eta_{3,4} = 0.436$
	0	89 ± 48.5	± 48.5 - 11	

Table 2. ¹⁴N EFG tensors in kHz in the I phase of TMATC-Zn expressed in the crystal fixed a, b, c frame:

$T(x) = T_0 + T_{1''} \cos[\Phi(x) - \Phi_1] + \frac{1}{2} T_{2'} + \frac{1}{2} T_{2''} \cos 2[\Phi(x) - \Phi_2]$											
$T_{1''}$			$T_{2'}$			$T_{2''}$			Φ_1		$\Phi_2 = 0$
$i = 1,2$	0	6	12	0	0	0	-5	0	0	45°	45°
	6	0	0	0	2	± 0.5	0	9	4	45°	0
	12	0	0	0	± 0.5	-0.5	0	4	-4	45°	0
$i = 3,4$	0	5.5	7	-1	0	0	2	0	0	45°	45°
	5.5	0	0	0	1.5	± 5	0	2.5	5	45°	0
	7	0	0	0	± 5	1	0	5	-3	45°	0

0340-4811 / 87 / 0300-0305 \$ 01.30/0. – Please order a reprint rather than making your own copy.

and in the I phase at 16 °C. In that paper the b and c rotations did not correspond to precise rotations about the b and c crystallographic axes, but instead those two axes were tilted for a small angle (θ ≈ 4°) with respect to the rotation axes. This lead to a slight misinterpretation of the ¹⁴N EFG tensors which we would like to correct here.

In the paraphase, which has the space group Pmcn, there are four physically and two chemically nonequivalent ¹⁴N sites in the unit cell. The ¹⁴N nuclei lie on the b–c mirror plane. The two groups of chemically nonequivalent ¹⁴N nuclei can be divided into two sub-groups of physically nonequivalent ¹⁴N nuclei. These two subgroups are related by the glide symmetry which requires the b and c principal axes of the two corresponding physically nonequivalent ¹⁴N EFG tensors to be rotated symmetrically about the a principal axis, where a lies normal to the b–c mirror plane.

One thus expects in the paraphase four different ¹⁴N EFG tensors, where each of the two physically nonequivalent EFG tensors is of the form

$$T_0^\pm = \begin{vmatrix} T_0^{aa} & 0 & 0 \\ 0 & T_0^{bb} & \pm T_0^{bc} \\ 0 & \pm T_0^{bc} & T_0^{cc} \end{vmatrix}, \quad T > T_1. \quad (1)$$

The symmetry of the particular physically non-equivalent tensor in the I phase is described in [1], where it is shown that each tensor element $T^{(\mu)}$ can be expanded in powers of the nuclear displacements from their high temperature equilibrium sites as:

$$T^{(\mu)}(x) = T_0^{(\mu)} + T_1^{(\mu)} \cos[\Phi(x) - \Phi_1^{(\mu)}] + \frac{1}{2} T_2^{(\mu)} + \frac{1}{2} T_2^{(\mu)} \cos 2[\Phi(x) - \Phi_2^{(\mu)}] + \dots \quad (2)$$




Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.


Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.



Figs. 1 a, b, c. Angular dependence of the quadrupole splitting of the ^{14}N spectra in TMATC-Zn at $26^\circ\text{C} > T_1$. The full line is the theoretical fit.



Figs. 2a, b, c. Angular dependence of the incommensurate frequency distribution singularities in the ^{14}N spectra of TMATC-Zn in the I phase at $16^\circ\text{C} < T_1$. The full line is the theoretical fit for the "non-local" model. The hatched area indicates the quasi-continuous distribution of the ^{14}N transition frequencies.

The angular dependence of the ^{14}N quadrupole splitting $2\Delta v_Q$ for $T = 26^\circ\text{C} > T_1$ is shown in Figs. 1a, b, c for rotation around the a , b and c crystal axes. The results show the existence of four physically (and two groups of chemically) nonequivalent ^{14}N sites (Table 1). The experimental error is about ± 2 kHz.

In the I phase at $T = 16^\circ\text{C} < T_1$, $T_0(i)$, $i = 1-4$ is not changed but $T_{1''}(i)$, $T_{2''}(i)$ and $T_{2'}(i)$ are non-

zero and can be determined from the angular variation [1] (Figs. 2a-c) of the incommensurate frequency distribution singularities. The results are collected in Table 2. The discussion of the results within the "non-local" model [2] is, however, correctly described in [1].

[1] J. Dolinšek and R. Blinc, Z. Naturforsch. **41a**, 265 (1986).

[2] R. Blinc, J. Seliger, and S. Žumer, J. Phys. C **18**, 2313 (1985).