

Densities and Molar Excess Volumes of Molten (Alk, Ag)Br- and (Alk, Ag)I-Mixtures

E. Boßmann, M. Müller und J. Richter
Institut für Physikalische Chemie, RWTH Aachen
Z. Naturforsch. **42 a**, 426–428 (1987);
received January 16, 1987

Measurements of densities (and from these molar excess volumes) of the molten salt systems (K,Ag)Br, (Na,Ag)Br, (Rb,Ag)Br, (Cs,Ag)Br, (Na,Ag)I, (K,Ag)I, and (Cs,Ag)I as functions of composition and temperature from the liquidus curve up to 1200 K are reported.

Our method of measuring the densities of molten salts is described in [1]. The results have been fitted to the equation

$$\rho(T, x_2) = (a + bx_2 + cx_2^2 + dx_2^3) + (a' + b'x_2 + c'x_2^2 + d'x_2^3) T, \quad (1)$$

where ρ denotes the density, T the absolute temperature, and x_2 the mole fraction of the silver component.

The obtained coefficients $a, b, c, d, a', b', c', d'$ are listed in Table 1. The molar excess volumes were

Reprint requests to Prof. Dr. J. Richter, Institut für Physikalische Chemie, RWTH Aachen, Templergraben 59, D-5100 Aachen.

evaluated from the equations

$$\bar{V}_{\text{add.}} = (1 - x_2) \frac{M_1}{\rho_1} + x_2 \frac{M_2}{\rho_2}, \quad (2)$$

$$\bar{V} = [(1 - x_2) M_1 + x_2 M_2] / \rho \quad (3)$$

and

$$\bar{V}^E = \bar{V} - \bar{V}_{\text{add.}}, \quad (4)$$

where M_1 and M_2 are the molar weights and ρ_1 and ρ_2 the densities of the pure salts.

The excess volumes of the (Alk, Ag)Br- and (Alk, Ag)I-mixtures are listed in Tables 2, and plotted in Figure 1. Depending on their composition, the (Alk, Ag)Br systems have positive and negative excess volumes. In contrast to the (Alk, Ag)Br- and (Alk, Ag)Cl-systems [1], all investigated (Alk, Ag)I systems have negative excess volumes.

The volume fraction of component 2 is defined as

$$\psi_2 = x_2 \frac{M_2}{\rho_2} \left[(1 - x_2) \frac{M_1}{\rho_1} + x_2 \frac{M_2}{\rho_2} \right]. \quad (5)$$

Some deviations from linearity are shown in Figures 2 and 3. With the exception of (Na, Ag)I, for the (Alk, Ag)Cl- [1], (Alk, Ag)Br-, (Alk, Ag)I-

Table 1. Constants for the calculation of the densities of the molten (Alk, Ag)I- and (Alk, Ag)Br-mixtures, depending on composition and temperature.

	(Na, Ag)I	(K, Ag)I	(Cs, Ag)I
$a \text{ (g cm}^{-3}\text{)}$	3.56847	3.17252	3.92596
$b \text{ (g cm}^{-3}\text{)}$	2.38356	3.58116	3.58490
$c \text{ (g cm}^{-3}\text{)}$	1.52402	−1.99325	−2.86987
$d \text{ (g cm}^{-3}\text{)}$	−1.11780	1.66507	1.77602
$a' (\cdot 10^3 \text{ g cm}^{-3})$	−0.89785	−0.79940	−0.91824
$b' (\cdot 10^3 \text{ g cm}^{-3})$	−0.18664	−1.15796	−1.90151
$c' (\cdot 10^3 \text{ g cm}^{-3})$	−0.76515	1.72076	2.26142
$d' (\cdot 10^3 \text{ g cm}^{-3})$	0.86072	−0.78801	−0.44250

	(Na, Ag)Br	(K, Ag)Br	(Rb, Ag)Br	(Cs, Ag)Br
$a \text{ (g cm}^{-3}\text{)}$	3.20989	2.96403	3.74662	4.27735
$b \text{ (g cm}^{-3}\text{)}$	2.47619	3.72788	1.58203	−0.39662
$c \text{ (g cm}^{-3}\text{)}$	1.39961	−1.67602	0.84282	3.70048
$d \text{ (g cm}^{-3}\text{)}$	−0.76801	1.26602	0.14070	−1.22769
$a' (\cdot 10^3 \text{ g cm}^{-3})$	−0.83611	−0.82130	−1.05919	−1.22946
$b' (\cdot 10^3 \text{ g cm}^{-3})$	−0.39076	−1.76399	−0.35752	1.27793
$c' (\cdot 10^3 \text{ g cm}^{-3})$	−0.76457	2.11354	0.18654	−3.82505
$d' (\cdot 10^3 \text{ g cm}^{-3})$	0.56445	−0.50536	0.18759	2.72832

0340-4811 / 87 / 0400-0426 \$ 01.30/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.




Fig. 1. Molar excess volumes of the (Alk, Ag)Br- and (Alk, Ag)I-mixtures at 1050 K.

Table 2. Molar excess volumes $\bar{V}^E$ of (Alk, Ag)Br- and (Alk, Ag)I-mixtures at 1050 K in units cm^3/mol .

	(K, Ag)Br	(Na, Ag)Br	(Rb, Ag)Br	(Cs, Ag)Br
0	0	0	0	0
0.1	0.302	0.413	0.525	-0.264
0.2	0.561	0.594	0.766	-0.118
0.3	0.731	0.620	0.806	0.150
0.4	0.800	0.553	0.702	0.354
0.5	0.770	0.438	0.513	0.397
0.6	0.663	0.308	0.296	0.275
0.7	0.501	0.187	0.099	0.052
0.8	0.317	0.088	-0.035	-0.160
0.9	0.141	0.024	-0.076	-0.222
1	0	0	0	0

	(Na, Ag)I	(K, Ag)I	(Cs, Ag)I
0	0	0	0
0.1	-0.249	-1.165	-0.744
0.2	-0.498	-1.669	-0.955
0.3	-0.716	-1.777	-0.883
0.4	-0.871	-1.658	-0.696
0.5	-0.949	-1.423	-0.506
0.6	-0.941	-1.130	-0.362
0.7	-0.842	-0.847	-0.275
0.8	-0.652	-0.560	-0.229
0.9	-0.371	-0.282	-0.151
1	0	0	0




Fig. 2. The densities ρ of (Na, Ag)Br plotted against x_2 and ψ_2 at 1050 K.




Fig. 3. The densities ρ of (Na, Ag)I plotted against x_2 and ψ_2 at 1050 K.

systems the dependence of ϱ on Ψ_2 is more nearly linear than that of ϱ on x_2 .

With our experimental setup we also tried to investigate the system (Rb, Ag)I. The molten salt mixtures of this system heavily evolve vapour, which influences the measurements so that they are not reported here.

Acknowledgements

We thank the Minister für Wissenschaft und Forschung des Landes Nordrhein-Westfalen, Düsseldorf, and the Fonds der Chemischen Industrie, Frankfurt, for financial support of this work.

- [1] A. Krekelberg, W. Merkens, and J. Richter, Z. Naturforsch. **38a**, 890 (1983).