

## Oxidation of a Cyanine Dye with o-Cl-Dibenzoyl Peroxide and Peroxydisulphate

A. M. Habib, M. H. Abdel-Kader, M. A. Salem, and S. T. Abdel-Halim

Chemistry Department, Faculty of Science, Tanta University, Tanta, Egypt

Z. Naturforsch. **43a**, 614–616 (1988); received March 31, 1988

The oxidation of the cyanine dye 1-methyl-4-N-dimethylamine- $\gamma$ -stilbazolium iodide with o-Cl dibenzoyl peroxide and peroxydisulphate ion is kinetically investigated. In both cases the oxidation process obeys second order kinetics, being first order in the oxidant molecule and first order in the cyanine dye. The thermodynamic parameters evaluated for the reaction between the dye and o-Cl-DBP are  $E_a = 45.6 \text{ kJ mol}^{-1}$ ,  $\Delta H^\ddagger = 43.09 \text{ kJ mol}^{-1}$ , and  $\Delta S^\ddagger = -115.7 \text{ J deg}^{-1} \text{ mol}^{-1}$ . The corresponding values of  $E_a$ ,  $\Delta H^\ddagger$ , and  $\Delta S^\ddagger$  obtained for the dye/ $\text{S}_2\text{O}_8^{2-}$  system are  $58.01 \text{ kJ mol}^{-1}$ ,  $55.5 \text{ kJ mol}^{-1}$ , and  $-147.8 \text{ J deg}^{-1} \text{ mol}^{-1}$ . Proposed mechanisms are discussed.

**Key words:** Oxidation of cyanine dyes, peroxydisulphate and organic peroxides.

### Introduction

Merocyanine and cyanine dyes are photoreactive compounds whose solvatochromism is interesting as to colour and constitution [1–3]. The solvents play a great role in the behaviour of the charge transfer absorption bands. Thus, such acts are ideal solvent-indicators. The ground and excited states of some merocyanine dyes and their photosensitization have been investigated [4]. However information about the thermally induced oxidation of cyanine dyes seems to be lacking. This prompted us to study such oxidation reactions using o-Cl-dibenzoyl peroxide and peroxydisulphate ion as oxidizing agents in neutral media.

### Experimental

The cyanine dye (1-methyl-4-N-dimethylamine- $\gamma$ -stilbazolium iodide) was prepared by mixing p-N-dimethylamino benzaldehyde with N-methyl picolinium iodide in the presence of piperidine as catalyst in a way similar to the method described by Minch and Shah [5]. o-Cl-dibenzoyl peroxide prepared and purified according to the method described by Price and Krobs [6]. All other chemical reagents were analytical grade substances and used as received.

The kinetics were studied with two methods. In the first, method the order of the reaction of the cyanine dye was determined. This was done spectrophotometrically under an at least tenfold excess of the oxidant (i.e. under pseudo-first order conditions). The two solutions were mixed and the rate of the decrease in the absorbance of the dye (at 450 nm) was

recorded with a uv/vis Pye Unicam SP 8000 spectrophotometer. The solvent was ethanol in case of cyanine/o-Cl-DBP and water in case of cyanine/ $\text{S}_2\text{O}_8^{2-}$ . The second method was used for the evaluation of the kinetic order for the oxidants. This was done iodometrically following the procedure reported by Kolthoff and Carr [7].




Fig. 1. The change in the absorbance of cyanine dye during the reaction with o-Cl-DBP at 50°C in ethanol.




Fig. 2. Logarithm of the absorbance of the cyanine dye vs. time at four temperatures: (○) 45°C; (●) 50°C; (△) 55°C; (□) 60°C.




Fig. 3. Arrhenius plots for the oxidation of cyanine dye with: (●) o-Cl-DBP; and (○)  $\text{S}_2\text{O}_8^{2-}$ .

Reprint requests to Prof. Dr. A. M. Habib, Chemistry Department, Faculty of Science, Tanta University, Tanta/Ägypten.

0932-0784 / 88 / 0600-0614 \$ 01.30/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

Table 1. Thermodynamic parameters obtained for the oxidation reaction of cyanine/o-Cl-DBP and cyanine/S<sub>2</sub>O<sub>8</sub><sup>2-</sup> systems

| Substrate                                           | <i>T</i> /C° | <i>k</i> × 10 <sup>3</sup><br>mol l <sup>-1</sup> min <sup>-1</sup> | <i>E</i> <sub>a</sub><br>[kJ mol <sup>-1</sup> ] | <i>ΔG</i> <sup>*</sup><br>[kJ mol <sup>-1</sup> ] | <i>ΔH</i> <sup>*</sup><br>[kJ mol <sup>-1</sup> ] | <i>ΔS</i> <sup>*</sup><br>[J deg <sup>-1</sup> mol <sup>-1</sup> ] |
|-----------------------------------------------------|--------------|---------------------------------------------------------------------|--------------------------------------------------|---------------------------------------------------|---------------------------------------------------|--------------------------------------------------------------------|
| Cyanine/o-Cl-DBP                                    | 45           | 3.6                                                                 | 58.01                                            | 103.62                                            | 55.5                                              | -147.8                                                             |
|                                                     | 50           | 5.8                                                                 |                                                  |                                                   |                                                   |                                                                    |
|                                                     | 55           | 8.3                                                                 |                                                  |                                                   |                                                   |                                                                    |
|                                                     | 60           | 15.0                                                                |                                                  |                                                   |                                                   |                                                                    |
| Cyanine/S <sub>2</sub> O <sub>8</sub> <sup>2-</sup> | 15           | 12.0                                                                | 45.6                                             | 77.29                                             | 43.1                                              | -115.7                                                             |
|                                                     | 24           | 17.6                                                                |                                                  |                                                   |                                                   |                                                                    |
|                                                     | 29           | 24.0                                                                |                                                  |                                                   |                                                   |                                                                    |
|                                                     | 34           | 37.0                                                                |                                                  |                                                   |                                                   |                                                                    |

## Results and Discussion

The absorbance of the cyanine dye in the presence of o-Cl-DBP changed with time as shown in Figure 1. On plotting the logarithm of the absorbance of unreacted cyanine vs. time, straight lines were obtained (Fig. 2), indicating a first order reaction with respect to the dye molecule. The plot obeys the equation

$$\log A_t = \log A_0 - \frac{k' t}{2.303}$$

*k'* equals *k* [P] where *k* is the true rate constant and [P] the initial concentration of the o-Cl-DBP. The measurements were repeatedly taken at various temperatures in the range 45–60°C. At temperatures below 65°C spontaneous thermal decomposition of the peroxide was undetectable. When the peroxydisulphate ion was substituted for the organic peroxide and the kinetic runs were carried out in the temperature range 15–34°C, a first order reaction for the dye was again observed. Also in the iodometric measurements, where the molar ratio of the dye and either of the oxidizing agents was held as high as 10:1, the reaction was found to be of first order. Thus, the overall reaction between the cyanine dye and o-Cl-DBP or S<sub>2</sub>O<sub>8</sub><sup>2-</sup>-ion follows second order kinetics

according to the equation

$$\text{rate} = k_2 [\text{oxidant}] [\text{cyanine}],$$

where *k*<sub>2</sub> is the second order rate constant. Linear relationships were obtained when plotting log *k*<sub>2</sub> vs. 1/*T* (Figure 3). The derived activation parameters are summarized in Table 1.


## Mechanism

### a) Oxidation with o-Cl-DBP

The mechanism resembles that reported by Tokumaru and Simamura [8]. The donating property of the solvent is important for this type of oxidation reactions. Thus, with donating solvents the reaction proceeds more rapidly compared with less donating ones [8, 9].

### b) Oxidation with S<sub>2</sub>O<sub>8</sub><sup>2-</sup>

Since the reaction of cyanine dye with peroxydisulphate followed first-order kinetics in both the dye and the S<sub>2</sub>O<sub>8</sub><sup>2-</sup>-ion, and this is in good agreement with the results reported by Choudary et al. [10] on the aromatic amines/S<sub>2</sub>O<sub>8</sub><sup>2-</sup> reactions, the mechanism may be as follows:



Inspection of Table 1 shows that the energy of activation of the cyanine/S<sub>2</sub>O<sub>8</sub><sup>2-</sup> reaction is lower than that of the cyanine/o-Cl-DBP reaction. This may be due to several factors, viz, steric effects, molecular sizes, diffusion coefficients and bond dissociation energies. The O-O-bond dissociation

energies characteristic of the peroxydisulphate [11] ion and peroxide [12] molecule are 110–120 and 130 kJ mol<sup>-1</sup>, respectively. This would explain the faster oxidation rate observed in the presence of S<sub>2</sub>O<sub>8</sub><sup>2-</sup>-ion than in the presence of the peroxide.

- [1] S. Dähne, *Science* **199**, 1163 (1978).
- [2] K. S. Nolte and S. Dähne, *Adv. Molec. Relax. Processes* **10**, 299 (1977).
- [3] J. Griffiths, "Colour and Constitution of Organic Molecules", Academic Press London 1976.
- [4] F. Dietz, G. Müller, G. Bach, and J. von Grossman, *J. Signal A. M.* **3**, 1, 35 (1975).
- [5] M. J. Minch and S. S. Shah, *J. Chem. Educ.* **54**, 707 (1977).
- [6] C. C. Price and E. Krobs, *Org. Synth.* **23**, 65 (1943).
- [7] I. M. Kolthoff and E. M. Carr, *Anal. Chem.* **25**, 298 (1953).
- [8] K. Tokumarn and O. Simamura, *Bull. Chem. Soc. Japan* **36**, 333 (1963).
- [9] F. Minisci, C. Giordano, E. Vismara, S. Levi, and V. Tortelli, *J. Amer. Chem. Soc.* **106**, 7146 (1984).
- [10] V. Choudary and A. K. Bhattacharya, *J. Chem. Soc., Perkin Trans. II* **1983**, 849.
- [11] V. K. Gupta and S. P. Srivastava, *Oxidation Commun.* **4**, 75 (1983).
- [12] K. Nozaki and P. D. Bartlett, *J. Amer. Chem. Soc.* **68**, 1686 (1946).