

Hydrogen in Electrodynamics II. Mathematical Means

Hans Sallhofer

Forschungs- und Versuchsantalt
der AUSTRIA METALL AG, Braunau, Austria

Z. Naturforsch. **44a**, 167–168 (1989);
received January 20, 1989

After a discussion of the one-component Schrödinger (1926) and the four-component Dirac (1928) representation of hydrogen it is shown that the six-component electrodynamic picture turns out to be considerably simpler and clearer. The computational effort is reduced to a fraction.

For the hydrogen-solution of electrodynamics, as we shall see later, we need the simplest form in polar coordinates for the expressions

$$(\partial_1 + i \partial_2) \Psi = (\partial_1 + i \partial_2) C R(r) P_l^m(\cos \theta) e^{im\phi} \quad (1)$$

and $(C: \text{constant})$

$$\partial_3 \Psi = \partial_3 C R(r) P_l^m(\cos \theta) e^{im\phi}, \quad (2)$$

In order to obtain these two simplicia we consider the well-known operators

$$(\partial_1 \pm i \partial_2) = e^{\pm i\phi} \left(\sin \theta \partial_r + \frac{\cos \theta}{r} \partial_\theta \pm \frac{i}{r \sin \theta} \partial_\phi \right) \quad (3)$$

and

$$\partial_3 = \cos \theta \partial_r - \frac{\sin \theta}{r} \partial_\theta \quad (4)$$

as well as the associated spherical harmonics

$$P_l^m(\cos \theta) = \sin^m \theta \frac{d^m P_l(\cos \theta)}{d \cos^m \theta}, \quad 0 \leq |m| \leq l \quad (5)$$

In addition we need the well-known recurrence relations for the spherical harmonics and the simple representation of the trigonometric functions by spherical harmonics [1]:

$$(2l+1) \sin \theta P_l^m = (l+m-1)(l+m) P_{l-1}^{m-1} - (l-m+1)(l-m+2) P_{l+1}^{m-1}, \quad (6)$$

$$(2l+1) \left(\frac{m}{\sin \theta} P_l^m - \sin \theta \cos \theta d_{\cos \theta} P_l^m \right) = (l+1)(l+m)(l+m-1) P_{l-1}^{m-1} + l(l-m+1)(l-m+2) P_{l+1}^{m-1} \quad (7)$$

$$(2l+1) \sin^2 \theta d_{\cos \theta} P_l^m = (l+1)(l+m) P_{l-1}^m - l(l-m+1) P_{l+1}^m, \quad (8)$$

$$(2l+1) \cos \theta P_l^m = (l+m) P_{l-1}^m + (l-m+1) P_{l+1}^m, \quad (9)$$

$$\sin \theta = \frac{P_{l+1}^{m+1} - P_{l-1}^{m+1}}{(2l+1) P_l^m}, \quad (10)$$

$$\cos \theta = \frac{(l-m+1) P_{l+1}^m + (l+m) P_{l-1}^m}{(2l+1) P_l^m}. \quad (11)$$

If we replace m by $(-m)$ in (6) and (7) and employ the well-known relation [2]

$$P_l^{-m} = (-1)^m \frac{(l-m)!}{(l+m)!} P_l^m, \quad (12)$$

we then obtain the even simpler recurrence relations

$$(2l+1) \sin \theta P_l^m = -P_{l-1}^{m+1} + P_{l+1}^{m+1}, \quad (13)$$

$$(2l+1) \left(\frac{m}{\sin \theta} P_l^m + \sin \theta \cos \theta d_{\cos \theta} P_l^m \right) = (l+1) P_{l-1}^{m+1} + l P_{l+1}^{m+1}. \quad (14)$$

In order to finally obtain the desired form in polar coordinates for (1) we insert (3) and use (13) and (14) in the resulting expression. Then we get

$$(\partial_1 + i \partial_2) C R P_l^m e^{im\phi} = \frac{C e^{i(m+1)\phi}}{2l+1} \left(- \left(d_r + \frac{l+1}{r} \right) R P_{l-1}^{m+1} + \left(d_r - \frac{l}{r} \right) R P_{l+1}^{m+1} \right). \quad (15)$$

Taking into account also negative values of ϕ , this becomes

$$(\partial_1 \pm i \partial_2) C R P_l^m e^{\pm im\phi} = \frac{C e^{\pm i(m+1)\phi}}{2l+1} \left(- \left(d_r + \frac{l+1}{r} \right) R P_{l-1}^{m+1} + \left(d_r - \frac{l}{r} \right) R P_{l+1}^{m+1} \right). \quad (16)$$

For more clarity we introduce the operator

$$(d_r + a/r) R = R_{,a} \quad (a: \text{arbitrary}) \quad (17)$$

Reprint requests to Prof. Dr. H. Sallhofer, Bahnhofstrasse 36, A-5280 Braunau, Austria.

0932-0784 / 89 / 0200-0167 \$ 01.30/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

and thus obtain for (1) the desired form

$$(\partial_1 \pm i \partial_2) CRP_l^m e^{\pm im\phi} = \frac{C e^{\pm i(m+1)\phi}}{2l+1} (-R_{,l+1} P_{l-1}^{m+1} + R_{,-l} P_{l+1}^{m+1}). \quad (18)$$

In order to obtain (2) also in an analogous form, we first insert from (4) and then from (8), (9) and (17). Then we get

$$\partial_3 CRP_l^m e^{im\phi} = \frac{C e^{im\phi}}{2l+1} [R_{,l+1}(l+m) P_{l-1}^m + R_{,-l}(l-m+1) P_{l+1}^m]. \quad (19)$$

Taking into account negative values of ϕ , we then obtain the desired relation also in this case

$$\partial_3 CRP_l^m e^{\pm im\phi} \quad (20)$$

$$= \frac{C e^{\pm im\phi}}{2l+1} [R_{,l+1}(l+m) P_{l-1}^m + R_{,-l}(l-m+1) P_{l+1}^m].$$

Moreover we note that the spherical harmonics have been generalized some time ago such that the most general form turns out to an entire function of m and l [3]. In such a generality the functions are therefore unique and continuous and can be differentiated after both indices. Halfinteger forms as, e.g.,

$$P_l^{m+1/2} = P_l^{m+1/2}(\cos \theta) \quad (21)$$

thus are uniquely defined.

[1] Proved in closed form in: Handbuch der Radiologie, Bd. VI, p.102, Leipzig 1933, – or in modern form in: H. A. Bauer, Grundlagen der Atomphysik, Springer, Wien 1951.

[2] K. Jänich, Analysis, p. 353, Springer, Berlin 1983.

[3] F. W. Schäfke, Einführung in die Theorie der speziellen Funktionen der Mathematischen Physik, Springer, Berlin 1963.