

On Electromagnetic Spinors and Electron Theory

Warren H. Inskeep

Z. Naturforsch. **44a**, 327–328 (1989);
received February 23, 1989

The relationship between the Dirac theory and electromagnetic spinors is extended to the case of finite mass. Certain products of the electromagnetic fields give rise to the Dirac differential operator upon the usual substitutions for the energy and momentum. By placing mass in the proper place for the wave mechanical approach to quantum theory, the algebra of the fields, interpreted as quantum operators, may be deduced.

It is nearly as easy to derive the Dirac equation starting from electrodynamics as it is by starting from mechanics. The electromagnetic fields always satisfy the relation

$$(H^2 + E^2)^2 - (2\mathbf{E} \times \mathbf{H})^2 \equiv \mathcal{E}^2 - P^2 \\ = (H^2 - E^2)^2 + 4(\mathbf{E} \cdot \mathbf{H})^2. \quad (1)$$

We may set $\mathbf{E} \cdot \mathbf{H} = 0$, identify $(H^2 - E^2)$ as the mass, eventually use the substitutions $\mathcal{E} \rightarrow i \frac{\partial}{\partial t}$, $\mathbf{P} \rightarrow -i \nabla$, and proceed in the usual manner [1]. The only new feature here is the possibility of speculating that mass may have an electromagnetic origin. The electromagnetic approach, however, becomes considerably more interesting by introducing the concept of an electromagnetic spinor [2].

A spinor may be viewed as arising from a vector of zero length [3]. For a three-vector \mathbf{x} , we thus require

$$x_1^2 + x_2^2 + x_3^2 = 0. \quad (2)$$

Then define two numbers ξ_0, ξ_1 such that

$$x_1 = \xi_0^2 - \xi_1^2, \quad x_2 = i(\xi_0^2 + \xi_1^2), \quad x_3 = -2\xi_0\xi_1, \quad (3)$$

which are consistent with

$$\xi_0 = \pm \sqrt{\frac{x_1 - ix_2}{2}}, \quad \xi_1 = \sqrt{\frac{-x_1 - ix_2}{2}}. \quad (4)$$

The two component object $\xi = (\xi_0, \xi_1)$ transforms as a spinor. The zero length vector and its associated

spinor are also related by

$$(\boldsymbol{\sigma} \cdot \mathbf{x}) \xi = 0, \quad (5)$$

which will prove useful below.

Let us define the fields

$$\Phi_R = \boldsymbol{\sigma} \cdot (\mathbf{H} - i\mathbf{E}), \quad \Phi_L = \boldsymbol{\sigma} \cdot (\mathbf{H} + i\mathbf{E}), \quad (6)$$

where the subscripts are suggested by the fact that \mathbf{E} changes sign under spatial inversion, but \mathbf{H} is invariant. Then with the identity

$$(\boldsymbol{\sigma} \cdot \mathbf{a})(\boldsymbol{\sigma} \cdot \mathbf{b}) = \mathbf{a} \cdot \mathbf{b} + i\boldsymbol{\sigma} \cdot (\mathbf{a} \times \mathbf{b}) \quad (7)$$

the requirement that the electromagnetic field have zero length at any point of space-time may be written as

$$\Phi_L \Phi_R + \Phi_R \Phi_L = 2(H^2 + E^2) = 0. \quad (8)$$

The corresponding electromagnetic spinors φ_L, φ_R will, in view of (5), satisfy

$$\Phi_L \varphi_L = 0, \quad \Phi_R \varphi_R = 0. \quad (9)$$

The individual terms in (8) are

$$\Phi_L \Phi_R = H^2 + E^2 - 2\boldsymbol{\sigma} \cdot (\mathbf{E} \times \mathbf{H}) = \mathcal{E} - \boldsymbol{\sigma} \cdot \mathbf{P}, \quad (10)$$

$$\Phi_R \Phi_L = H^2 + E^2 + 2\boldsymbol{\sigma} \cdot (\mathbf{E} \times \mathbf{H}) = \mathcal{E} + \boldsymbol{\sigma} \cdot \mathbf{P}, \quad (11)$$

which clearly give the Dirac differential operator with the substitutions for \mathcal{E}, \mathbf{P} as above. We further note that the Lagrangian density may be written as

$$\Phi_L \Phi_L = (H^2 - E^2) + 2i(\mathbf{E} \cdot \mathbf{H}), \quad (12)$$

$$\Phi_R \Phi_R = (H^2 - E^2) - 2i(\mathbf{E} \cdot \mathbf{H}), \quad (13)$$

From (10)–(13) it follows that

$$\Phi_L \Phi_R \Phi_R \Phi_L = \Phi_R \Phi_R \Phi_L \Phi_L, \quad (14)$$

which is identical to (1) (and presumably corresponds to the Klein-Gordon equation in the quantum interpretation).

Giving the wave mechanical interpretation to the right sides of (10) and (11), and following Feynman [4], applying $(\mathcal{E} - \boldsymbol{\sigma} \cdot \mathbf{P})$ to the right-handed spinor gives an object that transforms as a left handed spinor, and similarly for $(\mathcal{E} + \boldsymbol{\sigma} \cdot \mathbf{P})$. Setting

$$(\mathcal{E} - \boldsymbol{\sigma} \cdot \mathbf{P}) \varphi_R = m \varphi_L, \quad (15)$$

$$(\mathcal{E} + \boldsymbol{\sigma} \cdot \mathbf{P}) \varphi_L = m \varphi_R, \quad (16)$$

gives coupled equations that transform correctly, and these together are the Dirac equation in the Weyl

Reprint requests to W. H. Inskeep, 4305 SW 185th Ave., Aloha, Oregon, USA 97007.

0932-0784 / 89 / 0400-0327 \$ 01.30/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

representation. But from the left sides of (10) and (11) we can equally well write these as

$$(\Phi_L \Phi_R) \varphi_R = m \varphi_L, \quad (17)$$

$$(\Phi_R \Phi_L) \varphi_L = m \varphi_R. \quad (18)$$

Next, using (9) the left sides of these equations can only give zero. To avoid this let us introduce the interpretation familiar from quantum field theory according to which the fields (6) annihilate their correspond-

ing spinors, giving the vacuum state:

$$\Phi_L |\varphi_L\rangle = |0\rangle, \quad \Phi_R |\varphi_R\rangle = |0\rangle. \quad (19)$$

Now (17) and (18) imply

$$\Phi_L |0\rangle = m |\varphi_L\rangle, \quad \Phi_R |0\rangle = m |\varphi_R\rangle. \quad (20)$$

Whence, Φ_L and Φ_R both annihilate and create their corresponding spinors. Note that this result is consistent with the expressions for the Lagrangian density, (12) and (13), if $\mathbf{E} \cdot \mathbf{H} = 0$, and $(H^2 - E^2)$ is taken as the mass.

- [1] P. A. M. Dirac, The Principles of Quantum Mechanics, 4th ed., Oxford University Press, Oxford 1958, pp. 254–258.
- [2] W. H. Inskeep, Z. Naturforsch. **43a**, 695 (1988).

- [3] E. Cartan, The Theory of Spinors, M.I.T. Press, Cambridge 1966.
- [4] R. P. Feynman, Theory of Fundamental Processes, W. A. Benjamin, New York 1962, pp. 112–115.