

Deuterium Isotope Effects on the Silver Ion-Promoted Hydrolyses of S,S-Diethylacetals of p-Substituted Benzaldehydes

Derek P. N. Satchell and Rosemary S. Satchell

Z. Naturforsch. **44a**, 492–493 (1889);


received January 3, 1989

This paper is dedicated to Professor Jacob Bigeleisen on the occasion of his 70th birthday



Solvent deuterium isotope effects support the view that the p-NO₂ and p-Me derivatives of benzaldehyde S,S-diethylacetal undergo silver ion-promoted hydrolysis in water by different mechanisms.

Introduction

Silver ions promote the hydrolysis of S-acetals in aqueous solution. The kinetic and mechanistic details depend upon the structure of the S-acetal [1, 2]. For the (open-chain) S,S-diethylacetals of substituted benzaldehydes,



we have found [2] that the dominant path leading to hydrolysis when $[\text{Ag}^+] \geq 0.1 \text{ M}$ is that given in outline in the equations



In the presence of an excess of silver ions, the observed first-order rate constant for loss of acetal, k_{obs} , is given by

$$k_{\text{obs}} = k_1 K_1 [\text{Ag}^+] / (1 + K_1 [\text{Ag}^+]). \quad (5)$$

Using solutions in 1% (v/v) dioxane-water we have shown [2] that plots of (5) lead to values of K_1 and k_1 for various S,S-diethylacetals. Comparisons of the k_1 values, and of the corresponding activation parameters, led us to suggest [2] that there occurs a change in the detailed mechanism of step (3) on passing from the S,S-acetals with R=p-MeO and p-Me to the p-NO₂ derivative: for the former acetals we postulated a unimolecular mechanism analogous to the A1

scheme common [3] for O,O-acetals in the presence of H₃O⁺ ions, while for the latter we suggested a bimolecular A2-type mechanism with one (or more) water molecules in the slow phase of step (3). The compound with R=H may be an intermediate case [1, 2].

Solvent deuterium isotope effects are sometimes helpful in distinguishing between reaction mechanisms in solution [4]. Relatively few studies, however, have examined such isotope effects in systems involving metal ion-promotion or catalysis. No general pattern of behaviour has been established in the way that it has for H₃O⁺-catalysed processes [4]. We have now studied the effect of using D₂O in place of H₂O on some of the results for two of these S,S-acetals: the p-Me and p-NO₂ compounds.

Experimental

Deuterium oxide was the Goss product (99.8%). Our methods, concentration and other conditions were similar to those used [1, 2] for H₂O solutions, and the reproducibility of k_{obs} ($\pm 10\%$) was also similar to that obtained in H₂O. Our results are compared with those for H₂O in Figure 1. The derived K_1 and k_1 values were obtained via the slopes and intercepts of plots both of $1/k_{\text{obs}}$ against $1/[\text{Ag}^+]$, and of $[\text{Ag}^+]/k_{\text{obs}}$ against $[\text{Ag}^+]$. Good rectilinear behaviour was observed, in keeping with (5).

Discussion

It is evident from Fig. 1 that the change to D₂O has little effect on the reaction rate of the p-Me compound. At 25.0 °C the derived values of K_1 (dm³ mol⁻¹) and k_1 (s⁻¹) for the hydrolyses in H₂O and D₂O are $(K_1)_{\text{H}_2\text{O}} = 230 \pm 10$, $(K_1)_{\text{D}_2\text{O}} = 235 \pm 10$, $(k_1)_{\text{H}_2\text{O}} = 1.6 \pm 0.5$, $(k_1)_{\text{D}_2\text{O}} = 1.6 \pm 0.5$. The absence of a significant isotope effect on k_1 is compatible [4] with our suggestion [2] that water species are not involved in the slow phase of step (3) for the p-Me derivative. That the K_1 values are also so close is interesting: it could imply that the net effects on ΔG^\ddagger of the enthalpy and entropy changes accompanying the return of one (possibly more) water molecule(s) from the silver ion solvation water to the bulk on forming the adduct (I) are, in this case, much the same for both H₂O and D₂O.

For the p-NO₂ derivative the change of solvent has a clearly detectable effect (Figure 1). At 44.2 °C the derived parameters are $(K_1)_{\text{H}_2\text{O}} = 9 \pm 1$, $(K_1)_{\text{D}_2\text{O}} = 18 \pm 2$, $(k_1)_{\text{H}_2\text{O}} = 7.0 \pm 0.8$, and $(k_1)_{\text{D}_2\text{O}} = 4.1 \pm 0.5$. The relative k_1 values appear compatible [4] with our suggestion that one or more water molecules are involved in a nucleophilic attack on the pro-carbonyl carbon atom in the slow phase of step (3) for the p-NO₂ compound and point to some change in mechanism compared with the p-Me derivative. For the p-NO₂ compound the K_1 values in the two solvents are also notably different. In previous work [2] we found that the thermodynamic quantities ΔH^\ddagger and ΔS^\ddagger corresponding to equilibrium (2) in H₂O have values $(77 \pm 5) \text{ kJ mol}^{-1}$ and (259 ± 10)

Reprint requests to Dr. D. P. N. Satchell, King's College, London, Strand, WC2R 2LS, United Kingdom.

0932-0784 / 89 / 0500-0492 \$ 01.30/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.




Fig. 1. Variation of k_{obs} with $[\text{Ag}^+]$ in H_2O and in D_2O . **A:** p-methyl derivative at 25.0 °C; **B:** p-nitro derivative at 44.2 °C. $[\text{S,S-acetal}]_{\text{initial}} \cong 10^{-5} \text{ M}$, $[\text{H}_3\text{O}^+] = 0.05 \text{ M}$, and ionic strength = 0.51 M in all runs; k_{obs} insensitive to $[\text{H}_3\text{O}^+]$, see [1]; D_2O ○, H_2O ●.

J K mole^{-1} , respectively, for the p- NO_2 derivative, whereas for the p-Me (and the p-MeO) compound both ΔH° and ΔS° are negative. We interpreted this difference in thermodynamic parameters in terms of a much greater degree of desolvation of the reactants (especially the Ag^+ ion) being required for formation of (I) with the p- NO_2 compound owing to the latter's relative weakness as an S-base. The isotope

effect now found seems to us possibly compatible with that picture, and certainly indicates some difference in the details of adduct formation for the p- NO_2 and p-Me compounds. In all, the isotope effects evidently support our view that the p- NO_2 and p-Me compounds react via mechanisms containing significant differences in detail.

- [1] D. Penn, Z. Saidi, D. P. N. Satchell, and R. S. Satchell, *J. Chem. Res. (S)* 200 (1987) and earlier papers.
 [2] D. P. N. Satchell and R. S. Satchell, *J. Chem. Res. (S)* **1989**, 102.

- [3] E. H. Cordes and H. G. Bull, *Chem. Rev.* **74**, 581 (1974).
 [4] P. M. Laughton and R. E. Robinson, in: *Solute-Solvent Interactions* (J. F. Coetzee and D. C. Ritchie, eds.), Marcel Dekker, New York 1969, Chapt. 7.

