

On the Spectra of Turbulent Fluids at Large k

H. Tasso

Max-Planck-Institut für Plasmaphysik,
Euratom association, Garching bei München, F.R.G.

Z. Naturforsch. **45a**, 928 (1990); received May 4, 1990

The turbulence spectra of continuous fluids at large k are discussed. A basic difference in behaviour appears between conservative and dissipative systems. Only in the latter case can one eliminate higher order ultraviolet divergences.

In previous work on the equilibrium statistics of the Korteweg de Vries equation (see [1, 2]) and other 1-d non-integrable but dispersive equations [3], it was possible to prove that the space correlation function has an exponential decay like $e^{-\alpha|x|}$. This was the result of the simultaneous presence of nonlinearity and dispersion in the equation. Otherwise the correlation function would have a $\delta(x)$ behaviour.

The k -spectrum being the Fourier transform of the correlation function, the behaviour at large k of the spectrum corresponds to the behaviour at small x of the correlation function. The $e^{-\alpha|x|}$ behaviour suppresses the crude ultraviolet divergence in the energy spectrum and leads to a Lorenz shape as k^{-2} (see [4]). This crude divergence occurs in case of a $\delta(x)$ behaviour of the correlation function, which corresponds to equipartition. Since the derivatives of $e^{-\alpha|x|}$

are not continuous or even defined, divergences can appear in the spectra of the derivatives of the fluid variables. This situation seems to be unavoidable for conservative continuous systems.

Dissipative fluids have a fundamentally different behaviour. It has been proved for 2-d Navier-Stokes equations that the time-independent space correlation function is smooth for all x (see [5]). This has been used recently to estimate the large k behaviour of the energy spectrum, which turns out to be exponentially decaying (see [6]). In [6] the smoothness (C^∞) of the correlation function has been assumed for the 3-d case without proof. This assumption seems plausible, however, for dissipative fluids which usually have attractors with finite (see [7]) Hausdorff dimension, though proofs in the 3-d case are missing. This suggests that they behave like a system with a finite number of degrees of freedom for which no ultraviolet-like divergences of any type can occur.

In the dissipative case, the divergences are avoided whatever the ensemble average is, as long as ergodicity is assumed, while in the conservative case the kind of averaging is important because suitable weighting among the infinite number of degrees of freedom penalizes, for instance, the large k (see [2]). The penalization is, however, suitable for certain quantities but not for others, especially those containing higher derivatives, so it cannot be guaranteed that all ultraviolet divergences are suppressed. Viscous-like dissipation seems to be the way to eliminate completely the ultraviolet problem.

- [1] H. Tasso, Phys. Lett. **96 A**, 33 (1983).
- [2] H. Tasso and K. Lerbinger, Phys. Lett. **97 A**, 384 (1983).
- [3] H. Tasso, Phys. Lett. **120 A**, 464 (1987).
- [4] H. Tasso, Z. Naturforsch. **41 a**, 1258 (1986).
- [5] C. Foias and G. Prodi, Rend. Semin. Univ. Padova **39**, 1 (1967).

- [6] C. Foias, O. Manley, and L. Sirovich, Phys. Fluids **A 2**, 464 (1990).
- [7] R. Temam, Infinite-Dimensional Dynamical Systems in Mechanics and Physics, Springer-Verlag, New York 1988.

Reprint requests to Max-Planck-Institut für Plasmaphysik – Bibliothek – zu Hdn. von Frau Hohaus, D-8046 Garching bei München.

0932-0784 / 90 / 0700-0928 \$ 01.30/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.