

Modification of the Generalized tanh Method with sech Function for Generalized Hamiltonian Equations

Bo Tian, KeYi Zhao, and Yi-Tian Gao

Department of Applied Mathematics and Physics,
Beijing University of Aeronautics and Astronautics,
Beijing 100083, China

and

Laboratory of Computational Physics,
Institute of Applied Physics and Computational Mathematics,
P.O. Box 8009, Beijing 100088, China

Z. Naturforsch. **52a**, 372 (1997);
received January 9, 1997

We modify the generalized tanh method with the sech function, so as to obtain a new class of the soliton-like solutions to a coupled set of generalized Hamiltonian equations.

The generalized tanh method and its extensions [1–3] are a tool to directly construct the tanh-profiled soliton-like solutions for certain equations of mathematical physics.

The sech function, like tanh, does not diverge at infinity either (for real arguments). We hereby try to modify the generalized tanh method by assuming that certain soliton-like solutions, physically localized, be of the form

$$u(x, t) = \sum_{l=0}^L \mathcal{A}_l(x, t) \cdot \tanh^l[\Psi(x, t)] + \sum_{j=0}^J \mathcal{B}_j(x, t) \cdot \operatorname{sech}[\Psi(x, t)] \cdot \tanh^j[\Psi(x, t)], \quad (1)$$

where L and J are the integers determined via the balance of the highest-order contributions from both the linear and nonlinear terms of the original equation, while $\mathcal{A}_l(x, t)$'s, $\mathcal{B}_j(x, t)$'s, and $\Psi(x, t)$ are differentiable functions. $\Psi(x, t)$ should be real to make sure that the soliton profiles hold.

It is noted that in the generalized tanh method, the soliton-like solutions only come from the first summation in Ansatz (1). What we do is to make the power

series more complete via the introduction of the second summation.

We now apply the above modification to a coupled set of generalized Hamiltonian equations,

$$u_t = u_x + 2v, \quad (2)$$

$$v_t = 2\varepsilon uv, \quad \text{where } \varepsilon = \pm 1, \quad (3)$$

which has an infinite number of conserved densities, the nondegenerate Hamiltonian structure, certain Bäcklund transformations and exact solutions [4–8].

We use Ansatz (1) for both $u(x, t)$ and $v(x, t)$, and then equate to zero the coefficients of like powers of $\mathcal{A}_l(x, t)$'s and $\mathcal{B}_j(x, t)$'s. For simplicity, we consider a trial $\Psi(x, t) = \sum_{j=0}^3 \Psi_j(x) t^j$, and after computerized symbolic computation end up with $\Psi(x, t) = \beta t + \gamma(x)$, where β is a real, non-zero constant, while $\gamma(x)$ is a real, differentiable function with $\gamma_x(x) \rightarrow \pm \infty$ when $x \rightarrow \pm \infty$. Thus, we obtain a new class of the soliton-like solutions as

$$u(x, t) = \frac{1}{2} \{ \pm i |\beta| \operatorname{sech}[\beta t + \gamma(x)] - \varepsilon \beta \tanh[\beta t + \gamma(x)] \}, \quad (4)$$

$$v(x, t) = \frac{\gamma_x(x) - \beta}{4} \cdot \{ \pm i |\beta| \operatorname{sech}[\beta t + \gamma(x)] \cdot \tanh[\beta t + \gamma(x)] + \varepsilon \beta \operatorname{sech}^2[\beta t + \gamma(x)] \}. \quad (5)$$

This class is different from that in [2, 7, 8]. It can also be reduced to solitary waves when $\gamma(x) = \alpha x + \delta$, where α and δ are real constants.

Acknowledgements

This work has been supported by the Outstanding Young Faculty Fellowship & the Research Grants for the Scholars Returning from Abroad, State Education Commission of China.

- [1] B. Tian and Y. T. Gao, *Computer Phys. Comm.* **95**, 139 (1996).
- [2] B. Tian and Y. T. Gao, to appear in *Appl. Math. Lett.* (1997).

- [3] B. Tian and Y. T. Gao, to appear in *Chaos, Solitons & Fractals* (1997).
- [4] G. Tu, *Phys. Lett. A* **94**, 340 (1983).
- [5] W. Ma, *Kexue Tongbao* **32**, 1003 (1987).
- [6] Z. Chen, *Comm. on Appl. Math. & Comput.* **4**, 71 (1990).
- [7] W. Ma, *J. Fudan Univ. (Natural Sci.)* **33**, 319 (1994).
- [8] Y. Li and C. Tian, *Kexue Tongbao (Lett.)* **29**, 1556 (1984).

Reprint requests and correspondence to Prof. Dr. Y.-T. Gao.

0932-0784 / 97 / 0400-0372 \$ 06.00 © – Verlag der Zeitschrift für Naturforschung, D-72072 Tübingen



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.