

# Higher Dimensional Integrable Models with Painlevé Property Obtained from (1+1)-Dimensional Schwarz KdV Equation

Li-li Chen and Sen-yue Lou<sup>a</sup>

Department of Physics, Normal College, Ningbo University, Ningbo 315211, P. R. China

<sup>a</sup> Institute of Modern Physics, Normal College, Ningbo University, Ningbo 315211, P. R. China;  
Institute of Theoretical Physics, Academia Sinica, P. O. Box 2735, Beijing 100080, P. R. China

Z. Naturforsch. **53 a**, 689–692 (1998); received March 24, 1998

Using the extended Painlevé analysis, we obtained some higher dimensional integrable models with the Painlevé property from the (1+1)-dimensional Schwarz KdV equation.

The lower-dimensional soliton phenomena have widely been studied both by theory [1] and experiment [2, 3]. However, there are many difficulties in the study of the higher dimensional soliton theory. Recently, one of the present authors pointed out that, if one wants to find some higher dimensional integrable models, equations with conformal invariance (invariance under the Möbius transformation) [4] may be the most suitable candidates [5]. Some types of general higher dimensional Schwarz equations are proved to be integrable in the sense that they can be changed to some forms with the Painlevé property (PP) [6]. Now, one of the important problems is what kind of Schwarz equations can be obtained from the known physical equations. In [7] it is also shown that higher dimensional integrable models may also be obtained from lower dimensional ones. In [8], one of the present authors has extended Conte's invariant Painlevé analysis [9] to a *noninvariant* (under the Möbius transformation) but more generalized form in the *same dimensions* to get some exact solutions of various nonlinear equations by means of a nonstandard truncation approach. In this paper we extend Conte's Painlevé analysis to a different but still *invariant* form in *higher dimensions* such that we can find many higher dimensional integrable models from a known lower dimensional integrable model, say, the Schwarz KdV (SKdV) equation

$$\{\phi; x\} + \frac{\phi_t}{\phi_x} + \lambda = 0, \quad \left(\{\phi; x\} \equiv \frac{\phi_{xxx}}{\phi_x} - \frac{3}{2} \frac{\phi_{xx}^2}{\phi_x^2}\right) \quad (1)$$

which is a variant form of the usual KdV equation.

In order to show the PP of (1), we take the transformations

$$\phi = \exp \rho, \quad \rho_x = u, \quad \rho_t = g \quad (2)$$

at first. The final equations for  $u$  and  $g$  read

$$uu_{xx} - \frac{1}{2}u^4 - \frac{3}{2}u_x^2 + gu + \lambda u^2 = 0, \quad u_t - g_x = 0. \quad (3)$$

Equation (3) comes from the consistence condition of the transformation (2). The PP of (3) is obvious because of the KdV equation possesses the PP. However, in order to get more information from the Painlevé analysis, we assume that  $u$  and  $g$  are not only the functions of the explicit space-time variables  $\{x, t\}$  but also the functions of the inner space variables  $\{y, z\}$ . It is known that, if a partial differential equation (PDE) possesses the PP, then the singular manifold  $\phi$  in the usual Painlevé expansion is arbitrary. Because of the arbitrariness of the singular manifold, one can take some different types of forms, say, Conte changes the expansion function (singular manifold  $\phi$ ) as

$$\chi \equiv \left( \frac{\phi_x}{\phi} - \frac{\phi_{xx}}{2\phi_x} \right)^{-1}$$

such that all the coefficients in the new Painlevé expansion possess the Möbius transformation invariance [10]. In order to include the inner parameters in our further results, we can take

$$\xi \equiv \left( \frac{\phi_y}{\phi} - \frac{\phi_{yy}}{2\phi_y} \right)^{-1} \quad (4)$$

as a new expansion function. Differentiating (4) with respect to  $x$  and  $t$ , respectively, we can obtain two identities:

Reprint requests to Dr. Sen-yue Lou;  
E-mail: sylou@fudan.ac.cn.

0932-0784 / 98 / 0800-0689 \$ 06.00 © – Verlag der Zeitschrift für Naturforschung, D-72072 Tübingen



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

$$\xi_x = p_1 - p_{1y}\xi + \frac{1}{2}(p_1s + p_{1yy})\xi^2, \quad (5)$$

$$\xi_t = p_0 - p_{0y}\xi + \frac{1}{2}(p_0s + p_{0yy})\xi^2, \quad (6)$$

where

$$p_0 \equiv \frac{\phi_t}{\phi_y}, p_1 \equiv \frac{\phi_x}{\phi_y}, s \equiv \{\phi; y\} \equiv \frac{\phi_{yyy}}{\phi_y} - \frac{3}{2} \frac{\phi_{yy}^2}{\phi_y^2} \quad (7)$$

are all conformal invariant, invariant under the Möbius transformation:

$$\phi \longrightarrow \frac{a\phi + b}{c\phi + d}.$$

With help of the leading order analysis of (3), we know that the functions  $\{u, g\}$  should be expanded as

$$u = \sum_{j=0}^{\infty} u_j \phi^{j-1}, \quad g = \sum_{j=0}^{\infty} g_j \phi^{j-1} \quad (8)$$

with

$$g_0^2 = p_0^2, \quad u_0 = g_0 p_1 / p_0. \quad (9)$$

Substituting (8) into (3) and using (5) and (6), we have

$$(j+1)(j-1)u_0u_j = f(u_i, g_i, i \leq j-1, p_0, p_1, s), \quad (10)$$

$$(j-1)(g_ju_0 - u_jg_0) = u_{j-1,t} - g_{j-1,x}, \quad (11)$$

where  $f$  is a complicated function of  $\{u_i, g_i, i \leq j-1, p_0, p_1, s\}$  and its derivatives. From (10) and (11), it is not difficult to see that the resonances are located at

$$j = -1, 1, 1. \quad (12)$$

The resonance at  $j = -1$  corresponds to the arbitrariness of the singularity manifold  $\xi$  (and then  $\phi$ ). At two resonances,  $j = 1$ , there are two compatibility conditions

$$-u_0^2 p_{1x} + u_0 u_{0x} p_1 - 2u_0^3 u_1 + 2u_1 u_0 p_1^2 = 0, \quad (13)$$

$$p_{1t} - p_0 p_{1y} - p_{0x} + p_1 p_{0y} = 0 \quad (14)$$

which must be satisfied. After substituting (9) and (7) into (13) and (14) one can see that (13) and (14) are satisfied identically. In the standard approach, in order to obtain the Bäcklund transformation of (3), one takes

$$\begin{aligned} u_j(u_i, g_i, i \leq j-1, p_0, p_1, s) &= 0, \\ g_j(u_i, g_i, i \leq j-1, p_0, p_1, s) &= 0, \quad j \geq 2. \end{aligned} \quad (15)$$

Because (15) is conformal invariant for all  $j \geq 2$  and the original Schwarz KdV equation (1) is integrable, we believe that every one of (15) is integrable. Even if this idea is not true for all  $j$ , it is still possible to get some higher dimensional integrable models from (15) by checking their PP for small  $j$ . Here we check the PP only for  $u_2 = 0$  which has the concrete form

$$\begin{aligned} \frac{\phi_t}{\phi_x} + \{\phi; x\} + \lambda + 3 \frac{\phi_x \phi_{xy} \phi_{yy}}{\phi_y^3} - 3 \frac{\phi_x^2 \phi_{yy}^2}{4\phi_y^4} - 3 \frac{\phi_x \phi_{xyy}}{2\phi_y^2} \\ + 3F_x - 3F^2 + 3F \left( \frac{\phi_x \phi_{yy}}{\phi_y^2} - \frac{\phi_{xx}}{\phi_x} \right) = 0, \end{aligned} \quad (16)$$

where  $F(\equiv u_1)$  is an arbitrary function. From (16), we see that the (2+1)-dimensional model will be reduced back to the original (1+1)-dimensional SKdV equation (1) when we drop the  $y$ -dependent terms or take  $y \sim x$ . Using the transformations

$$\phi = \exp \rho, \quad \rho_x = u, \quad \rho_y = v, \quad \rho_t = g, \quad (17)$$

(16) becomes

$$\begin{aligned} \frac{1}{4} u^4 v^4 - \frac{3}{4} u^4 v_y^2 - \frac{3}{2} v^4 u_x^2 + u v^4 u_{xx} - \frac{3}{2} u^3 v^2 u_{yy} \\ + 3F_x u^2 v^4 + \lambda u^2 v^4 - 3F^2 u^2 v^4 + g u v^4 \end{aligned} \quad (18)$$

$$+ 3u^3 v u_y v_y + 3u^3 F v^2 v_y - 3F u v^4 u_x = 0,$$

$$u_t = g_x, \quad (19)$$

$$v_t = g_y. \quad (20)$$

It is easy to prove the PP of (18) - (20) by taking the traditional WTC (Weiss-Tabor-Carnevale) approach [10]. However, in order to get (3+1)-dimensional models with conformal invariance and the PP, we introduce

$$\eta \equiv \left( \frac{\phi_z}{\phi} - \frac{\phi_{zz}}{2\phi_z} \right)^{-1} \quad (21)$$

as a new expansion function. Differentiating (21) with respect to  $x, y$  and  $t$ , respectively, we can obtain the three identities

$$\eta_x = P_1 - P_{1y}\eta + \frac{1}{2}(P_1S + P_{1yy})\eta^2, \quad (22)$$

$$\eta_y = P_2 - P_{2y}\eta + \frac{1}{2}(P_2S + P_{2yy})\eta^2, \quad (23)$$

$$\eta_t = P_0 - P_{0y}\eta + \frac{1}{2}(P_0S + P_{0yy})\eta^2, \quad (24)$$

where

$$P_0 \equiv \frac{\phi_t}{\phi_z}, \quad P_1 \equiv \frac{\phi_x}{\phi_z}, \quad P_2 \equiv \frac{\phi_y}{\phi_z}, \quad S \equiv \{\phi; z\}. \quad (25)$$

By means of the leading order analysis of (18) - (20), we know that the functions  $\{u, v, g\}$  should be expanded as

$$u = \sum_{j=0}^{\infty} u_j \phi^{j-1}, \quad v = \sum_{j=0}^{\infty} v_j \phi^{j-1}, \quad g = \sum_{j=0}^{\infty} g_j \phi^{j-1} \quad (26)$$

with

$$g_0^2 = P_0^2, \quad u_0 = P_1 \frac{g_0}{P_0}, \quad v_0 = P_2 \frac{g_0}{P_0}. \quad (27)$$

Substituting (26) into (18) - (20) and using (22) - (24), we have

$$(j+1)(j-1)(u_0u_j + v_0v_j) \quad (28)$$

$$= f(u_i, g_i, i \leq j-1, P_0, P_1, P_2, S),$$

$$(j-1)(g_ju_0 - u_jg_0) = u_{j-1,t} - g_{j-1,x}, \quad (29)$$

$$(j-1)(g_jv_0 - v_jg_0) = v_{j-1,t} - g_{j-1,y}. \quad (30)$$

From (28) - (30) it is not difficult to see that the resonances are located at

$$j = -1, 1, 1, 1. \quad (31)$$

The three compatibility conditions at the resonances,  $j = 1$ , now read

$$\begin{aligned} G = & \left( -6u_1 + 6\frac{\phi_x}{\phi_y}v_1 \right) F + 3u_{1x} - 3u_1^2 + \frac{3}{2}\frac{\phi_x^2}{\phi_y^2}v_1^2 - \frac{3}{2}\frac{\phi_x}{\phi_t}u_{1t} + 3\frac{\phi_x}{\phi_y}u_1v_1 + \frac{3}{2}\frac{\phi_x^2}{\phi_t\phi_y}v_{1t} - \frac{3}{2}\frac{\phi_x^2}{\phi_t\phi_y}g_{1y} \\ & - 3\frac{\phi_x}{\phi_y}u_{1y} - \frac{3}{2}\phi_x^2\phi_y^2v_{1y} - 3\frac{\phi_{xx}}{\phi_x}u_1 - 3\frac{\phi_x^2\phi_{zz}}{\phi_z^2\phi_y}v_1 + \frac{3}{2}\frac{\phi_x}{\phi_t}g_{1x} + 3\frac{\phi_x\phi_{xy}}{\phi_y^2}v_1 + \frac{3}{2}\frac{\phi_x\phi_{zz}}{\phi_z^2}u_1 \\ & + \frac{3}{2}\frac{\phi_x\phi_{yy}}{\phi_y^2}u_1 + 3F_x - 3F^2 + 3\left(\frac{\phi_x\phi_{yy}}{\phi_y^2} - \frac{\phi_{xx}}{\phi_x}\right)F = 0. \end{aligned} \quad (37)$$

$$-3u_0^3u_1v_0^2P_2^2 + \frac{3}{2}u_0^4v_0^2P_{2y} + u_0^4v_0^3v_1 + 3Fu_0^2v_0^4P_1$$

$$+ 2v_0^3v_1u_0^2P_1^2 - 3u_0^4v_0v_1P_2^2 - 3Fu_0^3v_0^3P_2 \quad (32)$$

$$- \frac{3}{2}u_0^4v_{0y}v_0P_2 + u_0^3u_1v_0^4 = 0,$$

$$u_0P_{0z} - g_{0x} + u_{0t} - g_0P_{1z} = 0, \quad (33)$$

$$v_0P_{0z} - g_{0y} + v_{0t} - g_0P_{2z} = 0, \quad (34)$$

and these equations are satisfied identically because of (25) and (27). So the (2+1)-dimensional system (18) - (20) (and then its Schwarz form (16)) is integrable under the meaning that it possesses the PP. In the same way, if we take

$$u_j = u_j(P_0, P_1, P_2, S, u_1, v_1, g_1) = 0,$$

$$v_j = v_j(P_0, P_1, P_2, S, u_1, v_1, g_1) = 0, \quad (35)$$

$$g_j = g_j(P_0, P_1, P_2, S, u_1, v_1, g_1) = 0, \quad (j \geq 2)$$

we can get the Bäcklund transformation for the system (18) - (20). Furthermore, it is interesting that every one of the equations (35) for a fixed  $j$  may be integrable because all these equations are conformal invariant and are derived from the integrable equation (16). The simplest one of them is given by  $u_2 = 0$ , i. e.,

$$\begin{aligned} & \frac{\phi_t}{\phi_x} + \{\phi; x\} + \lambda + \frac{9}{2}\frac{\phi_x\phi_{zz}\phi_{xz}}{\phi_z^3} - \frac{3}{4}\frac{\phi_x^2\phi_{yy}^2}{\phi_y^4} \\ & + \frac{3}{8}\frac{\phi_x^2\phi_{zz}^2}{\phi_z^4} + 3\frac{\phi_x^2\phi_{yzz}}{\phi_z^2\phi_y} - \frac{3}{2}\frac{\phi_x\phi_{xyy}}{\phi_y^2} + 3\frac{\phi_x\phi_{xy}\phi_{yy}}{\phi_y^3} \\ & - \frac{9}{4}\frac{\phi_x\phi_{xzz}}{\phi_z^2} - 6\frac{\phi_x^2\phi_{yz}\phi_{zz}}{\phi_z^3\phi_y} + G = 0, \end{aligned} \quad (36)$$

where the arbitrary function  $G$  is related to other arbitrary functions  $F, u_1, v_1$  and  $g_1$  by

If we drop  $y$  and  $z$  related terms or if fixed  $y$  and  $z$  are proportional to  $x$ , we reobtain the (1+1)-dimensional SKdV equation (1) from the (3+1)-dimensional equation (36).

We have proved the PP of (36) by using the Painlevé analysis and that some (4+1)-dimensional integrable models with the PP can be obtained when we introduce a further inner parameter into the procedure of proving the PP of the (3+1)-dimensional model (36). One of the simplest models obtained in this way reads

$$\begin{aligned} \frac{\phi_t}{\phi_x} + \{\phi; x\} + \lambda - \frac{3}{4} \frac{\phi_x^2 \phi_{yy}^2}{\phi_y^4} + \frac{9}{2} \frac{\phi_x \phi_{zz} \phi_{xz}}{\phi_z^3} - \frac{9}{4} \frac{\phi_x \phi_{xz}^2}{\phi_z^2} + 3 \frac{\phi_x \phi_{xy} \phi_{yy}}{\phi_y^3} + \frac{3}{8} \frac{\phi_x^2 \phi_{zz}^2}{\phi_z^4} - \frac{9}{2} \frac{\phi_x^2 \phi_{x_4 x_4} \phi_{y x_4}}{\phi_{x_4}^3 \phi_y} \\ + \frac{9}{4} \frac{\phi_x^2 \phi_{y x_4 x_4}}{\phi_{x_4}^2 \phi_y} - 6 \frac{\phi_x^2 \phi_{yz} \phi_{zz}}{\phi_z^3 \phi_y} - \frac{3}{16} \frac{\phi_x^2 \phi_{x_4 x_4}^2}{\phi_{x_4}^4} + 3 \frac{\phi_x \phi_{x x_4} \phi_{x_4 x_4}}{\phi_{x_4}^3} + \frac{9}{4} \frac{\phi_x^2 \phi_{x_4 x_4} \phi_{z x_4}}{\phi_{x_4}^3 \phi_z} - \frac{9}{8} \frac{\phi_x^2 \phi_{z x_4 x_4}}{\phi_{x_4}^2 \phi_z} \\ + 3 \frac{\phi_x^2 \phi_{y z z}}{\phi_z^2 \phi_y} - \frac{3}{2} \frac{\phi_x \phi_{x x_4 x_4}}{\phi_{x_4}^2} - \frac{3}{2} \frac{\phi_x \phi_{x y y}}{\phi_y^2} = 0. \end{aligned} \quad (38)$$

Equation (38) is a (4+1)-dimensional generalization of the SKdV (1). If we use some reductions on (38), say,  $x_4 = z$ ,  $x_4 = y$ , ..., we can get some further (3+1)-dimensional SKdV equations.

Using the above procedure step by step, we can get various integrable models with the PP in any dimensions.

In summary, after introducing some inner parameters explicitly in the expansion variables, we

can get many higher dimensional integrable models from lower dimensional ones. Some concrete SKdV equations with the PP in (2+1)-, (3+1)- and (4+1)-dimensions are listed here for further investigations.

The work was supported by the National Nature Science Foundation of China and the Nature Science foundation of Zhejiang province in China. One of the authors (Lou) thanks Professors Q.-p. Liu, X.-b. Hu, J.-j. Xu and G.-j. Ni for their helpful discussions.

- [1] M. J. Ablowitz and P. A. Clarkson, *Solitons Nonlinear Evolution Equations and Inverse Scattering*, London Mathematical Society Lecture Note Series 149 (Cambridge University Press 1991).
- [2] Y. S. Kivshar and B. A. Malomed, *Rev. Mod. Phys.* **61**, 763 (1989) and references therein.
- [3] S.-y. Lou and Guoxiang Huang, *Mod. Phys. Lett.* **B9**, (1995) 1231 (1995); S.-y. Lou, J. Yu, J. Lin, Guoxiang Huang, and J.-k. Zhang, *Mod. Phys. Lett.* **B10**, 11 (1996).
- [4] S.-y. Lou, *J. Phys. A: Math. Gen.* **30**, 4803 (1997).
- [5] S.-y. Lou, *Science in China (series A)*, **34**, 1317 (1997).
- [6] S.-y. Lou, *J. Math. Phys.* **31**, 2112 (1998); S.-y. Lou, *Commun. Theor. Phys.* **28**, 129 (1997).
- [7] S.-y. Lou, *J. Phys. A: Math. Gen.* **30**, 4803 (1997).
- [8] S.-y. Lou, *Z. Naturforsch.* **53a**, 251 (1998).
- [9] R. Conte, *Phys. Lett.* **A140**, 383 (1989).
- [10] J. Weiss, M. Tabor, and G. Carnevale, *J. Math. Phys.* **24**, 522 (1983).