

A New Type of Chaotic Attractor

Normann Kleiner, Sebastian Fischer, and Dieter Fröhlich

Division of Theoretical Chemistry, University of Tübingen
Auf der Morgenstelle 8, D-72076 Tübingen

Z. Naturforsch. **53a**, 265 (1998); received February 24, 1998

An experimentally discovered inverted spiral-type chaotic attractor is reproduced by a model equation.

Does there exist a simple equation for the attractor recently found both in an electronic and a hydrodynamical system [1]? After a first, unpublished attempt by Sven Sahle to mirror a classical spiral-type attractor using a tube put into the middle, which yielded "messy" equations, a fairly simple ordinary differential equation (ODE) was found and will be presented in the following.

An experimental result is reproduced in Figure 1. It

shows a typical inverted spiral-type attractor obtained in the hydrodynamical system described in [1].

The 3-variable ODE:

$$\begin{aligned}\dot{x} &= -y + z, \\ \dot{y} &= x - ay, \\ \dot{z} &= b - (x^2 + y^2)z + cz.\end{aligned}\quad (1)$$

Figures 1 and 2 show an analogous type of chaotic attractor. The main point in common is that the spiral is traversed "from the outside to the inside", in contrast to the attractor described in 1976 by Rössler [2].

Two questions are currently open:

1) Can the equations be simplified (so that eventually perhaps only a single quadratic term is left)?

2) Is a Shil'nikov flow, with a homoclinic trajectory emerging from a central saddle focus [3], as it has been observed experimentally by N. Kleiner (work in preparation), reproducible by a similarly simple equation?

To conclude, we have presented a simple 3-variable ODE showing a prototype "inverted spiral attractor".

We thank Sven Sahle and Otto E. Rössler for discussions.




Fig. 1. Experimental attractor obtained with a hydrodynamical system, cf. [1]




Fig. 2. Numerical simulation of (1), using $a = 0.1$, $b = 0.1$, and $c = 4.2$.

Reprint requests to S. Fischer;
E-mail: sebastian.fischer@student.uni-tuebingen.de

- [1] N. Kleiner, Das deterministische Chaos im Experiment, in: Komplexität und Selbstorganisation, ed. by H. Krapp and T. Wägenbaur (Fink, München 1997), p. 259.
[2] O. E. Rössler, An equation for continuous chaos, Phys. Lett. A **57**, 397 (1976).

- [3] L. P. Sil'nikov, A contribution to the problem of the structure of an extended neighborhood of a structurally stable equilibrium of saddle-focus type, Math. USSR Sbornik **10**, p. 91 (1970).

0932-0784 / 98 / 0500-0265 \$ 06.00 © – Verlag der Zeitschrift für Naturforschung, D-72027 Tübingen



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

