

Dielectric Relaxation of Hexadeutero Dimethylsulfoxide

H. Betting and M. Stockhausen

Institut für Physikalische Chemie der Universität Münster,
Schlossplatz 4/7

Reprint requests to Prof. M. St.; Fax: (+0251)8323441.

Z. Naturforsch. **54a**, 661 (1999);
received September 7, 1999

The dielectric relaxation parameters of the title substance (DMSO- d_6) in its pure liquid state are determined from measurements up to 72 GHz at 20 °C in comparison to protonated DMSO. While the relaxation strengths do not differ, the relaxation time of DMSO- d_6 is significantly longer (21.3 ps) than that of DMSO (19.5 ps).

Key words: Dielectrics; Isotopic Substitution; Liquids; Relaxation.

Dimethylsulfoxide (DMSO) is a liquid exhibiting some peculiar physical properties which point to specific intermolecular interactions, the nature of which is a matter of continuing debate. Such a peculiarity is also found in the dielectric relaxation behaviour inasmuch as the relaxation time is longer than to be expected from a comparison with other liquids consisting of quasi-rigid molecules for which the main relaxation contribution is due to the reorientational tumbling motion of single molecules [1]. Thus some kind of transient associative effect is likely to be operative, on average slowing down the reorientational motion (note that, in principle, dielectric spectroscopy observes molecular collectives rather than single molecule motion). In that context it seems worthwhile to regard the alteration of the relaxation behaviour caused by isotopic substitution $H \rightarrow D$, thus comparing DMSO with DMSO- d_6 . For a small, rigid molecule like water it is known that deuteration leads to a noticeable increase in relaxation time τ [2]: At 20 °C, the ratio for deuterated/protonated water is $\tau(D)/\tau(H)=1.25$. Concerning DMSO, the far-infrared absorption spectra of DMSO and DMSO- d_6 have been studied with respect to the libra-

tional modes effective in that region [3], but such a comparison has not yet been published for the "classical" dielectric relaxation process, viz. the microwave region. Therefore we report here the respective relaxation data.

We have studied both DMSO and DMSO- d_6 (Merck, deuteration >99%) at 20 °C over the frequency range between 90 MHz and 72 GHz, paying particular attention to the differences in the (negative) imaginary part of permittivity, ϵ'' , in dependence on the frequency ν . In both cases, the $\epsilon''(\nu)$ spectra can satisfactorily be fitted by a Debye function, disregarding the indication of an additional but very weak contribution on the high frequency side of the spectrum. The parameters of the main relaxation region, that is the relaxation time τ and relaxation strength S , are given in Table 1.

Table 1. Relaxation time τ and relaxation strength S for DMSO and DMSO- d_6 at 20 °C.

	τ ps	S
DMSO	19.5	44.2
DMSO- d_6	21.3	44.2

In view of the only slight increase of molar concentration on deuteration (less than 0.5 percent) it is not unexpected that the relaxation strength S stays practically unaltered. The relaxation time τ , on the other hand, is significantly increased: $\tau(D)/\tau(H)=1.09$. It is interesting to compare that finding with NMR results from which the correlation time τ_c of the single molecule tumbling motion can be obtained. The respective ratio is $\tau_c(D)/\tau_c(H)=1.12$ [4]. Even taking into account experimental uncertainties, this value seems to be slightly larger than $\tau(D)/\tau(H)$. If so, this would be not inconsistent with the above-mentioned conjecture that the dielectrically observed process comprises some kind of associative effect, which here may cause a reduction of the collective characteristic $\tau(D)/\tau(H)$ in comparison to the single molecule isotopic correlation time ratio.

- [1] G. Turkey, G. Wilke, U. Witt, A. Ghoneim, and M. Stockhausen, Proc. 3rd Conf. Dielectric and Related Phenomena, Zakopane 1994, p. 146.
- [2] P. S. Yastremskii, Zh. Strukt. Khim. **12**, 532 (1971); E. H. Grant and R. Shak, Trans. Faraday Soc. **65**, 1519 (1969).

- [3] I. W. Larkin, J. Chem. Soc. Faraday Trans. II **70**, 1457 (1974).
- [4] Mao Xi'an and M. Holz, Sci. China Ser. B **38**, 393 (1995); M. Holz, Xi-an Mao, D. Seiferling, and A. Sacco, J. Chem. Phys. **104**, 669 (1996).

0932-0784 / 99 / 1000-0661 \$ 06.00 © Verlag der Zeitschrift für Naturforschung, Tübingen · www.znaturforsch.com



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.