## Properties of Nanocrystalline Nickel Particles in Ni-SiO<sub>2</sub> Composites

G. Ennas, A. Falqui<sup>a</sup>, G. Piccaluga, S. Solinas, D. Gatteschi<sup>b</sup>, C. Sangregorio<sup>b</sup>, and A. Benedetti<sup>c</sup>

Dipartimento di Scienze Chimiche, Università di Cagliari, 09100 Cagliari, Italy

Reprint requests to Prof. G. E; Fax: +39 070 6754388, E-mail: ennas@unica.it

Z. Naturforsch. 55 a, 581–588 (2000); received March 9, 2000

Nickel-Silica nanocomposites with a nickel content equal to 10, 15, 20 wt% have been prepared by a sol-gel method starting from ethanolic solutions of tetraethoxysilane and nickel nitrate hexahydrate. After gelation the samples were reduced in  $H_2$  flow at selected temperature (450 °C < T < 600 °C).

The morphological, structural and magnetic properties were investigated by transmission electron microscopy (TEM), wide and small angle X-ray scattering (WAXS, SAXS), magnetic susceptibility in zero field cooled and field cooled mode (ZFC and FC), and magnetic hysteresis loop.

Nanometric nickel particles are observed in all the investigated samples. TEM, WAXS and SAXS

the nickel concentration. TEM results moreover indicate that also the width of the particle size distribution, which can be simulated by log-normal functions, follows this trend.

All the sample treated in hydrogen show superparamagnetic behaviour. The magnetisation falls to reach saturation up to highest measuring field of 70 kOe even at 3 K, while the observed

techniques indicate that the average nickel particle size grows slightly but almost regularly with

to reach saturation up to highest measuring field of 70 kOe even at 3 K, while the observed coercivity  $H_c$  is much higher than the theoretical bulk one. Some uncertainty in the complete interpretation of the sequence of magnetic measurements is attributed to a progressive oxidation of the samples when these are air exposed.

Key words: Nanocomposites; Nickel-Silica; X-ray Diffraction; TEM; Magnetic Properties.

<sup>&</sup>lt;sup>a</sup> Dipartimento di Fisica, Università di Cagliari, Italy

<sup>&</sup>lt;sup>b</sup> Dipartimento di Chimica, Università di Firenze, Italy

<sup>&</sup>lt;sup>c</sup> Dipartimento di Chimica Fisica, Università di Venezia, Italy