(2+1)-Dimensional Davey-Stewartson II Equation for a Two-Dimensional Nonlinear Monatomic Lattice

Zhi-Fang Li^a and Hang-Yu Ruan^{a,b,c}

- ^a Department of Physics, Ningbo University, Ningbo 315211, China
- ^b Nonlinear Science Center, Ningbo University, Ningbo 315211, China
- ^c State Key Laboratory of Scientific and Engineering Computing, Institute of Computational Mathematics and Scientific Engineering Computing, Academy of Mathematics and System Sciences, the Chinese Academy of Sciences, P.O. Box 2719, Beijing 100080, China

Reprint requests to H.-Y.R.; E-mail: hyruan@mail.nbip.net

Z. Naturforsch. **61a**, 45 – 52 (2006); received October 12, 2005

A two-dimensional monatomic lattice with nearest-neighbor interaction is investigated by the method of multiple scales combined with a quasidiscreteness approximation. The Davey-Stewartson II equation (DS-II) is obtained from the original two-dimensional (2D) differential-difference system. By solving the DS-II, explicit periodic solutions, soliton solutions and rational function solutions are obtained, and the leading order approximated solutions of the 2D monatomic lattice are constructed by explicit solutions of the DS-II.

Key words: Two-Dimensional Monatomic Lattice; Method of Multiple Scales; Quasi-Discreteness Approximation; Davey-Stewartson II Equation.