

NOTIZEN

***n*-Paraffins from the Flowers and Leaves of *Parabenzoin praecox* (Sieb. et Zucc.) Nakai (Lauraceae)**

N. Hayashi and H. Komae

Department of Chemistry, Faculty of General Education,
Hiroshima University, Hiroshima, Japan

(Z. Naturforsch. **28 c**, 772 [1973] ; received July 3, 1973)

P. praecox (Sieb. et Zucc.) Nakai (Lauraceae), flowers and
leaves, *n*-paraffins (C₁₇ to C₃₄)

In the course of the chemotaxonomical study of
Lauraceae family, *n*-paraffin constituents of the
leaves and flowers of *P. praecox* were investigated.

Flowers. The flowers (12 g), collected on March
31st at Hiroshima Prefecture in Japan, were ex-
tracted with *n*-hexane (500 ml) for one week. The
n-paraffin fraction (5.7 mg) was isolated from
the extractives by means of column chromatography
over *n*-hexane and then by Molecular Sieb 5A
treatment.

The following *n*-paraffins were detected by gas
chromatography (SE-30 10% on Chromosorb W,
temperature programmed 190 to 320 °C, 2 °C/
min): *n*-Paraffins C₁₇ to C₁₉ trace, C₂₀ 0.1%, C₂₁
11.6%, C₂₂ 0.7%, C₂₃ 30.1%, C₂₄ 1.6%, C₂₅ 25.8%,
C₂₆ 0.7%, C₂₇ 14.7%, C₂₈ 0.3%, C₂₉ 13.3%, C₃₀
0.1%, C₃₁ 1.0%. Odd paraffins : even paraffins =
96.5 : 3.5.

Leaves. The leaves (250 g) which collected on
May 25th in Hiroshima Prefecture were extracted
with *n*-hexane for one week. The following *n*-paraf-
fins (479.8 mg) were detected by gas chromato-
graphy as the same method mentioned above:
n-Paraffins C₂₁ 0.2%, C₂₂ 0.3%, C₂₃ 1.4%, C₂₄ 0.5%,
C₂₅ 3.8%, C₂₆ 0.7%, C₂₇ 19.1%, C₂₈ 1.0%, C₂₉
55.1%, C₃₀ 1.3%, C₃₁ 16.2%, C₃₂ trace, C₃₃ 0.4%,
C₃₄ trate. Odd paraffins : even paraffins = 96.2 : 3.8

The *n*-paraffins of *P. praecox* are similar to those
of other Lauraceae plants^{1, 2} as shown in Table I.
The main constituent of the leaves is C₂₉ paraffin
and that of the flowers is C₂₃ paraffin.

Table I. *n*-Paraffin constituents of lauraceae plants. A: *Cinnamomum camphora* Sieb. B: *Lindera obtusiloba* Blume. C: *Lit-
sea japonica* (Thunb.) Juss. D: *Lindera umbellata* Thunb. E: *Parabenzoin praecox* (Sieb. et Zucc.) Nakai. F: *Lindera umbel-
lata* Thunb.-flowers. G: *Parabenzoin praecox* (Sieb. et Zucc.) Nakai-flowers.

<i>n</i> -Paraffins	Constituents [%]*						
	A	B	C	D	E	F	G
C ₁₆	trace	1.1	—	0.1	—	—	—
C ₁₇	trace	0.3	—	0.2	—	—	trace
C ₁₈	trace	1.3	—	0.3	—	—	trace
C ₁₉	trace	0.4	trace	0.3	—	—	trace
C ₂₀	0.2	1.3	trace	0.6	—	0.3	0.1
C ₂₁	0.5	0.4	trace	0.3	0.2	36.5	11.6
C ₂₂	0.8	1.3	trace	0.6	0.3	2.8	0.7
C ₂₃	2.6	1.4	0.6	1.3	1.4	46.7	30.1
C ₂₄	2.4	1.3	2.1	1.2	0.5	0.7	1.6
C ₂₅	7.1	2.9	3.4	4.5	3.8	8.5	25.8
C ₂₆	—	1.9	7.0	1.5	0.7	—	0.7
C ₂₇	23.9	25.6	14.4	18.8	19.1	2.8	14.7
C ₂₈	4.3	1.6	18.8	1.5	1.0	—	0.3
C ₂₉	43.3	45.1	23.8	53.3	53.1	0.9	13.3
C ₃₀	1.2	1.6	14.2	1.3	1.3	0.05	0.1
C ₃₁	8.8	11.5	14.8	15.0	16.2	0.25	1.0
C ₃₂	trace	trace	trace	—	trace	—	—
C ₃₃	trace	trace	0.9	—	0.4	—	—
C ₃₄	trace	trace	—	—	trace	—	—

* The percentage of the constituents of the oils was calculated from the areas of the peaks of the gas chromatogram.

Requests for reprints should be sent to Dr. Nanao Hayashi,
Department of Chemistry, Faculty of General Education,
Hiroshima University, Hiroshima, Japan.

¹ H. Komae and N. Hayashi, Phytochem. **10**, 2834 [1971].
² N. Hayashi and H. Komae, Z. Naturforsch. **27 b**, 1438
[1972].

