

The Effect of D-Amphetamine on the Concentration of Phospholipids in the Rat Hippocampus

B. E. Leonard * and V. Neuhoff

Max-Planck-Institut für Experimentelle Medizin,
Arbeitsgruppe Neurochemie, Göttingen

(Z. Naturforsch. **29** c, 187–188 [1974]; received
December 12, 1973)

Amphetamine, Phospholipids, Hippocampus

It is generally accepted that the behaviourally stimulant properties of D-amphetamine can be attributed to its ability to release catecholamines from intraneuronal storage sites and block their reuptake^{1, 2}. Since lipids constitute an essential part of cellular membranes³, and there is some evidence that they may be involved in ion transport processes in nerve membranes⁴, it seems possible that amphetamine could be producing its effect on amine metabolism by a direct action on membrane phospholipid metabolism. The present investigation was designed to assess this possibility.

The fluorometric microdetermination of Schiefer and Neuhoff⁵ was used to determine the concentration of phospholipids in the hippocampus. This brain region was studied because it could be clearly and reproducibly identified, rapidly dissected from surrounding brain regions and contained sufficient tissue for multiple determinations to be made.

Groups of 5 adult rats (♂, 180–200 g) were injected with D-amphetamine (10 mg/kg i.p.) or physiological saline (control group) and killed by decapitation 1 hour later. At the time of killing, the amphetamine treated rats were hyperactive; there is evidence from our previous studies that the drug produces its maximal effect on brain amine metabolism after this period of treatment⁶. The brains were rapidly removed, placed on a cold dish, the hippocampi rapidly dissected and frozen on solid

carbon dioxide. Small pieces of tissue, weighing approximately 1 mg, were removed by means of a metal punch⁷ and the phospholipids extracted by chloroform : methanol (2 : 1). The extract was then submitted to two-dimensional chromatography on silica gel microplates and the phospholipids visualized with iodine vapour. The exact position of phosphatidyl choline, phosphatidyl ethanolamine, sphingomyelin, phosphatidyl serine and phosphatidyl inositides could then be located, removed with a micro spatula and estimated fluorometrically using rhodamine 6 GO. This procedure has been described in detail elsewhere^{5, 7}. At least 4 separate estimations of the phospholipid content were made on each hippocampus.

The results are given in Table I. In this investigation it can be seen that, in the hippocampus, D-amphetamine produces a significant decrease in the concentration of phosphatidyl choline and an increase in that of phosphatidyl ethanolamine. The slight rise in the total phospholipid concentration can be partly attributed to the rise in phosphatidyl ethanolamine as the summation of the concentrations of the individual phospholipids as a percentage of the total phospholipid is the same in both groups (54%). The reason for the increase in phosphatidyl ethanolamine must await further investigation but preliminary results show that D-amphetamine significantly reduces (approx. 30%) the concentration of ethanolamine in the hippocampus⁸. It thus seems possible that the turnover of phosphatidyl ethanolamine is increased by D-amphetamine.

Hitzemann and Loh⁹ have recently shown that the same dose of D-amphetamine inhibits the synthesis of phosphatidyl choline in the rat brain, probably by inhibiting the cytidine diphosphorylcholine diglyceride transferase activity. These investigators also found that the turnover of phosphatidyl choline was

Table I. Effect of D-amphetamine on the concentration of phospholipids in the rat hippocampus.

Group	Total P. lipid	P'tidyl choline	P'tidyl serine	P'tidyl inositides	P'tidyl ethanolamine
Controls (N=5)	2.93 ± 0.29	0.726 ± 0.064	0.032 ± 0.007	0.032 ± 0.006	0.758 ± 0.23
Amphetamine (N=5)	3.88 ± 0.32	0.468 ± 0.072	0.050 ± 0.008	0.028 ± 0.006	1.52 * ± 0.20

All values, as the Mean ± s.e.m., are expressed as mg phospholipid/g wet weight of tissue. * P<0.05.

* Present address: Pharmacology Department, Organon Scientific Development Group, Oss, The Netherlands.

Requests for reprints should be sent to Prof. Dr. V. Neuhoff, Max-Planck-Institut für Experimentelle Medizin, Arbeitsgruppe Neurochemie, D-3400 Göttingen, Hermann-Rein-Str. 3.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

reduced in several regions of the rat brain but they did not study the effects of this drug on the metabolism of this phospholipid in the hippocampus.

The concentration of sphingomyelin in the hippocampus is very low in comparison to that of the other phospholipids which were estimated. Reliable and reproducible results for the concentration of

this phospholipid in the hippocampus were not obtained.

In conclusion, these and the results of other investigators⁹, suggest that the D-amphetamine has an effect on membrane phospholipid metabolism; this could play a contributory role in the effect of this drug release and reuptake of neurotransmitter substance in the brain.

¹ A. Carlsson, M. Lindquists, A. Dahlström, K. Fuxe, and D. Masuoka, *J. Pharmacy Pharmacol.* **17**, 521 [1965].

² L. L. Iversen, J. Axelrod, and J. Glowinski, *J. Proc. 5th Int. Congr.: Coll. Inr. Neuropsychopharmac.*, p. 362, Washington 1967.

³ W. Stoeckenius and D. M. Engelman, *J. Cell Biol.* **42**, 613 [1969].

⁴ L. E. Hokin and M. R. Hokin, *Proc. nat. Acad. Sci. USA* **53**, 1369 [1965].

⁵ H. G. Schiefer and V. Neuhoff, *Hoppe-Seyler's Z. physiol. Chem.* **352**, 913 [1971].

⁶ B. E. Leonard and S. A. Shallice, *Br. J. Pharmacol. Chemotherapy* **41**, 198 [1971].

⁷ V. Neuhoff, *Molecular Biology, Biochemistry and Biophysics*, eds. A. Kleinzeller, G. F. Springer, H. G. Wittmann, **Vol. 14**, pp. 149–178, Springer-Verlag, Berlin, Heidelberg, New York 1973.

⁸ B. E. Leonard, unpublished observation.

⁹ R. J. Hitzemann and H. H. Loh, *Biochem. Pharmacol. Chemotherapy* **22**, 2731 [1973].