

## Notizen

# NMR Relaxation Study of the Thermal Denaturation of Lysozyme

D. Pumpernik, B. Fajt, S. Lapanje, and A. Azman

Chemical Institute Boris Kidrič and Department of Chemistry, University of Ljubljana, Ljubljana

(Z. Naturforsch. **30 c**, 294–295 [1975]; received November 25, 1974)

Relaxation, Lysozyme, Thermal Denaturation

Lysozyme relaxation times  $T_1$  and  $T_2$  have been measured. Methyl groups  $T_1$  are sensitive to thermal denaturation of the lysozyme. The exchange of the water molecules between bounded and free is slow.

The thermal denaturation of lysozyme was very extensively studied<sup>1</sup> by various methods including high resolution NMR spectroscopy<sup>2</sup>. The purpose of this contribution is to report the results of NMR measurements of the proton spin-lattice ( $T_1$ ) and spin-spin ( $T_2$ ) relaxation times. The measurements have been made on lysozyme dissolved in  $D_2O$  and  $H_2O$ . Salt-free egg-white lysozyme (6 x cryst., Lot 7102) was supplied by Miles Laboratories, Inc. The concentration of lysozyme was in both cases 10% and pD 5.6. The relaxation times were measured on a conventional Bruker pulse spectrometer B-KR 322 s.

## Lysozyme in $D_2O$

A. The measured spin-lattice relaxation time can be accurately separated into three contributions. In Figs 1 and 2, two of them are depicted. The third is temperature independent with the value 1.6 sec. The dominant relaxation mechanism for the protons is the dipole-dipole one. From the minimum of  $T_1^1$  one can calculate the reorientational correlation time  $\tau = 9.8 \times 10^{-10}$  sec and the proton-proton distance  $r = 2 \text{ \AA}$ . Near the minimum of  $T_1^1$  the activation energy has the value 2.24 kcal/mol. All these three numbers suggest that the main contributors to the  $T_1^1$  minimum are the methyl groups.

The contributions of other groups to  $T_1^1$  are not negligible. The shape of  $T_1^1$  is not symmetrical around the minimum, and the activation energy far from it has a value of approximately 12 kcal/mol which strongly indicates that larger groups than the methyl ones influence the relaxation time. The

Requests for reprints should be sent to Dr. A. Azman, Department of Chemistry, University of Ljubljana, Mur-nikova 6 – P.O.B. 537, 61001 Ljubljana, Yugoslavia.




Fig. 1. Spin-lattice relaxation time  $T_1^1$  of lysozyme in  $D_2O$ .




Fig. 2. Spin-lattice relaxation time  $T_1^3$  of lysozyme in  $D_2O$ .

methyl groups as monitored by  $T_1^1$  are a sensitive indicator of pretransition changes occurring in the thermal transition<sup>3</sup>.  $T_1^3$  results from the constituent motions that are not activated in the thermal transition.

B. The spin-spin relaxation time  $T_2$  can be separated into two contributions, Figs 3 and 4. From the temperature dependence of  $T_2^1$  the conclusion can be drawn that the main contributions stem from larger groups whose ordering in lysozyme is constant between  $T = 320$  and  $T = 340$  K. At higher temperatures motions of these groups are uncooperative like in simple liquids with  $T_2$  approaching  $T_1$ . The maximum of  $T_2^2$  is at the same temperature as the minimum of  $T_1^1$ . From the known




Fig. 3. Spin-spin relaxation time  $T_2^1$  of lysozyme in  $D_2O$ .



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.




Fig. 4. Spin-spin relaxation time  $T_2^2$  of lysozyme in  $D_2O$ .

expression<sup>4</sup> for  $T_2$  and from the data calculated from  $T_1$ <sup>1</sup>, the  $T_2^2$  can be reproduced in the temperature interval  $T = 315 - 335$  K. Thus the re-orientational motions of methyl groups contribute to  $T_2^2$  in this temperature interval.

#### Lysozyme in $H_2O$

Measurements of  $T_1$  ( $T_2$ ) of lysozyme in  $H_2O$  have given only one value for  $T_1$  ( $T_2$ ), Figs 5 and 6. The relaxation times  $T_1$  and  $T_2$  are due to the proton motion of water. From the difference between  $T_1$  of lysozyme in  $H_2O$  and pure water the fraction of the "immobilized" water at 25 °C is estimated to be about 14%. This is the upper limit if the assumption is made that  $T_1$  of the bound water is much smaller than that of the free one. It is interesting that this is not supported by measurements of the diffusion coefficients. Within the experimental error, the diffusion coefficients of pure water and the water in lysozyme-water solution are the same. The sensitivity of  $T_1$  and the diffusion coefficients to the amount of the bound water is apparently not the same.




Fig. 5. Spin-lattice relaxation time  $T_1$  of lysozyme in  $H_2O$ .




Fig. 6. Spin-spin relaxation time  $T_2$  of lysozyme in  $H_2O$ .

The spin-spin relaxation time  $T_2$  decreases with temperature. This behaviour suggests that the exchange of the "free" and "bound" water molecules is slow<sup>5</sup>. The difference  $(1/T_2 - 1/T_1)$  increases logarithmically with temperature, the exchange energy being 3.9 kcal/mol.

This work was supported by the Boris Kidrič Fund.

<sup>1</sup> T. Imoto, L. N. Johnson, A. C. T. North, D. C. Phillips, and J. A. Rupley, *The Enzymes* (P. B. Boyer, ed.), Vol. VII, Chapter 21, Academic Press, New York and London 1973.

<sup>2</sup> C. C. McDonald, W. D. Phillips, and D. D. Glickson, *J. Amer. Chem. Soc.* **93**, 235 [1971].

<sup>3</sup> N. N. Khechinashvili, P. L. Privalov, and E. I. Tiktopulo, *FEBS Lett.* **30**, 57 [1973].

<sup>4</sup> A. Abragam, *The Principles of Nuclear Magnetism*, Oxford University Press, London 1961.

<sup>5</sup> B. D. Sykes, P. G. Schmidt, and G. R. Stark, *J. Biol. Chem.* **245**, 1180 [1970].