

## Over-Production of Porphyrins and Heme in Heterotrophic Bacteria

W. K. Philipp-Dormston and M. Doss

Abteilung für Klinische Biochemie, Arbeitsgruppe Biochemische Mikrobiologie, am Hygiene-Institut des Fachbereichs Humanmedizin der Philipps-Universität, Marburg a. d. Lahn

(Z. Naturforsch. 30 c, 425–426 [1975]; received September 18, 1974/January 10, 1975)

Porphyrins, Heme,  $\delta$ -Aminolevulinic Acid, Porphobilinogen, Microorganisms

In *Escherichia coli*, *Pseudomonas aeruginosa*, and *Achromobacter metalcaligenes*  $\delta$ -aminolevulinic acid synthase and  $\delta$ -aminolevulinic acid dehydratase appear to be the rate limiting enzymes of porphyrin and heme biosynthesis. Bypassing of these enzymes by addition of  $\delta$ -aminolevulinic acid or porphobilinogen results in overproduction of tetrapyrroles.

Comparative studies of porphyrin and heme biosynthesis indicate that  $\delta$ -aminolevulinic acid synthase (EC 2.3.1.37) is the rate limiting enzyme in this biosynthetic pathway in all microorganisms tested so far<sup>1–3</sup>. There is little information about the regulatory function of  $\delta$ -aminolevulinic acid dehydratase (EC 4.2.1.24), which has been isolated from *Saccharomyces cerevisiae*<sup>4</sup>, *Propionibacterium shermanii*<sup>5</sup>, and *Rhodospseudomonas spheroides*<sup>6</sup>. In the present study concentrations of porphyrins and heme were determined in *Escherichia coli*, *Pseudomonas aeruginosa*, and *Achromobacter metalcaligenes* following biosynthesis from the direct precursors of uroporphyrinogen, either  $\delta$ -aminolevulinic acid (ALA) or porphobilinogen (PBG). Thus the corresponding enzymes, which might have a regulatory function in the tetrapyrrole synthesis of heterotrophic bacteria, were bypassed.

Sources of the organisms *E. coli*, *Ps. aeruginosa*, and *A. metalcaligenes* as well as the composition of a defined medium have been reported previously<sup>7</sup>.

Supplements added to the medium in various experiments are given in the legends to the tables. ALA was obtained from Serva, Heidelberg, and PBG produced by *Propionibacterium shermanii*<sup>9</sup> was the kind gift of Dr. G. Müller, University of Stuttgart. All cultures were grown in 500 ml Erlenmeyer flasks at 37 °C with moderate aeration for up to 42 hours<sup>1</sup>. Cell density (dry weight/ml) was determined by turbidity measurement<sup>9</sup>.

Porphyrins and heme were measured spectrophotometrically as methyl esters after separation by silica gel thin-layer chromatography<sup>10</sup>. Protoheme was converted into its pyridine hemochrome and assayed by difference spectrophotometry<sup>11</sup>. Small amounts of porphyrin esters were analysed as copper chelates<sup>12</sup> or by fluorescence measurement<sup>13</sup>.

The three species synthesised heme and all porphyrins in the biosynthetic chain in varying amounts, as summarized in the Tables. The supplements had no significant influence upon growth of the cells.

The results reported in Table I indicated, that during biosynthesis from DL-lactate the porphyrin patterns of *E. coli* and *Ps. aeruginosa* are similar in that coproporphyrin is the main component formed in each, whereas *A. metalcaligenes* synthesised predominantly protoporphyrin. Addition of ALA or PBG elevated porphyrin and heme biosynthesis considerably, but in all cultures overproduction from ALA appeared to be greater than from PBG (Table II, III).

Iron(III) citrate (10  $\mu$ M) enhanced heme formation in all culture systems about 2-fold, whereas total porphyrin synthesis both from DL-lactate and ALA was reduced by exogenous iron by about 40% (Table I, II). Porphyrin synthesis from PBG was not reduced by iron (Table III). In addition iron also altered the ratio of copro-/protoporphyrin as described previously<sup>7</sup>.

Table I. Porphyrin and heme biosynthesis from DL-lactate with and without addition of iron(III) citrate (10  $\mu$ M) in *Escherichia coli* (1), *Pseudomonas aeruginosa* (2), and *Achromobacter metalcaligenes* (3). T = traces.

Organism	Addition of iron	Heme [pmol/mg]	Total porphyrins [pmol/mg]	Distribution of porphyrins [%]						
				Uro-	7 CO <sub>2</sub> H-	6 CO <sub>2</sub> H-	5 CO <sub>2</sub> H-	Copro-	3 CO <sub>2</sub> H-	Proto-
1	—	21	60	15	11	6	9	41	8	10
	+	38	28	12	4	T	2	33	10	39
2	—	14	157	17	10	6	8	49	5	5
	+	32	63	14	4	2	6	47	8	19
3	—	28	38	10	7	2	8	19	16	38
	+	66	18	7	1	T	T	49	17	26

Requests for reprints should be sent to Prof. Dr. M. Doss, Division of Clinical Biochemistry, Hygiene Institute of the University, Pilgrimstein 2, D-3550 Marburg a. d. Lahn, Federal Republic of Germany (FRG).



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

Table II. Porphyrin and heme biosynthesis from  $\delta$ -aminolevulinic acid (0.2 mM) with and without addition of iron(III)-citrate (10  $\mu$ M) in *Escherichia coli* (1), *Pseudomonas aeruginosa* (2), and *Achromobacter metalcaligenes* (3).

Organism	Addition of iron	Heme [pmol/mg]	Total porphyrins [pmol/mg]	Distribution of porphyrins [%]						
				Uro-	7 CO <sub>2</sub> H-	6 CO <sub>2</sub> H-	5 CO <sub>2</sub> H-	Copro-	3 CO <sub>2</sub> H-	Proto-
1	—	69	607	36	13	3	7	33	3	5
	+	116	446	41	10	5	8	27	3	6
2	—	61	937	42	12	3	5	35	1	2
	+	109	686	49	9	2	3	31	2	4
3	—	82	450	38	7	6	8	17	7	17
	+	149	283	31	5	2	4	41	8	9

Table III. Porphyrin and heme biosynthesis from porphobilinogen (0.1 mM) with and without addition of iron(III)-citrate (10  $\mu$ M) in *Escherichia coli* (1), *Pseudomonas aeruginosa* (2), and *Achromobacter metalcaligenes* (3).

Organism	Addition of iron	Heme [pmol/mg]	Total porphyrins [pmol/mg]	Distribution of porphyrins [%]						
				Uro-	7 CO <sub>2</sub> H-	6 CO <sub>2</sub> H-	5 CO <sub>2</sub> H-	Copro-	3 CO <sub>2</sub> H-	Proto-
1	—	43	370	33	7	3	8	42	3	4
	+	80	356	36	3	2	6	36	4	13
2	—	39	419	34	8	4	7	38	4	5
	+	68	397	40	7	2	5	29	4	13
3	—	55	287	36	9	3	5	14	10	23
	+	99	297	31	9	2	8	24	10	16

The formation of smaller amounts of porphyrin and heme from PBG than from ALA might be due to the rate of passage of the different molecules through the membranes. The amount of endogenous heme appears to be closely related to the iron concentration in the culture fluid<sup>7</sup>. The observation that porphyrin formation from DL-lactate drops when endogenous heme concentrations is increased and that tetrapyrrole synthesis is enhanced by addition of ALA favours ALA-synthase as the rate limiting enzyme in this biosynthetic chain<sup>3,14</sup>. During incubation with exogenous ALA porphyrin formation is also depressed by higher heme levels, whereas synthesis from PBG appears to be unaffected by heme. This suggests repression of ALA-dehydratase by heme, which has also been observed in photosynthetic bacteria<sup>6</sup>. It is conceivable too that

ALA-dehydratase is repressed by high concentrations of ALA, as has been shown in enzyme preparation from *Rhodospseudomonas spheroides*<sup>15</sup>, although the physiological significance of this observation is difficult to assess.

The present results suggest that, in analogy to tetrapyrrole biosynthesis in phototrophic bacteria<sup>3,6</sup>, regulation in the heterotrophic bacteria of the overall synthesis seems to be associated with ALA-synthase and ALA-dehydratase. The effect of iron, which alters the copro-/protoporphyrin ratio, might be due to species-specific differences and appears to be of secondary nature.

This study was supported by the Deutsche Forschungsgemeinschaft (Do 134/5).

<sup>1</sup> M. Doss and W. K. Philipp-Dormston, Hoppe-Seyler's Z. Physiol. Chem. **352**, 725–733 [1971].

<sup>2</sup> W. K. Philipp-Dormston and M. Doss, Enzyme (Basel) **16**, 57–64 [1973].

<sup>3</sup> J. Lascelles, Tetrapyrrole biosynthesis and its regulation. W. A. Benjamin, New York-Amsterdam 1964.

<sup>4</sup> O. I. Clara De Barreiro, Biochim. Biophys. Acta **139**, 479–486 [1967].

<sup>5</sup> W. Walerych, Acta Biochim. Pol. **10**, 243–252 [1963].

<sup>6</sup> B. F. Burnham and J. Lascelles, Biochem. J. **87**, 462–472 [1963].

<sup>7</sup> M. Doss and W. K. Philipp-Dormston, Hoppe-Seyler's Z. Physiol. Chem. **352**, 43–51 [1971].

<sup>8</sup> G. Müller and G. Bezold, Z. Naturforsch. **24b**, 47–53 [1969].

<sup>9</sup> F. Kavanach, Analytical Microbiology, Vol. II, pp. 43–121, Academic Press, New York-London 1972.

<sup>10</sup> M. Doss, Hoppe-Seyler's Z. Physiol. Chem. **350**, 499–502 [1969].

<sup>11</sup> M. Doss, Klin. Wschr. **46**, 731–732 [1968].

<sup>12</sup> M. Doss, Anal. Biochem. **39**, 7–14 [1971].

<sup>13</sup> M. Doss, B. Ulshöfer, and W. K. Philipp-Dormston, J. Chromatogr. **63**, 113–120 [1971].

<sup>14</sup> A. Neuberger, Regulation of Porphyrin and Heme Biosynthesis (ed. M. Doss), pp. 74–78, S. Karger, Basel 1974.

<sup>15</sup> J. Lascelles, J. Gen. Microbiol. **23**, 487–498 [1960].