

Phase Shifting Effect of Caffeine in the Circadian Rhythm of *Phaseolus coccineus* L.

W. Mayer and I. Scherer

Institut für Biologie I, Tübingen

(Z. Naturforsch. 30 c, 855–856 [1975]; received August 27, 1975)

Caffeine, Circadian Rhythm, Sleep and Wakefulness, *Phaseolus coccineus* L.

4-hour caffeine pulses (10 mM) offered *via* the transpiration stream advances or delays the phase of the circadian leaf movement rhythm of *Phaseolus coccineus* as a function of the phase of application. It is hypothesized that the caffeine effect upon sleep and wakefulness in man is partly due to this phase-shifting effect.

Light and temperature are the best known Zeitgeber for circadian rhythms in plants and animals^{1,2}. Administered as pulses they shift the phase of the circadian oscillation as a function of the phase of application.

Many chemicals were tested, but only a few act on the circadian clock. A clear phase-shifting effect has been shown only for D₂O^{2,3}, ethanol^{2,4,5}, Valinomycin^{2,5}, K⁺^{2,4} and CN⁻⁶.

Recently, a phase-shifting effect of theophylline and pentobarbital has been described for the deep body temperature rhythm in the rat⁷. Furthermore, a phase-shifting and period-lengthening effect of

theophylline could be shown in the case of the circadian leaf movement rhythm in *Phaseolus coccineus*⁸. In addition to theophylline (1,3-dimethylxanthine) we tested the phase-shifting effect of caffeine (1,3,7-trimethylxanthine) on the leaf movement rhythm of *Phaseolus coccineus*.

Fig. 1 shows typical examples of movements of two isolated primary leaves of the same plant under continuous light (LL) of 5600 lx and 23 °C. One leaf served as a control. To the other leaf 10 mM caffeine was offered *via* the transpiration stream for 4 h. A pulse starting 4.4 h after "0" time (corresponding to the second maximum of the night position in LL) delayed the rhythm by 2 h (Fig. 1, above), a pulse starting 21.2 h after "0" delayed by about 4.5 h (Fig. 1, center) and a pulse starting 24.2 h after "0" advanced the rhythm by 2.5 h (Fig. 1, below).

These results show that not only theophylline, but also caffeine has in addition to its well known pharmacological effects a further one: it rephases a circadian rhythm.

Since circadian rhythms in unicellular organisms, plants, animals, and man have similar properties², we hypothesize, that not only the theophylline effect⁷, but also the caffeine effect upon sleep and wakefulness in man is partly due to the phase-shifting effect of these substances. This could be of




Fig. 1. Typical examples for circadian leaf movements of two isolated primary leaves of one individual plant of *Phaseolus coccineus* under LL 5600 lx and 23 ± 1 °C showing the phase-shifting effect of 4 h pulses of caffeine (10 mM) (····) control leaves; (—) leaves are treated in the noted phase (above 4.4 h, center 21.2 h, below 24.2 h after "o" corresponding to the second maximum of the night position in LL).

Requests for reprints should be sent to Dr. W. Mayer, Institut für Biologie I, D-7400 Tübingen, Auf der Morgenstelle 1.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

pharmacological interest, since it opens the possibility to use caffeine and theophylline as drugs for the phase control of circadian rhythms *e.g.* of transmeridional travellers and shift-workers.

This work has been supported by the Schwerpunktprogramm "Biologie der Zeitmessung" Deutsche Forschungsgemeinschaft (Ma 386/4).

¹ J. Aschoff, Circadian Clocks, North-Holland Publishing Co., Amsterdam 1965.

² E. Bünning, The Physiological Clock, revised third edit., Springer, Berlin-Heidelberg-New York 1973.

³ A. Maurer and W. Engelmann, Z. Naturforsch. **29 c**, 36–38 [1974].

⁴ E. Bünning and I. Moser, Proc. Nat. Acad. Sci. U.S. **70**, 3387–3389 [1973].

⁵ B. M. Sweeney, Plant Physiol. **53**, 337–342 [1974].

⁶ W. Steinheil, Z. Pflanzenphysiol. **62**, 204–215 [1970].

⁷ C. F. Ehret, V. R. Potter, and K. W. Dobra, Science **188**, 1212–1215 [1975].

⁸ W. Mayer, R. Gruner, and H. Strubel, Planta, in press.