

Chemical Constituents of the Essential Oils of Aroliaceae and Lauraceae Families

Nanao Hayashi, Kôki Yokochyô and Hisashi Komae

Faculty of Integrated Arts and Sciences, Hiroshima University, Hiroshima

(Z. Naturforsch. 30 c, 421 [1975]; received February 24, 1975)

Aroliaceae, Leaf Oil, Mono- and Sesquiterpenes, Aromatic Compounds, Lauraceae

In the course of the chemosystematic study of Aroliaceae and Lauraceae families, the leaf oils of some plants were investigated. The leaf oils were obtained by steam distillation of the fresh leaves. The individual constituents were isolated by column chromatography followed by preparative gas chromatography, and identified by gas chromatography and by infrared spectra.

Acanthopanax innovans Franch. et Sav. (Aroliaceae). Source. Hiroshima Prefecture, Japan. Leaf Oil. Steam distillation of the fresh leaves gave a yield of 0.03% oil which possess the following physical properties: n_D^{25} 1.4895, α_D^{25} -25.15° . Constituents. Individual constituents were isolated by the preparative gas chromatography (GLC) and identified by infrared spectra: α -pinene (0.6%), camphene (0.3%), *cis*-ocimene (0.4%), *trans*-ocimene (5.2%), myrcene (3.9%), caryophyllene (14.5%), β -farnesene (50.5%).

Acanthopanax sciadophylloides Franch. et Sav. (Aroliaceae). Source. Hiroshima Prefecture. Leaf Oil. The leaf oil of this plant was obtained in 0.025% yield. The physical properties determined

as follows: n_D^{25} 1.4872, α_D^{25} $+7.55^\circ$. Constituents. α -pinene (2.0%), β -pinene (3.0%), myrcene (21.4%), caryophyllene (31.6%), β -farnesene (24.0%), α -humulene (5.0%), ethylbenzene (trace), *m*-xylene (trace), *o*-xylene (trace) were isolated by preparative GLC and identified by IR spectra.

Lindera communis var. *Okinaensis* (Kamik) Hatusima (Lauraceae). Source. Hyakuna in Okinawa Prefecture. Leaf Oil. Steam distillation of the fresh leaves gave a yield 0.19% leaf oil which possess the following properties: d_4^{25} 0.8913, n_D^{25} 1.4951, α_D^{25} $+12.3^\circ$. Constituents: α -pinene (9.8%), camphene (4.9%), β -pinene (12.5%), *cis*-ocimene (17.5%), *trans*-ocimene (5.5%), caryophyllene (17.2%).

Cassytha filiformis Linn. (Lauraceae). Source. Hyakuna in Okinawa Prefecture. Stem Oil. The essential oil of the stems of this plant was obtained by the steam distillation in 0.075% yield. The physical properties determined as follows: d_4^{25} 0.9080, n_D^{25} 1.4860, α_D^{25} $+2.7^\circ$. Constituents. A sesquiterpene hydrocarbon, caryophyllene, was isolated by preparative GLC and identified by IR spectrum.

Lindera sericea var. *glabrata* Blume. (Lauraceae). Source. Yuki in Hiroshima Prefecture. Leaf Oil. The essential oil from the leaves of the plant by steam distillation has been obtained in 0.32% yield and has the following physical properties: d_4^{25} 0.9736, n_D^{25} 1.4955, α_D^{25} $+34.1^\circ$. Constituents: α -corocalene, β -eudesmol, cadalene, curcumenone and shyobunone were isolated by preparative GLC and identified by IR spectra in addition to the previous paper¹.

Requests for reprints should be sent to Dr. Nanao Hayashi, Faculty of Integrated Arts and Sciences, Hiroshima University, Hiroshima 730, Japan.

¹ Nanao Hayashi and Hisashi Komae, Z. Naturforsch. 28 c, 227 [1973].



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.