

## Conductance Characteristics of Interactive Subunit Membranes

A. H. Bretag and D. I. B. Kerr

School of Pharmacy, South Australian Institute of Technology and Department of Human Physiology and Pharmacology, University of Adelaide, Adelaide

(Z. Naturforsch. 30 c, 535–537 [1975]; received April 1, 1975)

### Subunit Interactions, Membrane Conductance Models

Membrane channels formed by groups of subunits with a special form of interaction between the subunits provide a simple basis for models of membrane conductance. It is shown how the subunit interaction can explain conductance kinetics; particular attention being paid to characteristics of the potassium conductance in the squid giant axon.

We have arrived at an explanation for some of the unique features of ion conductance through cell membranes. Here we show how "induction" and "superposition" of potassium conductance in the squid axon<sup>1,2</sup> can result from a special form of interaction between subunits of a planar array membrane. Various forms of interaction and cooperativity have been considered previously in attempts to describe potassium conductance<sup>2–11</sup>, but few of these have proved any more successful than the original non-interactive Hodgkin-Huxley<sup>12</sup> formulation.

In several cases where claims have been made for potassium conductance models involving membrane subunit or pore interactions<sup>4–9</sup>, it has been shown that the mathematical descriptions of the models are approximations only<sup>2–5</sup>. In particular, accurate computations of Adam's model<sup>6</sup> by Hill and Chen<sup>2,3</sup> have shown his approximation to be invalid and his model to be inadequate.

By contrast, the approximations shared by the models of Bretag, Davis and Kerr<sup>4,5</sup>, Tille<sup>7</sup> and Starzak<sup>8,9</sup> remain, so far undisputed. The validity of these approximations and their relation to the exact mathematical description of the interaction in these models will be discussed in a forthcoming publication. Meanwhile, further consideration of the form of the interaction has provided a simple mechanistic explanation for the conductance characteristics of the models.

The interaction in these models occurs between nearest neighbour sites in a two dimensional lattice representing the membrane. Sites may exist in two states, active and inactive. The special constraint is that a site cannot undergo a change of state unless it is situated adjacent to a site in the active state.

Requests for reprints should be sent to Dr. A. H. Bretag, School of Pharmacy, South Australian Institute of Technology, North Terrace, Adelaide, South Australia 5000.

"Seeding" or "nucleation" is a natural consequence of this form of interaction with the rate of increase in the number of active sites being low while the number of such sites is low and increasing as the number of active sites increases. In other words, if the external determinant of site activation (e.g. membrane potential) is increased, then the initial rate of increase in the number of active sites will depend upon the number of such sites existing at that time. Consequently, the models demonstrate "induction" and the duration of the induction period depends on the initial number of activated sites (Fig. 1). Inflexion occurs in the rate of increase when further increase in the number of active sites becomes limited by either the level of the external activation determinant or, eventually, by the shortage of activable sites. This, of course, always assumes that the proportion of active sites is being increased from a low level where not all inactive sites are nearest neighbours of active sites; which leads us to the second important property of these models.




Fig. 1. Induction behaviour. a. Induction illustrated by an arbitrary function

$$x = f(t),$$

where  $x$  has been plotted against

$$t' = t - t_0,$$

$t_0$  being the point chosen for the origin of the abscissa. The smaller the value of  $x$  at  $t = t_0$ , the longer the induction period. b. Induction illustrated by activation in a square lattice where only nearest neighbours may be activated. Beginning with one active site, four further sites may be activated, while beginning with two, three or four; six, at least seven, or at least eight respectively, may be activated. Active sites are indicated by "8", inactive nearest neighbours by "4" and other inactive sites by "1".

Once the proportion of active sites is great enough so that effectively all inactive sites are nearest neighbours of active sites, then further activation is free of our constraint regarding interaction: further activation then being dependent



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

solely on the level of the external determinant and the availability of activable sites. The rate of activation is thereafter constant at any specified proportion of active sites, for a fixed external determinant.

a 41111484448484888844114888484148414484884488448441  
8411148444848488884114844441484111148444144884484  
41141488484484414441114888841484414114848414848441  
11484844484144848444144848414484841484411414411  
14844884148484448848488884148444844448411144841  
484148414484484111444448484148488448848411484414  
48841444848441111484441414848484144441114841448  
144148844484841114884884144844141114884148844848  
1111484844884114144844844884111148444148448414  
4114844488444414848411414846414111484884448484441  
844441488444484844484111414848411441488484444884  
4488441484844444114411411148414841444444111441  
1488484844844484114844484114411488484448844111  
48484884144844141144844484148411448448844484141  
14844888448841484411448444114111444148841141484  
114484484144411448411444844841488448841414848  
14888414841484148441114844844141444484414848844  
14884848484141484114448841488448411141114844841  
41444844141484114114884484488414114114141448411  
841488841148414114848884414848441484148448484411  
4111488484144148411484484148414484884148414441411  
414414414148414414148884114414884441488488448414  
84844841144448448448884114884484488414884488448  
848488844484484484444441448484444144484484488  
4844444484844884414848484448844144484488441488844  
1484414414144484444848844844484448448448444448  
148488488444888448488484114114888414484144848448  
4144841444844484844484441141444484484441484414  
848484484148414441484144844484884141444884141148  
41414148411414841148411448841484111484484841484

b 4444884888444488888484484448484488488884488888888  
4888841484844484844414888888848888484484848888  
84888414848848848448844888484414844484888848888  
41488888884884888844888488484848444488884888  
844888488488488848448488448844888841488848844884  
8884484848884488888888848848448448888484888848  
8484888488841448448888848848884144488441444488  
44848888848488888444444148888888448488884888444  
11488848448448448488484144848884844148448884488  
11448888448848844848888484884484848484884884484  
14848848444848848448488888448888888488488884884  
48848888848448488888484884488488848488488484844  
88488848488488444484848444888884888848488844888  
8444441444484888884844884484444884888448884484  
88888444888848884448441448441484884488848448448  
848488888484884444441488848884848488488488484  
8888414844888488848888484488448414884888488448  
4844888488448488888448884848484848448488414448488  
844848884484148841488448448888844848484844448484  
44844144884888848848884844448484484848888441  
888484888448448488488844848844844848884848844884  
44884488844884888444488884488488448884488888441  
484844488484888884448888448888444484484888884484  
48484888884848884484484488848844488844888844884  
4888444848848844884448888844848888484888848884484  
484448448448844484484484884884484884488884888488  
8888484841484848848888844884144888484848884484  
888414884844888488488844884844888444884484884888




Fig. 2. Square array structure for a potassium-conducting subunit membrane. The rounded form of a subunit indicates the activated state, "8", which, when present in all subunits of a minimum functional group (all subunits adjacent to A), allows the formation of a conducting channel or pore at A. Openings at B and C are considered inadequate for the passage of ions.

Fig. 3. Parts of two  $50 \times 50$  square arrays representing potassium-conducting patches of membrane. Subunits are labelled as in Fig. 1. In the first array (a) 26.2% of subunits were randomly set as active. Of those remaining 67.8% were then nearest neighbours of the active subunits. In the second array (b) 55.9% of subunits were randomly set as active. This time 98.8% of the remainder were nearest neighbours. These two cases represent equilibrium potassium conductances, 0.4% and 9.8% of maximum, respectively, in the BDK square array model.

Thus rates of further activation to a fixed higher level show "superposition". That this is a legitimate property of the Bretag, Davis and Kerr<sup>4</sup> (BDK) potassium conductance model can be illustrated in the following way:

Let us consider a square array for the lattice where four neighbouring sites are required to be simultaneously activated to produce a potassium conductance channel (Fig. 2). Potassium conductance thus depends (if the site distribution is random) on the fourth power of the relative number of active sites. Then, for the resting potassium conductance to be close to that observed experimentally at the normal resting potential<sup>13</sup>, it is necessary for the proportion of active sites to be about 25% (c.f.  $n_0 = 0.24$  for the Hodgkin-Huxley<sup>12</sup> activating particle model). This proportion is already close to that where effectively all inactive sites are nearest neighbours of active sites (Fig. 3). Hence, kinetics for potassium currents, even only slightly above the resting level, will be approximately superimposable for a potential step to a fixed higher potential level. Any initial deviation would rapidly become insignificant as the potassium current increased.

The BDK model is therefore able to fit the experimental data of Cole and Moore<sup>1</sup> with respect to both induction and superposition, even though it does not strictly obey the conditions of stochastic independence of sites imposed by both Hill and Chen<sup>2</sup> and Albano<sup>14</sup>. This is because the working range of the model includes two distinctly different regions: one where there is significant interaction which specifies induction and another where there is no significant interaction which allows superposition.

Although the principles involved in the BDK model are therefore sound, it should be noted that

the parameters used in the model were derived by fitting curves to the experimental potassium conductance data of Hodgkin and Huxley<sup>12</sup>.

Recent work<sup>15,16</sup> indicates that such modelling is in error because the conductance curves of Hodgkin and Huxley<sup>12</sup> are not true axolemmal conductance curves; these latter authors having failed to consider potassium ion accumulation in the Schwann Cell Layer (S.C.L.). However, conflicting views of the seriousness of the error are presented<sup>15,16</sup>. Adam<sup>15</sup> even implies that axolemmal conductance kinetics may indeed not be superimposable and that the superposition of current curves may result from modifications imposed by the S.C.L. We feel that his extreme view is argued against by the experimental results of Cole and Moore<sup>1</sup>. Potassium currents following conditioning hyperpolarisation to various levels were just as superimposable as those following conditioning depolarisation. In the prior case, S.C.L. effects would be identical and independent of the level of conditioning hyperpolarisation, which strongly suggests that axolemmal conductance is superimposable. In fact, Adelman, Palti and Senft<sup>16</sup> take the view that axolemmal conductance is satisfactorily modelled merely by adjusting the maximum conductance  $\bar{g}_K$  and the rate constants  $\alpha_n$  and  $\beta_n$  in the Hodgkin-Huxley formulation. We are therefore confident that similar minor alterations to our model would enable it also to describe these recent findings as well as maintaining its ability to describe Cole-Moore induction and superposition.

Finally, we feel that the form of interaction specified in the BDK model may also prove more widely applicable: especially if the constraint is relaxed to allow some probability of change of state in those sites not situated adjacent to an active site.

<sup>1</sup> K. S. Cole and J. W. Moore, *Biophys. J.* **1**, 1 [1960].

<sup>2</sup> T. L. Hill and Y. Chen, *Proc. Nat. Acad. Sci. U.S.* **68**, 1711 [1971].

<sup>3</sup> Y. Chen and T. L. Hill, *Proc. Nat. Acad. Sci. U.S.* **70**, 62 [1973].

<sup>4</sup> A. H. Bretag, B. R. Davis, and D. I. B. Kerr, *J. Membr. Biol.* **16**, 363 [1974].

<sup>5</sup> A. H. Bretag, C. A. Hurst, and D. I. B. Kerr, *J. Theor. Biol.* **48**, 477 [1974].

<sup>6</sup> G. Adam, *Physical Principles of Biological Membranes* (F. Snell, J. Wolken, G. Iversen, and J. Lam, eds.), p. 35, Gordon and Breach, New York 1970.

<sup>7</sup> J. Tille, *Biophys. J.* **5**, 163 [1965].

<sup>8</sup> M. E. Starzak, *J. Theor. Biol.* **39**, 487 [1973].

<sup>9</sup> M. E. Starzak, *J. Theor. Biol.* **39**, 505 [1973].

<sup>10</sup> D. M. J. Quastel, *Physiology Canada* **4**, 203 [1973].

<sup>11</sup> T. L. Hill and Y. Chen, *Proc. Nat. Acad. Sci. U.S.* **68**, 2488 [1971].

<sup>12</sup> A. L. Hodgkin and A. F. Huxley, *J. Physiol. (London)* **117**, 500 [1952].

<sup>13</sup> A. L. Hodgkin and A. F. Huxley, *J. Physiol. (London)* **116**, 473 [1952].

<sup>14</sup> A. M. Albano, *J. Theor. Biol.* **45**, 275 [1974].

<sup>15</sup> G. Adam, *J. Membr. Biol.* **13**, 353 [1973].

<sup>16</sup> W. J. Adelman, Y. Palti, and J. P. Senft, *J. Membr. Biol.* **13**, 387 [1973].