

**Some Aliphatic Methyl Esters
from *Aralia elata* Seemann (Araliaceae)**

S. Watanabe, Y. Hayashi, and Y. Murayama

Department of Pharmacology, Kawasaki University School
of Medicine, Kurashiki

(Z. Naturforsch. **30 c**, 825 [1975]; received
June 6, 1975)

Aralia elata Seemann, Aliphatic Methyl Ester, 1-Hexacosene

The dried bark of *Aralia elata* Seemann was extracted with *n*-hexane. Methyl pentadecanoate, methyl hexadecanoate, methyl octadecanoate, methyl eicosanoate, methyl docosanoate, methyl tetracosanoate, methyl hexacosanoate, and 1-hexacosene were identified from the extract by means of GC-Mass spectrometer.

In the course of the investigation of the bark components in *Aralia elata* Seemann which has been used as a folk medicine¹⁻³ for diabetes mellitus, we identified seven aliphatic methyl esters and 1-hexacosene from *n*-hexane extract by GC-Mass spectrometry.

Plant. The dried bark (1 kg) of *Aralia elata* Seemann, collected on May 1st at Okayama Prefecture in Japan, was extracted with methanol (9 l) for four weeks. The residue obtained after removal of the solvent was extracted with *n*-hexane. The hexane fraction was evaporated and chromato-

graphed on silica gel column (Merck, Kiesel gel 60). First elution with *n*-hexane – Et₂O (10:1) gave oily matter (2.23 g). The constituents of the oil were detected by means of GC-Mass spectrometer (GC: SE-52 0.75% on Chromosorb W, temperature programmed 50 to 250 °C, 5 °C/min; Mass: ionizing voltage 20 eV, ionaccelerating voltage 1000 V, ionization chamber temperature 220 °C, sample injection block temperature 160 °C) on the basis of the characteristic fragmentation patterns and the retention time as follows: methyl pentadecanoate 1.0%, methyl hexadecanoate 43.2%, methyl octadecanoate 27.5%, methyl eicosanoate 5.6%, methyl docosanoate 9.4%, methyl tetracosanoate 6.5%, and methyl hexacosanoate 6.8%. The common fragments of the above methyl esters are *m/e* 74 (base peak), 87, 143, 199 and M⁺-43. (The percentage of the constituents of the oil was calculated from the areas of the peaks of the gas chromatogram.) Among these constituents, 1-hexacosene (2.6 g) was isolated by column chromatography at second eluate and identified using mass spectrometry: *m/e* (relative abundance); 69 (49), 73 (100), 97 (89), 111 (52), 125 (28), 139 (14), 153 (8), 167 (6), 181 (4), 195 (3), 207 (2), 335 (5), 364 (5). The main components in *n*-hexane extract were seven methyl esters and 1-hexacosene.

Requests for reprints should be sent to Dr. S. Watanabe, Department of Pharmacology, Kawasaki University School of Medicine, 577 Matsushima, Kurashiki, 701-01, Japan.

¹ S. Kuwata, J. Pharm. Soc. Japan **49**, 660 [1929].

² S. Kuwata and M. Harima, *ibid.* **49**, 668 [1929].

³ S. Kuwata, *ibid.* **51**, 17 [1931].



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.