

Chirality of the Hydrogen Transfer to NAD Catalyzed by *myo*-Inositol Dehydrogenase from *Klebsiella pneumoniae*

Miguel A. Alizade, Karl Gaede

Centro de Biofísica y Bioquímica, Instituto Venezolano de Investigaciones Científicas (IVIC), Caracas

and

Klaus Brendel

Department of Pharmacology, College of Medicine, The University of Arizona, Tucson, Arizona

(Z. Naturforsch. 31c, 624–625 [1976]; received May 19/June 21, 1976)

Chirality, Hydrogen Transfer, *myo*-Inositol Dehydrogenase

The chirality of the hydrogen transfer to NAD catalyzed by *myo*-inositol dehydrogenase (*myo*-inositol: NAD 2-oxido-reductase, EC 1.1.1.18) from *Klebsiella pneumoniae* (formerly classified taxonomically as *Aerobacter aerogenes* or *Klebsiella aerogenes*) was investigated. $|4\text{-}^3\text{H}|$ NAD was enzymatically reduced to $|4\text{-}^3\text{H}|$ NADH with non-labeled *myo*-inositol and *myo*-inositol dehydrogenase. The stereochemistry of the prochiral center at C4 of the NADH produced was determined. It was found that the label was exclusively located at the (4S) position of the produced NADH. Since the hydrogen transferred from non-labeled *myo*-inositol to $|4\text{-}^3\text{H}|$ NAD must have entered the opposite or (R) position, *myo*-inositol dehydrogenase from *K. pneumoniae* should be classified as an (R) or A-type enzyme with respect to the stereochemistry of the hydrogen transfer to NAD.

The absolute stereochemistry of direct hydrogen transfer from substrates to the prochiral C₄ position of the nicotinamide ring of the coenzyme, and vice versa, catalyzed by pyridine nucleotide-linked dehydrogenases has been conclusively demonstrated^{2–5}. Dehydrogenases able to catalyze the hydrogen transfer to the pro(R) or pro(S) position of the C₄ prochiral center of the nicotinamide ring of the coenzyme⁴ were classified as A or B-type enzymes, respectively^{2–5}.

A large number of NAD and NADP-linked dehydrogenases have been investigated with regard to their stereochemistry of hydrogen transfer to the coenzyme. Results are summarized in references 6 to 8.

In this communication, the chirality of hydrogen transfer catalyzed by the inducible NAD-linked dehydrogenase *myo*-inositol dehydrogenase (EC 1.1.1.18) from *K. pneumoniae*¹ was investigated. $|4\text{-}^3\text{H}|$ NAD was enzymatically reduced to $|4\text{-}^3\text{H}|$ NADH with *myo*-inositol and *myo*-inositol dehydro-

genase from *K. pneumoniae*. The chirality at the C₄ position of the produced $|4\text{-}^3\text{H}|$ NADH was analyzed by transfer of the hydrogen of the (4S) position to α -ketoglutarate with the B-type (S) glutamate dehydrogenase. From Table I one can ascertain that more

Table I. Stereochemistry of the enzymic hydrogen transfer from *myo*-inositol to $|4\text{-}^3\text{H}|$ NAD catalyzed by *myo*-inositol dehydrogenase from *Klebsiella pneumoniae*.

Specific radioactivities * [dpm/ μ mol]			
$ 4\text{-}^3\text{H} $ NAD ^a	$ 4\text{-}^3\text{H} $ NADH ^a	(S) glutamate ^b	$ 4\text{-}^3\text{H} $ NADH ^c
9.2×10^6	9.0×10^6	8.7×10^6	2.8×10^5

* The specific radioactivities of Table I refer to the following steps in the following reaction scheme:

$$|4\text{-}^3\text{H}| \text{ NAD}^a + \text{myo-inositol} \xrightarrow{\text{IDH}} 2\text{-keto-myoinositol} + |4\text{-}^3\text{H}| \text{ NADH}^a$$

$$|4\text{-}^3\text{H}| \text{ NADH}^a + \alpha\text{-ketoglutarate} + \text{NH}_3 \xrightarrow{\text{GDH}} (\text{S}) \text{ glutamate}^b + |4\text{-}^3\text{H}| \text{ NAD}^c$$

$$|4\text{-}^3\text{H}| \text{ NAD}^c + (\text{S}) \text{ lactate} \xrightarrow{\text{LDH}} \text{pyruvate} + |4\text{-}^3\text{H}| \text{ NADH}^c$$

than 95% of the label originally present in the (4S)-position of the generated $|4\text{-}^3\text{H}|$ NADH is transferable to α -ketoglutarate by the reaction catalyzed by (S) glutamate dehydrogenase. This outcome was confirmed in another experiment, in which the specific radioactivity of the concomitantly produced NAD was determined. For that purpose the NAD content of another aliquot from the incubation mixture was enzymatically reduced to NADH with nonlabelled (S)lactate and (S)lactate dehydrogenase. As expected only a small fraction (3%) of the original label remains attached to the newly formed NADH (NADH^c from Table I). These results prove that the label of the originally produced $|4\text{-}^3\text{H}|$ NADH is located at the (4S) position of the nicotinamide ring. Hence the hydride transferred from nonlabelled *myo*-inositol to $|4\text{-}^3\text{H}|$ NAD catalyzed by *myo*-inositol dehydrogenase must have entered the (4R)-position of the produced (4S) $|4\text{-}^3\text{H}|$ NADH. This result allows the classification of the investigated *myo*-inositol dehydrogenase from *K. aerogenes* as an (R) or A-type enzyme.

Experimental Section

Myo-Inositol, pyruvic acid, (S)lactic acid, (S)-glutamic acid, (S)lactate dehydrogenase (EC 1.1.1.27) from rabbit muscle, (S)glutamate dehydrogenase (EC 1.4.1.3) from beef heart, and *myo*-inositol dehydrogenase (EC 1.1.1.18) from *Klebsiella pneumoniae*^{9,10} were purchased from Sigma Chemical Co. $|4\text{-}^3\text{H}|$ NAD with a specific radio-

Requests for reprints should be sent to Dr. M. A. Alizade, Instituto Venezolano de Investigaciones Científicas (IVIC-SAS), Apartado 1827, Caracas 101, Venezuela.

Abbreviations used: IDH, *myo*-inositol dehydrogenase (EC 1.1.1.18); GDH, (S)glutamate dehydrogenase (EC 1.4.1.3); LDH, (S)lactate dehydrogenase (EC 1.1.1.27).



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

activity of 50 Ci/mol was obtained from The Radiochemical Center, Amersham. *Myo*-inositol was recrystallized three times from water by addition of ethanol. *Myo*-Inositol dehydrogenase was further purified by chromatography at 5 °C on a 1 × 30 cm P2-column (Bio-Rad Lab.) previously equilibrated with 0.05 M Tris HCl buffer of pH 8.2 (measured at 25 °C) and obtained by elution with the same buffer.

The enzymatic hydrogen transfer from *myo*-inositol to $[4\text{-}^3\text{H}]\text{NAD}$ was performed incubating 9.0 μmol of $[4\text{-}^3\text{H}]\text{NAD}$ with a specific radioactivity

of 9.2×10^6 dpm/ μmol with 48 μmol of non-labelled *myo*-inositol and 0.4 units of *myo*-inositol dehydrogenase (19 U/mg) in a total volume of 3.0 ml of reaction medium containing in addition 600 μmol of Tri-HCl buffer previously adjusted to pH 8.4 at 25 °C. After a 10 minute incubation, 0.6 μmol $[4\text{-}^3\text{H}]\text{NADH}$ was isolated^{11,12} and its chirality at C₄ determined as already described¹².

Work supported in part by the Regional Program of Scientific and Technological Development of the Organization of the American States (OAS), the Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICIT) and USPHS Grant NIH AM 14977.

¹ Bergey's Manual of Determinant Bacteriology (1974) (R. E. Buchanan and N. E. Gibson, eds.), pp. 322–324, The Williams and Wilkins Company, Baltimore.

² F. H. Westheimer, F. H. Fischer, E. E. Conn, and B. Vennesland, J. Amer. Chem. Soc. **73**, 2403 [1951].

³ H. R. Levy, F. A. Loewus, and B. Vennesland, Progress in Stereochemistry (P. B. D. de la Mare and W. Klyne, eds.), Vol. 3, pp. 329, Butterworths, London 1962.

⁴ J. W. Cornforth, G. Ryback, G. Popjack, C. Donninger, and G. Schroeffer, Biochem. Biophys. Res. Commun. **9**, 371–375 [1962].

⁵ G. Krakow, J. Ludowieg, J. H. Mether, W. M. Narmore, L. Tosi, S. Udaka, and B. Vennesland, Biochemistry **2**, 1009–1014 [1963].

⁶ R. Bentley, Molecular Asymmetry in Biology, Vol. 2, pp. 1–89, Academic Press, New York 1970.

⁷ G. Popjack, Stereochemistry of Enzymic Reactions, in: The Enzymes (P. D. Boyer, ed.), Vol. 2, pp. 115–215, Academic Press, New York 1970.

⁸ B. Vennesland, Stereospecificity in Biology, In: Topics in Current Chemistry, Vol. 48, pp. 39–65, Springer Verlag, Berlin, New York 1974.

⁹ W. A. Anderson and B. Magasanik, J. Biol. Chem. **246**, 5653–5661 [1971].

¹⁰ T. K. Sundarama, J. Bacteriol. **111**, 284–286 [1972].

¹¹ E. Silverstein, Anal. Biochem. **12**, 199–212 [1965].

¹² M. A. Alizade, R. Bressler, and K. Brendel, Biochim. Biophys. Acta **364**, 204–208 [1974].