

Studies on the Retention of CF₁ with or without Induced ATPase Activity by Pyrophosphate Treated Thylakoids and Its Relation to the Regeneration of Photophosphorylation

R. Schopf and G. Harnischfeger

Lehrstuhl für Biochemie der Pflanze der Universität Göttingen

(Z. Naturforsch. **32 c**, 398–404 [1977]; received March 22, 1977)

CF₁/ATPase-Reconstitution, Photophosphorylation, Fluorescence Labeling

The reconstitution of chloroplast coupling factor 1 (CF₁) into thylakoid membranes was investigated by the fluorescence of the covalently attached label fluorescamine. In contrast to a functional regeneration of ATP synthesis, a rebinding of CF₁ was observed regardless if the protein was in its native, purified state or had been activated for ATPase activity by heat, dithiothreitol (DTT) or trypsin treatment. The reintegration of CF₁ into the thylakoid membrane was estimated to be almost quantitative.

Since the label binds covalently and irreversibly to free NH₂-groups, its effects on the various functional parameters were investigated. Blocking of NH₂ groups leads to inhibition of ATPase activity in isolated CF₁. Labeled factor used in a reconstitution experiment on the other hand, does still result in considerable regeneration of ATP formation.

Attaching the label to whole thylakoids leads to a differential effect on phosphorylation, which is inhibited, and the H⁺-uptake which seems largely unaffected. A differential effect is also observed on coupled and uncoupled electron transport.

The results are interpreted as evidence for an involvement of free NH₂-groups in the mechanism of photophosphorylation.

Introduction

The success of reconstituting phosphorylation activity by recombination of CF₁ and CF₁-depleted thylakoid membranes depends largely on the amount of proteins not extracted from the plastids during EDTA or pyrophosphate treatment (review by Nelson¹). The critical lower limit for sufficient regeneration is a loss of more than 50% of native CF₁ during the preparation of CF₁ depleted membrane particles². This observation has been attributed to an insufficient retention of the purified CF₁ due to changes in surface properties leading *e.g.* to aggregation. Alternative though related is the notion that its close to complete extraction through either EDTA or pyrophosphate treatment leads to irreparable destruction of membrane structure². A third possibility derives from the argument that the protein retained during the reconstitution experiment serves only in a structural capacity (re-

sealing of "holes") while the catalysis of ATP formation is restricted to the residual, native CF₁³. In this case the less residual CF₁, the less the maximum theoretical activity (on a chlorophyll basis) obtainable in a reconstitution experiment. It is possible that all three factors contribute to the aforementioned observation of a critical CF₁ concentration retained in the "CF₁-depleted" membranes.

One approach to investigate if retention of protein constitutes the critical factor is the use of fluorescent covalent labeling techniques which was employed by Kraayenhof and Slater⁴ to study conformational phenomena of CF₁ activity. Similarly, a double fluorescent label was suggested by Harnischfeger and Schopf⁵ to investigate the physical parameters of membrane-CF₁ interaction. CF₁ was labeled in both cases with fluorescamine which reacts specifically with exposed amino groups. Only the covalently bound dye possesses a strong fluorescence whereas the molecule itself and its hydrolyzation products are non-fluorescent.

This paper describes the influence of the fluorescamine label on the activity of isolated CF₁, the reconstitution of activity with labeled CF₁ and also the effects of ATPase-activation on labeled and unlabeled protein with regard to reconstitution. The experiments suggest that retention of protein during reconstitution is not the critical factor limiting ac-

This paper is dedicated to the memory of the late Professor Dr. G. Jacobi who initiated and encouraged many aspects of this work.

Requests for reprints should be sent to Dr. Götz Harnischfeger, Lehrstuhl für Biochemie der Pflanze, Untere Karspüle 2, D-3400 Göttingen.

Abbreviations: 9-AAC, 9-aminoacridine; CF₁, chloroplast coupling factor 1; DTT, dithiothreitol; PMS, phenazine-methosulfate; P.P., pyrophosphate.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

tivity for a regeneration of photophosphorylation. In addition applying the label to the thylakoids rather than to CF₁ suggests, that in the mechanism of phosphorylation and electron transport free amino groups play an important role.

Material and Methods

Coupling factor CF₁ was prepared according to Lien and Racker⁶ with the modification that the chloroplasts were washed prior to EDTA treatment with 5 mM pyrophosphate at pH 7.4 in order to remove stroma proteins and carboxydismutase⁷. The protein, pure according to electrophoretic observation, was stored in 20 mM Tris – SO₄ – 2 mM ATP – 2 mM EDTA at pH 7.1.

For the use in the experiments the CF₁ preparation was centrifuged, the precipitate resolved in 10 mM Tricine pH 8 and passed through a Sephadex G-25 column (20 × 1.5 cm). After elution with 20 mM Tricine pH 8.0 the CF₁ was effectively separated from (NH₄)₂SO₄ as determined by the Neßler test.

ATPase activity was determined through released inorganic phosphate using the method of Taussky and Shorr⁸. Activity of ATPase was induced through either trypsin incubation⁹, DTT treatment¹⁰ or heat exposure¹¹.

CF₁ was labeled with fluorescamine in the concentrations given in the legends. Since the dye is insoluble in water, it was added in acetone solution. The amount of acetone in the labeling assay did not exceed 5%. The labeling assay contained bicarbonate buffer (10 mM, pH 8.3) in order to prevent any possible influence through the amino groups of Tricine. Fluorescence was measured in the fluorimeter described by Harnischfeger¹², excitation was at 365 nm, the emission spectrum was recorded between 400 and 500 nm.

Uncoupled thylakoids which have lost between 25–30% of their native CF₁ were prepared according to Schopf *et al.*¹³. They were partially depleted of CF₁ by hypotonic treatment with 0.5 mM pyrophosphate of pH 7.4. Chlorophyll concentration at this step was 0.1 mg/ml.

Cyclic photophosphorylation was assessed in the PMS system. The assay contained in 2.2 ml: 100 μ mol Tricine pH 8, 10 μ mol MgCl₂, 5 μ mol ATP, 10 μ mol Na₂HPO₄ including ³²P, 100 μ mol PMS and 40 μ g chlorophyll eq. chloroplasts. Illumination was with 9.5×10^5 erg/cm² × sec of light with a wavelength of >610 nm. The determination of ATP³²P formed was according to Lindberg and

Ernster¹⁴ using Cerenkov emission in the liq. scintillation assay¹⁵.

The fluorescence of 9-aminoacridine, used as an indication of membrane energetization, was measured as described before¹². The determination of the size of the proton uptake was accomplished with a sensitive glass electrode (Ingold LOT 405-M5) attached to a Metrohm pH meter. Starting pH of the assay was 6.2, illumination by red light ($\lambda > 610$ nm) of an intensity of 9.5×10^5 erg × cm⁻¹ × sec⁻¹. The assay contents were similar to those described by Strotmann¹⁶.

In the reconstitution assay the uncoupled thylakoids and purified CF₁ were recombined. The medium contained in addition 0.1 M sucrose, 25 mM Tricine pH 8, and 10 mM MgCl₂. The reconstituted system was assayed for photochemical activity after an incubation time of 20 min at 0 °C.

Results

1. The retention of CF₁ activated for ATPase activity

Our experiments confirm the initial observation by Jagendorf¹⁷ that a regeneration of photophosphorylation is impossible with CF₁, in which ATPase activity was induced by treatment with trypsin or heat. It can only be achieved, if the activation was obtained by DTT exposure (Table I).

Table I. Regeneration of cyclic photophosphorylation using CF₁ in which ATPase activity was induced prior to the reconstitution experiment.

Thylakoids	CF ₁	Phosphorylation activity	ATPase activity of CF ₁
Control	—	419	—
P.P. washed	—	4	—
P.P. washed	+CF ₁	123	—
P.P. washed	+DTT-CF ₁	129	5.9
P.P. washed	+trypsin-CF ₁	3	6.4
P.P. washed	+heat-CF ₁	10	6.2

Activity of phosphorylation in the PMS system given in μ mol ATP formed × mg Chl⁻¹ h⁻¹; rate of ATP hydrolysis by activated CF₁ (type of activation indicated by prescript) given in μ mol ATP × mg protein⁻¹ min⁻¹. DTT-CF₁ was purified before reconstitution by Sephadex G-25, the removal of DTT did not influence ATPase activity.

Further investigation showed, that the reconstituting capacity is already lost long before the trypsin treatment yields the maximum rate of ATP hydrolysis (Fig. 1). In the same manner, the 9-AAC




Fig. 1. Influence of trypsin treatment of CF₁ on its ability to regenerate phosphorylation in a reconstitution experiment. The rate of ATP formation in the untreated control was $290 \mu\text{mol} \times \text{mg Chl}^{-1} \text{h}^{-1}$.




Fig. 2. Influence of trypsin treatment of CF₁ on its ability to regenerate the 9-AAC quench after reconstitution.

quench is diminished after reconstitution with the briefly activated coupling factor (Fig. 2).

These observations lead to the question whether the ATPase-activated CF₁ does not bind to the thylakoid or whether it is only inactive but fully integrated into the membrane. One possible solution is offered by reconstitution experiments with CF₁ to which a fluorescent label was attached through exposure to fluorescamine. After a sufficient incuba-




Fig. 3. Concentration dependence of CF₁ retention of thylakoids. CF₁ (0.9 mg/ml) was labeled with $250 \mu\text{M}$ fluorescamine and used in increasing amounts for this series of reconstitution experiments. Fluorescence was measured after two washes of the reconstitution assay to remove excess CF₁.

tion of the reconstitution mixture unreacted CF₁ can be removed by centrifugation and washing, and the amount of CF₁ retention can be estimated by the remaining fluorescence signal. (Two washes of the reconstituted particles in the incubation medium sufficed in our experiments to remove all fluorescence emission due to the fluorescamine-protein adduct in the supernatant.) Fig. 3 shows the result of such an experiment with purified, labeled, coupling factor. Approximately 3–4 mg CF₁/mg Chl sufficed to saturate retention. The actual amount of coupling factor incorporated into the membrane is quite small, only 3% of the added protein under our conditions. This value was calculated in Table II from the relative size of the fluorescence signal and estimated to be around 0.1 mg protein/mg chlorophyll. This corresponds to the amount of CF₁ released during pyrophosphate washing¹³.

Table II. Determination of the amount of CF₁ rebound to thylakoid membranes during reconstitution.

	Relative fluorescence signal eq. $\mu\text{g CF}_1$ ($F_{365} \rightarrow 470$)	
Entire reconstitution assay	820	655
Reconstituted thylakoids after removal of unreacted CF ₁	17	13.6

CF₁ (0.9 mg/ml) was marked with $250 \mu\text{M}$ fluorescamine prior to use in the reconstitution assay which contained besides the above amount of CF₁, $136 \mu\text{g}$ chlorophyll eq. uncoupled thylakoids. The fluorescence signal was determined before and after removal of non reacted CF₁ using aliquot samples.

The coupling of fluorescamine with purified CF₁ is influenced by a number of factors which can distort the information obtainable from reconstitution experiments. One poignant example is given in the model reaction of Fig. 4 where lysine labeled with fluorescamine was used. The presence of DTT or ATP during labeling reduces the emission of the adduct. ATP is itself susceptible to the label and can, therefore, influence the result. The DTT dependent decrease was, however, not observed when the compound was added after the labeling reaction. DTT must, therefore, interfere with the chemical mechanism forming the covalent bond.




Fig. 4. The influence of DTT and ATP on the fluorescence signal of the fluorescamine adduct. The label (250 μ M) was attached to lysine (10 mM). The two components were present in the conc. given during the labeling procedure.

This also can explain the observation of Kraayenhof and Slater⁴, that fluorescamine labeling of CF₁ is much diminished after DTT treatment. Their interpretation, that amino groups exposed in the native enzyme are buried after ATP-ase activation with DTT is, therefore, open to question. Consequently, in order to measure the retention of the ATPases, it was necessary to separate prior to reconstitution those substances which were added for the activation of CF₁.

In the case of DTT, the activated protein was separated by column chromatography with Sephadex G-25 and subsequently labeled with fluorescamine.

Taking this precaution the retention of activated CF₁ in reconstitution experiments is almost identical regardless of the procedure used to induce ATPase activity. Moreover, as summarized in Table III, the fluorescence signal of the rebound protein was comparable to that of the uninduced coupling factor. The retention of the trypsin treated protein did not change even after 5 min of trypsin exposure. This finding does not agree with experiments by Racker *et al.*¹⁸ who observed a diminished degree of retention of trypsin activated CF₁-ATPase, measured by ATP hydrolysis of the supernatant after reconstitution.

Table III. Influence of ATPase induction in CF₁ on its re-binding capacity in reconstitution experiments.

	Rel. fluorescence signal ($F_{365} \rightarrow 470$)
CF ₁ , purified	10.4
DTT-CF ₁	10.6
Heat-CF ₁	9.8
Trypsin-CF ₁	10.8

Incubation of CF₁ for full activation of ATPase activity: DTT — 3 h, trypsin — 1 min, heat (64°C) — 4 min. Labeling with 250 μ M fluorescamine after removal of by-products and activation reagents. Reconstitution and other procedures as described in methods and in the legend to Table II.

2. The reconstitution of the proton uptake and photophosphorylation with labeled CF₁

Although the retention of CF₁ in the reconstitution experiment is complete as shown in Table II, the capacity for energy conservation of the reconstituted system should nevertheless be influenced by the label of the coupling factor.

We observed, that the fluorescamine label affects markedly the ATPase activity of isolated CF₁. Fig. 5 shows that the hydrolytic capacity is inhibited around 90% at the concentration of 100 μ M fluorescamine in the labeling assay. If the fluorescence emission of the marked CF₁ serves as indicator, this concentration corresponds to a saturation of the exposed amino-groups (Fig. 6). Since trypsin by its proteolytic action generates new amino terminal groups, more fluorescamine can react with them and the respective curve in Fig. 6 is shifted towards higher concentrations. These results indicate, that the fluorescamine labeled CF₁ used in the reconstitution experiments is enzymatically inactive, pro-




Fig. 5. Influence of the fluorescamine label on the ATPase activity of isolated CF₁. The protein (0.2 mg/ml) was labeled after activation by the methods indicated. Activity of 100% is equal to 13.4 $\mu\text{mol ATP hydrolysed} \times \text{mg prot.}^{-1} \text{min}^{-1}$ for trypsin activated CF₁, 5.6 for heat activated and 6.1 for DTT activated protein.




Fig. 6. Fluorescence saturation curve for fluorescamine labeling for isolated CF₁. The protein (0.2 mg/ml) was labeled after activation as indicated.

vided, phosphorylation and hydrolysis of ATP proceed by a similar mechanism involving the same catalytic sites.

However, as demonstrated in Fig. 7, the proton uptake and the phosphorylation is similarly reconstituted if unlabeled CF₁ or a coupling factor which was marked with fluorescamine was used. Only higher concentrations of label inhibit slightly. At 250 μM fluorescamine, which was generally used in the experiments described before, the reconstitution is almost the same as with the unlabeled coupling factor. The absolute values for the controls were 138 $\mu\text{mol of ATP} \times \text{mg Chl}^{-1} \text{h}^{-1}$ and 168 nmol $\text{H}^+ \times \text{mg}^{-1}$ chlorophyll for the proton uptake corresponding to a 30% regeneration.




Fig. 7. Influence of fluorescamine labeling of CF₁ on its ability to regenerate the proton uptake and phosphorylation in reconstitution experiments. CF₁ (0.9 mg/ml) was labeled with the amount of fluorescamine indicated prior to the reconstitution assay. 100% activity were 138 $\mu\text{mol ATP} \times \text{mg Chl}^{-1} \text{h}^{-1}$ (7 μmol before reconstitution) and 168 nmol $\text{H}^+/\text{mg Chl}$ (16 nmol before reconstitution).

3. Reconstitution with labeled thylakoids

The free amino groups seem to play an important role in the coupling mechanism of phosphorylation. This is evident from the experiment shown in Fig. 8, where control type thylakoids (shocked but still possessing their full complement of CF₁) were exposed to various concentrations of fluorescamine. A differential effect on photophosphorylation and proton uptake is observed. At 250 μM the former was inhibited, the latter still unaffected. Basis electron transport (FeCy acceptor) is not influenced by the label, while the uncoupled activity is pronouncedly decreased (Fig. 9).




Fig. 8. Influence of fluorescamine labeling on phosphorylation and proton uptake of intact thylakoids. The thylakoids (0.2 mg Chl/ml) were labeled with the fluorescamine concentration indicated. The 100% values are 147 nmol $\text{H}^+ \cdot \text{mg Chl}^{-1}$ and 121 $\mu\text{mol ATP} \cdot \text{mg Chl}^{-1} \cdot \text{h}^{-1}$ for photophosphorylation (FeCy-acceptor).




Fig. 9. Influence of fluorescamine labeling on electron transport of intact thylakoids (FeCy-acceptor).




Fig. 10. Influence of thylakoid-membrane labeling by fluorescamine on the ability to regenerate cyclic photophosphorylation in reconstitution experiments. Chlorophyll concentration in the labeling assay was 0.2 mg/ml.

Fig. 10 describes the effect of labeling thylakoids with fluorescamine on the regeneration of activity in a reconstitution experiment. In analogy to the above results an inhibition of cyclic phosphorylation is observed in both controls and reconstituted thylakoids leading to the same final value, though on a relative basis the influence on the reconstituted system is much less. This finding argues, that in these experiments the active (coupling?) site of the residual CF₁, which is not removed by washing from the uncoupled thylakoids, is affected, since labeled CF₁, on the other hand, does not greatly impair the reconstituted activity (compare Fig. 6).

Discussion

The experiments give information about both the structural and the functional aspect of reconstitution. From a structural point of view the finding is important, that all types of CF₁, normal or activated by the various means available, bind to the thylakoid membrane. A rough quantitative estimate from the fluorescence data gives a retention of 0.1 mg protein/mg Chl. The total amount of CF₁ in intact thylakoids has been given as 0.42–0.45 mg CF₁/mg Chl^{7,19}. Taking this value into con-

sideration the amount rebound would correspond to about 30% of the total, *i.e.* exactly the protein removed through EDTA or 0.5 mM pyrophosphate treatment. The structural reconstitution thus seems quantitative.

Of special interest is the observation that CF₁ with trypsin induced ATPase activity is reintegrated into the membrane during reconstitution. Deters *et al.*²⁰ reported that under these conditions CF₁ loses its δ subunit and concluded from the absence of function that the protein did not rebound. Similarly Nelson and Karny²¹ implied a strong correlation between binding of CF₁ and the δ subunit. If we imply, — though we have not shown it —, that in analogy to Deters *et al.*²⁰ trypsin treatment in our case also removes the δ subunit from CF₁, we have to conclude that the δ subunit cannot be the only structural link between CF₁ and thylakoid membrane.

The functional aspect, *i.e.* the ability to regenerate photophosphorylation, is more complex. Hesse *et al.*² imply 3 successive degrees of membrane breakdown during CF₁ extraction, namely, disorganization of grana stacking — removal of CF₁ — membrane perforation. They report membrane perforation as starting when about 50% of native CF₁ is ex-

tracted. Since the uncoupled thylakoids used in our experiments contain still about 70% of their native CF₁, holes in the membrane (not to be confused with the normal ion transmitting pores) and thus a resealing through the offered protein during reconstitution is not the decisive factor for regeneration of phosphorylation activity. As discussed by Jagendorf¹⁷ the functional aspect of those CF₁ molecules that rebind is not clear. The regeneration of phosphorylation with inactive, fluorescamine labeled CF₁ implies, that the renewed activity rests mostly, but not necessarily exclusively, in the residual membrane factor (cp. also Berzborn and Schröder²²). The result suggests that in this case the rebound CF₁ restores only membrane integrity which is required for energy conservation.

The labeling experiments seem to indicate also that free amino-groups play a role in the coupling between phosphorylation and electron transport. These groups are virtually abolished through the covalent attachment of fluorescamine²³ in saturating amounts. Although the proton uptake is

unaffected at 100 μ M fluorescamine, the phosphorylation is only 10% its original size. Thus, the energy reservoir, if proton uptake serves as a rough indicator, is still present but cannot be diverted into ATP formation. Labeling the thylakoids before reconstitution is much more sensitive than labeling CF₁, which however can be easily explained by a strictly structural organizational role of the rebound CF₁. NH₂-groups must be involved at the coupling site proper, since fluorescamine labeling decreases uncoupled electron transport in class I chloroplasts but leaves basal rates practically unaffected (Fig. 9).

The results so far indicate that although in reconstitution experiments CF₁ is retained at all the binding sites, the crucial feature is the repair of energy transfer between membrane and coupling factor. This problem warrants further investigation.

The technical assistance of Ms. M. Koch is gratefully acknowledged. The study was supported by grants from the Deutsche Forschungsgemeinschaft (DFG).

- ¹ N. Nelson, *Biochim. Biophys. Acta* **456**, 314–338 [1976].
- ² H. Hesse, E. Jank-Ladwig, and H. Strotmann, *Z. Naturforsch.* **31 c**, 445–451 [1976].
- ³ R. Schmid and W. Junge, *Proc. IIIrd. Int. Congr. Photosynthesis*, Rehovoth (M. Avron, ed.), p. 821–830, Elsevier, Amsterdam 1975.
- ⁴ R. Kraayenhof and R. C. Slater, *Proc. Ird Int. Congr. Photosynthesis*, Rehovoth (M. Avron, ed.), p. 985–996, Elsevier, Amsterdam 1975.
- ⁵ G. Harnischfeger and R. Schopf, *Z. Naturforsch.* **32 c**, 398 [1977].
- ⁶ S. Lien and E. Racker, *Methods in Enzymology XXIII* (A. San Pietro, ed.), p. 547–551, Academic Press, New York 1971.
- ⁷ H. Strotmann, H. Hesse, and K. Edelmann, *Biochim. Biophys. Acta* **314**, 202–210 [1973].
- ⁸ H. H. Taussky and E. Shorr, *J. Biol. Chem.* **202**, 675–685 [1953].
- ⁹ V. K. Vambutas and E. Racker, *J. Biol. Chem.* **240**, 2660–2667 [1965].
- ¹⁰ R. E. McCarty and E. Racker, *J. Biol. Chem.* **243**, 129–137 [1968].
- ¹¹ F. Farron, *Biochemistry* **9**, 3823–3828 [1970].
- ¹² G. Harnischfeger, *Habilitationsschrift*, Göttingen 1976.
- ¹³ R. Schopf, K.-P. Heise, B. Schmidt, and G. Jacobi, *Proc. IIIrd Int. Congr. Photosynthesis* (M. Avron, ed.), p. 911–919, Elsevier, Amsterdam 1975.
- ¹⁴ O. Lindberg and L. Ernster, *Methods Biochem. Anal.* **3**, 1–22 [1956].
- ¹⁵ W. Hoffmann, *Radiochim. Acta* **4**, 117–119 [1965].
- ¹⁶ H. Strotmann, *Habilitationsschrift*, Hannover 1971.
- ¹⁷ A. T. Jagendorf, *Bioenergetics of Photosynthesis* (Govindjee, ed.), p. 414–419, Academic Press, New York 1975.
- ¹⁸ E. Racker, G. A. Hauska, S. Lien, R. J. Berzborn, and N. Nelson, *Proc. IInd Int. Congr. Photosynth.* (G. Forti, ed.), p. 1097–1113, The Gague 1972.
- ¹⁹ G. Hauska and P. V. Sane, *Z. Naturforsch.* **27 b**, 938–942 [1972].
- ²⁰ D. Deters, E. Racker, N. Nelson, and H. Nelson, *J. Biol. Chem.* **250**, 1041–1047 [1975].
- ²¹ N. Nelson and O. Karny, *FEBS-Letters* **70**, 249–253 [1976].
- ²² R. J. Berzborn and P. Schröder, *FEBS-Letters* **70**, 271–275 [1976].
- ²³ S. Udenfriend, S. Stein, P. Böhlen, W. Dairman, W. Leimgruber, and M. Weigele, *Science* **178**, 871–872 [1972].