

Minor Phenolic Constituents of *Grevillea robusta* and *Hakea saligna*

Manju Varma, R. S. Varma, and
M. R. Parthasarathy [1]

Department of Chemistry, University of Delhi, Delhi-110007 (India)

Z. Naturforsch. **35 c**, 344–345 (1980);
received November 15, 1979

Proteaceae, *Grevillea robusta*, *Hakea saligna*, Lignans

Three hitherto unknown phenolic constituents epilyoniresinol, its glucoside and rhamnocitrin-3-O-rutinoside have been isolated from *G. robusta* wood and leaves besides kaempferol-3-O-rutinoside whereas 2,6-dimethoxybenzoquinone, lyoniresinol and its rhamnoside are obtained from *H. saligna* wood. Their structures were assigned on the basis of chemical and spectral evidence.

Previous work has shown the presence of several 5-*n*-alkyl resorcinol and macrocyclic phenols in the wood [2] and leaves [3] of *Grevillea robusta*. Parallel studies on Proteaceae members naturalised in India revealed the presence of other new phenolic components from wood [4, 5] and leaves [6] of *G. robusta* and *Hakea saligna*. We wish to report now the occurrence of few minor constituents from wood and leaves of these species, extraction procedures being described earlier [4–6].

G. robusta, wood: The combined acetone and methanol extract after fractionation on silica gel column followed by purification on preparative SiO₂-TLC in solvent EtOAc-H₂O-MeOH (40:5.4:6.6) afforded compounds (A) and (B).

Compound A, pinkish prisms from EtOAc/C₆H₆ (150 mg from 2 kg of wood) m. p. 198–200 °C, $[\alpha]_D^{MeOH} = -28.9^\circ$ (Found: C, 63.0; H, 7.0; calculated for C₂₂H₂₈O₈ C, 62.8; H, 6.6). It gave colour reaction with FeCl₃ (changing green to dark blue), positive Mäule test and blue colour with *p*-NO₂C₆H₄N₂⁺BF₄⁻ and 20% Na₂CO₃ overspray, a test for syringyl lignans. The IR spectrum displayed the aromatic and phenolic hydroxyl absorptions (ν_{max}^{KBr} 3472, 1600, 1493 and 1449 cm⁻¹) confirmed by the bathochromic shift in UV (from λ_{max}^{MeOH} 278 nm to 290 nm) on addition of alkali. It formed a dimethyl ether (**1 a**), m. p. 148–149 °C (Me₂SO₄-K₂CO₃-Me₂CO) and tetraacetate (**1 b**), m. p. 145 °C with

Py/Ac₂O. PMR spectrum of acetate (60 MHz, CDCl₃, TMS, δ): 6.54 (s, 1H, Ar-H), 6.34 (s, 2H, Ar-H), 4.28 (m, 1H), 4.08 (m, 4H, -CH₂O-), 3.83 (s, 6H, ArO-CH₃), 3.76 (s, 3H ArO-CH₃), 3.23 (s, 3H, ArO-CH₃), 2.90–2.58 (m, 4H), 2.32 (s, 6H, ArO-COCH₃), 2.08 (s, 3H, -CH₂O-COCH₃) and 2.04 (s, 3H, -CH₂O-COCH₃). These data suggested it to be a disyringyl lignan, the physical properties being closely resembling to lyoniresinol [7]. However, its co-IR and co-TLC with authentic sample showed some minor differences. In view of these and its different $[\alpha]_D$, it was considered as an epimer of lyoniresinol with structure **1** or its enantiomer.

Compound B, brownish gum (90 mg) gave all tests for compound (A) and a positive Molisch's test. IR spectrum (ν_{max}^{KBr} 3448, 1608, 1493 and 1449 cm⁻¹) and UV (λ_{max}^{MeOH} 275 and $\lambda_{max}^{MeOH-NaOH}$ 280 nm) were almost identical with compound (A). The acid hydrolysis afforded glucose as sugar and compound (A) as aglycone m. p. 197–198 °C. The PMR spectrum of heptaacetate (Found: C, 57.2; H, 6.1; C₄₂H₅₂O₂₀ requires C, 57.7; H, 5.9) showed in addition to the signals for compound (A), a complex multiplet at δ 5.16–4.19 ascribable to sugar protons. The major difference, however, was in the methoxyl region where the highest field methoxyl of compound (A) (δ 3.23) was now appearing as a pair of peaks (δ 3.32 and 3.22). This behaviour, ascribed to the presence of rotamers, has been reported in the case of lyoniresinol-2 α -O-rhamnoside acetate [7]. For similar reasons, it was identified as epi-lyoniresinol-2-O-glucoside (**1 c**).

G. robusta leaves: The column chromatography of acetone extract afforded a glycosidic fraction from which two closely moving compounds (C) and (D) were obtained by preparative SiO₂-TLC in solvent CHCl₃-MeOH-H₂O (33:13.5:1.8). Both of them gave greenish brown colour with FeCl₃, colour tests for flavonoids and were found to be glycosidic in nature.

Compound C, yellow plates from MeOH, m. p. 167–168 °C (200 mg from 500 g of leaves). UV λ_{max}^{MeOH} 270, 343 nm with no shift on addition of NaOAc but a bathochromic shift of long wavelength maximum to 405 nm on addition of AlCl₃. On total acid hydrolysis with H₂SO₄ it gave glucose and rhamnose as sugars and rhamnocitrin as aglycone identified by comparison with authentic sample (co-TLC, co-IR and m. m. p.). Permethylolation and hydrolysis by Hakomori's method [8] yielded two meth-

Reprint requests to Dr. M. Varma, School of Physical and Mol. Sciences, University College of North Wales, Bangor, Gwynedd LL57 2DG, U. K.

0341-0382/80/0300-0344 \$ 01.00/0





Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.



ylated sugars identified as 2,3,4-trimethyl glucopyranose (R_{TMS} 0.84) and 2,3,4-trimethyl rhamnopyranose (R_{TMS} 0.98) confirmed by direct comparison with the samples obtained under similar conditions from rutin. That the compound (C) is rhamnocitrin-3-O-rutinoside (2) was inferred from PMR of its octaacetate m. p. 124–125 °C (Found: C, 55.7; H, 5.6; $C_{44}H_{48}O_{23}$ requires C, 55.9; H, 5.1): δ 8.13 (d, 2H, $J = 9$ Hz, 2',6'-H), 7.02 (d, 2H, $J = 9$ Hz, 3',5'-H), 6.82 (d, 1H, $J = 2$ Hz, 8-H), 6.61 (d, 1H, $J = 2$ Hz, 6-H), 5.16–4.89 (m, sugar protons), 3.85 (s, 3H, ArO-CH₃), 2.43 (s, 3H, ArO-COCH₃), 2.38 (s, 3H, ArO-COCH₃), 2.09, 2.03, 1.95 (m, 18H, -OCOCH₃) and 1.07 (br, 3H, C-CH₃).

Compound D, yellow needles from MeOH, m. p. 180–182 °C. It was identified as kaempferol-3-O-rutinoside (2a) by UV spectral analysis with usual shift reagents [9] and other experiments as described for compound (C). A direct comparison with authentic sample (co-TLC, co-IR and m. m. p.) confirmed the identity.

Hakea saligna wood: The combined acetone and methanol extract on SiO₂ column chromatography gave compounds (E), (F) and (G).

Compound E, from C₆H₆-EtOAc (1:4) eluates as pinkish prisms (125 mg from 2.8 kg of wood), m. p. 192–193 °C, $[\alpha]_D^{MeOH} = 0^\circ$. It answered tests for syringyl lignans and showed resemblance to compound (A) (TLC, IR). It was identified as (±) lyoni-resinol [7] by comparison with an authentic sample (co-TLC, co-IR and m. m. p.).

Compound F, from EtOAc eluates, yellow needles (140 mg), m. p. 258–260 °C. It gave typical tests for quinones [10] IR (ν_{max}^{KBr} 1698 and 1650 cm⁻¹). It was inferred as 2,6-dimethoxy benzoquinone (UV and PMR) and confirmed by comparison with authentic sample (co-TLC, co-IR and m. m. p.).

Compound G, from EtOAc-MeOH (19:1) eluates, purified by preparative SiO₂-TLC in solvent EtOAc-H₂O-MeOH (40:5.4:6.6), amorphous solid m. p. 108–109 °C. It answered tests for syringyl lignans and also the Molisch's test. On hydrolysis it gave rhamnose as sugar and lyoni-resinol as aglycone. Its identity as lyoni-resinol-2 α -O-rhamnoside [7] was confirmed by direct comparison with an authentic specimen (co-TLC, co-IR).

It is the first report of the natural occurrence of the epimer of lyoni-resinol and its glucoside as well as rhamnocitrin-3-O-rutinoside. The other compounds has been reported from different sources; this, however, is the first report of their occurrence in Proteaceae. The minor stereochemical differences in the lyoni-resinol based lignan constituents of both the woods and the occurrence of glucoside in *G. robusta* and rhamnoside in *H. saligna* are the main distinctive features. However, a definite conclusion on the role of any of these compounds as chemotaxonomic markers requires investigation of more representative samples from different regions.

- [1] We are grateful to Dr. John W. Rowe, U. S. D. A., Forest Products Laboratory; Madison, Wis., USA, for the Samples of Lyoni-resinol and Its Rhamnoside.
- [2] E. Ritchie, W. C. Taylor, and T. K. Vautin, Australian J. Chem. **18**, 2015 (1965); M. Rasmussen, D. D. Ridley, E. Ritchie, and W. C. Taylor, Australian J. Chem. **21**, 2989 (1968).
- [3] J. R. Cannon, P. W. Chow, M. W. Fuller, B. H. Hamilton, B. W. Metcalf, and A. J. Power, Australian J. Chem. **28**, 2257 (1973).
- [4] R. S. Varma, Manju, and M. R. Parthasarathy, Phytochemistry **15**, 1418 (1976).

- [5] Manju and M. R. Parthasarathy, Ind. J. Chem. Sec. B, **15**, 1090 (1977).
- [6] Manju, R. S. Varma, and M. R. Parthasarathy, Phytochemistry **16**, 793 (1977).
- [7] F. D. Hostettler and M. K. Seikel, Tetrahedron **25**, 2325 (1969).
- [8] S. Hakomori, J. Biochem. **55**, 205 (1964).
- [9] J. B. Harborne, T. J. Mabry, and H. Mabry, The Flavonoids, Chapman and Hall, London 1975.
- [10] R. H. Thomson, Naturally Occurring Quinones, Academic Press, London 1971.