

## On the Reactivity of Pyridoxal-5'-phosphate with Yeast tRNA<sup>Phe</sup> and tRNA<sup>Tyr</sup>

Nobuo Okabe\* and Friedrich Cramer

Max-Planck-Institut für experimentelle Medizin, Abteilung Chemie, Hermann-Rein-Straße 3, D-3400 Göttingen

Z. Naturforsch. **35 c**, 522–525 (1980);  
received February 11/March 17, 1980

Yeast tRNA<sup>Phe</sup>, Yeast tRNA<sup>Tyr</sup>, Pyridoxal-5'-phosphate

Yeast tRNA<sup>Phe</sup> and tRNA<sup>Tyr</sup> were reacted with the fluorescent reagent pyridoxal-5'-phosphate and the modified tRNAs were analysed with respect to the number and position of modified nucleosides and with respect to aminoacylation.

a) Following the intrinsic fluorescence of pyridoxal-5'-phosphate, the treatment of tRNA<sup>Tyr</sup> with increasing amounts of pyridoxal-5'-phosphate revealed about 50 mol of reagent or a even higher number bound per one mol of tRNA<sup>Tyr</sup>. After borohydride reduction (in order to stabilize the linkage) of this modified tRNA<sup>Tyr</sup> and purification with reverse phase chromatography a modified tRNA<sup>Tyr</sup> was obtained carrying about 2 mol of the reagent.

b) Both tRNA<sup>Tyr</sup> and tRNA<sup>Phe</sup> treated with pyridoxal-5'-phosphate and reduced exhibited almost unchanged aminoacylation as compared to the unmodified tRNAs.

c) Pyridoxal-5'-phosphate treated and reduced tRNA<sup>Phe</sup> and tRNA<sup>Tyr</sup> were digested with ribonuclease T<sub>1</sub> and the resulting oligonucleotides were separated. However, no fluorescent oligonucleotide and no difference to an oligonucleotide pattern obtained from unmodified tRNA were observed.

Thus, pyridoxal-5'-phosphate might have been bound to the highly purified yeast tRNA<sup>Phe</sup> and tRNA<sup>Tyr</sup> samples either via an unstable linkage or not covalently. This result is controversial with respect to the specific reaction of pyridoxal-5'-phosphate with unfractionated tRNAs from colon carcinoma and tRNAs from *E. coli* as reported in the literature.

## Introduction

The aim of modification of tRNA by a fluorescent reagent is to obtain information on tRNA structure as well as on tRNA-protein interaction. The fluorescent dye has to bind specifically to one or a few nucleosides in the tRNA [for reviews see 1–4] and the reaction conditions have to be mild so that neither damage to the structure of tRNA nor incorrect recognition resulting in misaminoacylation occurs.


\* Permanent address: Osaka University, Faculty of Pharmaceutical Sciences, Osaka 565, Japan.

Reprint requests to Dr. F. Cramer.

0341-0382/80/0500-0522 \$ 01.00/0

Recently Kopelovich and Wolfe [5] modified unfractionated tRNA from human colon carcinoma and some *Escherichia coli* tRNAs with pyridoxal-5'-phosphate. They suggested that the aldehyde group of the reagent might react with an amino group of the tRNA and the resulting Schiff base might be transformed in a stable covalent C-N linkage by borohydride reduction. In particular guanosine 20 in the dihydrouridine loop of the tRNA seemed to be reactive from inhibition studies with N-acetoxy-2-acetylaminofluorene [6] using unfractionated *E. coli* tRNA. This course of reaction was also suggested from the reaction of the related aldehydes kethoxal and glyoxal with tRNA<sup>Phe</sup> as reported in the literature [7].

Since pyridoxal-5'-phosphate should react mildly [5], is fluorescent [8], and is not very large, we aimed to prepare yeast tRNA<sup>Phe</sup> and tRNA<sup>Tyr</sup> modified in the dihydrouridine loop with this reporter group.



## Experimental

### Materials

Pyridoxal-5'-phosphate was purchased from Boehringer (Mannheim), sodium borohydride, salts and buffer substances (ultrapure grade) from Merck (Darmstadt), RNase T<sub>1</sub>, E.C. 3.1.4.8, from Sankyo (Tokyo, Japan), snake venom phosphodiesterase (1 mg/ml), E.C. 3.1.4.18, and alkaline phosphatase from *E. coli* (1 mg/ml), E.C. 3.1.3.1, from Boehringer (Mannheim), and [<sup>14</sup>C]phenylalanine and [<sup>14</sup>C]tyrosine (50 Ci/mol) from Schwarz BioResearch (Orangeburg, USA). tRNA<sup>Phe</sup> and phenylalanyl-tRNA synthetase, E.C. 6.1.1.20, from yeast were isolated according to [9, 10], tRNA<sup>Tyr</sup> and tyrosyl-tRNA synthetase, E.C. 6.1.1.1, from yeast according to [11].

### Spectroscopy

Ultraviolet absorbance measurements were performed with a Shimadzu double beam spectropho-



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

tometer UV-200 and a Zeiss PMQ-3 spectrophotometer. Relative fluorescence intensities were measured with a modified Farrand MK-1 spectrofluorometer equipped with a Varian-F-80 A X-Y-recorder.

#### Fluorescence titration

100  $\mu$ l samples containing 1  $A_{260}$  unit of tRNA and various amounts of reagent were incubated at 37 °C for 30 min in 20 mM borate buffer, pH 8.0, containing 5 mM  $MgSO_4$ . The reaction mixture was passed through a Sephadex G-25 column (1.7  $\times$  38 cm) equilibrated with water at pH 6.0. The fluorescence intensity of the eluate was measured by using 37  $\mu$ M pyridoxal-5'-phosphate in aqueous solution at pH 6.0 as a standard at 20 °C. The observed fluorescence intensity was then calculated as relative fluorescence intensity per  $A_{260}$  unit of tRNA.

#### Reaction of pyridoxal-5'-phosphate with tRNA

30–300 nmol tRNA (20–200  $A_{260}$  units) were reacted with pyridoxal-5'-phosphate dissolved in water of pH 8.0, in 20 mM borate buffer, pH 8.0, containing 5 mM  $MgSO_4$  in a total volume of 1–2 ml. The reaction was continued at 37 °C for 30–60 min. Then the mixture was cooled in the ice bath and tRNA was precipitated by adding three volumes of cold ethanol in 0.2 M KCl. After centrifugation, the precipitant was washed with cold ethanol followed by vacuum drying. Then it was dissolved in 1 ml of 0.2 M Tris-HCl, pH 7.5, containing 5 mM  $MgCl_2$  and reduced by adding a 3500–7000 fold molar excess of  $NaBH_4$  dissolved in 500  $\mu$ l of cold 0.2 M Tris-HCl, pH 7.5, containing 5 mM  $MgCl_2$ . Reduction was continued for 15–30 min at 0 °C in the dark, and then free  $NaBH_4$  was hydrolyzed by addition of 1 N acetic acid to pH 4 in the ice bath. tRNA was separated from the reagents on a Sephadex G-25 column (1.7  $\times$  38 cm) equilibrated with water. The pooled tRNA fractions were concentrated and loaded on a RPC-5 column (0.7  $\times$  90 cm) equilibrated with 50 mM phosphate buffer pH 7.0, containing 10 mM  $MgSO_4$  and 0.4 M NaCl. After washing with the same buffer, the tRNA was eluted with a linear gradient of 2  $\times$  250 ml, 0.4 M to 1.0 M NaCl in 50 mM phosphate buffer, pH 7.0, containing 10 mM  $MgSO_4$  at 40 °C. The appropriate fractions were pooled and dialyzed against water in the cold room, then these samples were concentrated and reprecipitated with ethanol in 0.2 M KCl to remove any contamination. The pre-

cipitant was dissolved in a small volume of water and desalted on a Biogel P-2 column (1.5  $\times$  30 cm) equilibrated with water. The pyridoxal-5'-phosphate treated tRNAs were stocked at –20 °C.

#### Aminoacylation

100  $\mu$ l of the reaction mixture contained 150 mM Tris-HCl, pH 7.5, 50 mM KCl, 5 mM ATP, 0.05 mM [ $^{14}C$ ]phenylalanine or [ $^{14}C$ ]tyrosine and about 0.2  $A_{260}$  units of tRNA. It was started by adding 3.4  $\mu$ g of phenylalanyl-tRNA synthetase or 10.5  $\mu$ g of tyrosyl-tRNA synthetase. In 10  $\mu$ l aliquots the acid-precipitable radioactivity was determined [9–11].

#### Ribonuclease $T_1$ cleavage of modified tRNA and oligonucleotide separation

3  $A_{260}$  units of tRNA<sup>Phe</sup> treated with  $NaBH_4$  and 15  $A_{260}$  units of tRNA<sup>Phe</sup> treated with pyridoxal-5'-phosphate and  $NaBH_4$  as described above were incubated with 100 and 250 units of RNase  $T_1$  in 1.5 ml of 50 mM Tris-HCl, pH 7.5 at 37 °C for 16 h, respectively. The reaction mixture was adjusted to 20 mM Tris-HCl, pH 7.5 and 7 M urea and then loaded on a DEAE-cellulose column (0.7  $\times$  85 cm) equilibrated with 20 mM Tris-HCl, pH 7.5. The column was eluted with a linear gradient of 2  $\times$  200 ml 0 M to 0.3 M NaCl in 20 mM Tris-HCl, pH 7.5 at a flow rate of 16 ml/h.

34  $A_{260}$  units of tRNA<sup>Tyr</sup> treated with pyridoxal-5'-phosphate and  $NaBH_4$  were digested with 500 units of RNase  $T_1$  and further dealt with as above.

## Results

#### Reaction of tRNA<sup>Tyr</sup> and tRNA<sup>Phe</sup> with pyridoxal-5'-phosphate

tRNA<sup>Tyr</sup> was reacted with a 10–500 fold excess of pyridoxal-5'-phosphate in an aqueous borate buffer in the presence of  $Mg^{2+}$  at pH 8 and the excess of reagent was removed by gel filtration. This pyridoxal-5'-phosphate treated tRNA<sup>Tyr</sup> exhibited fluorescence excitation and emission maxima of 330 nm and 420 nm (Fig. 1), which correspond to the respective maxima of free pyridoxal-5'-phosphate in aqueous solution. The measured fluorescence was normalized to a standard solution of the reagent. The number of pyridoxal-5'-phosphate molecules bound to tRNA<sup>Tyr</sup> was estimated assuming that its




Fig. 1. Relative fluorescence intensity of pyridoxal-5'-phosphate treated  $\text{tRNA}^{\text{Tyr}}$  at various initial molar ratios of reagent to  $\text{tRNA}^{\text{Tyr}}$ . Excitation was done at 330 nm, emission was measured at 420 nm. Insert: uncorrected fluorescence excitation and emission spectra of pyridoxal-5'-phosphate treated  $\text{tRNA}^{\text{Tyr}}$  (in  $10^{-4}$  M aqueous solution at pH 3.8 and  $20^\circ\text{C}$ ). Band widths of 5 and 10 nm at the excitation and emission side, respectively, were used.

fluorescence quantum yield is unchanged upon binding to tRNA as is the quantum yield of pyridoxamine-5'-phosphate upon complex formation with human or bovine serum albumin [8]. As is evident from Fig. 1 saturation of  $\text{tRNA}^{\text{Tyr}}$  was observed at a molar ratio of pyridoxal-5'-phosphate over  $\text{tRNA}^{\text{Tyr}}$  in the range of 50–200.

In contrast the number of pyridoxal-5'-phosphate molecules bound to  $\text{tRNA}^{\text{Tyr}}$  considerably changed when borohydride reduction [12] was performed after the pyridoxal-5'-phosphate treatment and the  $\text{tRNA}^{\text{Tyr}}$  was then purified by RPC-5 chromatography. It eluted in a major fraction corresponding to 95% of the applied  $\text{tRNA}^{\text{Tyr}}$ . The mean number of pyridoxal-5'-phosphate molecules bound to  $\text{tRNA}^{\text{Tyr}}$  in this fraction is 2.3, irrespective of the initial molar ratio of the reagent to  $\text{tRNA}^{\text{Tyr}}$ . No absorbance change was observed at all with this modified yeast  $\text{tRNA}^{\text{Tyr}}$  around 325 nm, whereas

Kopelovich and Wolfe [5] with human and *E. coli* tRNAs reported appearance of an absorption peak at that wavelength.

With yeast  $\text{tRNA}^{\text{Phe}}$  the estimation of the number of pyridoxal-5'-phosphate molecules bound per molecule of  $\text{tRNA}^{\text{Phe}}$  by means of fluorescence was not possible, because of the overlapping emission spectra of pyridoxal-5'-phosphate and the Y base of  $\text{tRNA}^{\text{Phe}}$ . Nevertheless  $\text{tRNA}^{\text{Phe}}$  was treated in the same way as  $\text{tRNA}^{\text{Tyr}}$  and identified by oligonucleotide analysis.

#### Aminoacylation of pyridoxal-5'-phosphate treated $\text{tRNA}^{\text{Phe}}$ and $\text{tRNA}^{\text{Tyr}}$

The aminoacylation was investigated under standard aminoacylation conditions [9–11] and determined to be 1600 pmol tyrosine per  $A_{260}$  unit to  $\text{tRNA}^{\text{Tyr}}$  and 1750 pmol phenylalanine per  $A_{260}$  unit of  $\text{tRNA}^{\text{Phe}}$ . Aminoacylation of the modified tRNAs as described above led to a small enhancement of 2% with  $\text{tRNA}^{\text{Tyr}}$ , whereas a reduction of 8% in aminoacylation was observed with  $\text{tRNA}^{\text{Phe}}$ . Thus the modification of  $\text{tRNA}^{\text{Phe}}$  and  $\text{tRNA}^{\text{Tyr}}$  with pyridoxal-5'-phosphate has only minor influence on the extent of aminoacylation compared to the native tRNAs.

#### Separation of oligonucleotides from ribonuclease $T_1$ digestion of pyridoxal-5'-phosphate treated $\text{tRNA}^{\text{Phe}}$ and $\text{tRNA}^{\text{Tyr}}$

Native  $\text{tRNA}^{\text{Phe}}$  was treated with sodium borohydride, digested with RNase  $T_1$ , and the resulting oligonucleotides were separated on DEAE cellulose (Fig. 2a). Concomitantly  $\text{tRNA}^{\text{Phe}}$  which was first




Fig. 2. Separation of oligonucleotides obtained by digestion with ribonuclease  $T_1$  of a)  $\text{tRNA}^{\text{Phe}}$  treated with sodium borohydride, b)  $\text{tRNA}^{\text{Phe}}$  treated with pyridoxal-5'-phosphate followed by borohydride reduction and c)  $\text{tRNA}^{\text{Tyr}}$  treated with pyridoxal-5'-phosphate followed by borohydride reduction. Relative fluorescence at 420 nm ( $\cdots$ ), UV absorbance at 254 nm ( $---$ ).

reacted with excess pyridoxal-5'-phosphate, subsequently reduced by sodium borohydride, was then digested with RNase T<sub>1</sub> and the resulting oligonucleotides were separated analogously (Fig. 2b). The composition of oligonucleotides and their respective elution volumes are nearly identical in both tRNA<sup>Phe</sup> samples. However, no fluorescent oligonucleotide, resulting from modification with pyridoxal-5'-phosphate, was detected. Only one fluorescent oligonucleotide was observed in both elution patterns and was identified by nucleoside analysis as being the Y base containing dodecanucleotide.

tRNA<sup>Tyr</sup> was treated analogously as tRNA<sup>Phe</sup>, reduced with borohydride and digested in one case (data not shown), and reacted with pyridoxal-5'-phosphate prior to reduction and digested in the other case (Fig. 2c). Again, no fluorescent oligonucleotide, revealing the fluorescence characteristics of pyridoxal-5'-phosphate, could be detected.

### Concluding Remarks

In the present investigation evidence is presented, that pyridoxal-5'-phosphate is not reacting with yeast tRNA<sup>Phe</sup> and tRNA<sup>Tyr</sup> with formation of a stable covalent linkage. Since linking via reduced Schiff base seems to be stable upon RNase T<sub>1</sub> digestion as reported in the case of proflavine and ethidium bromide linkage to tRNA<sup>Phe</sup> [12], we have to conclude that with yeast tRNA<sup>Phe</sup> and tRNA<sup>Tyr</sup> pyridoxal-5'-phosphate may have not so a specific reactivity towards guanosines as has been reported previously [5] for unfractionated human colon carcinoma tRNA and some *E. coli* tRNAs.

### Acknowledgements

Dr. N. Okabe thanks the Alexander von Humboldt-Stiftung for supporting his stay in Germany and Dr. D. Gauss for discussions during the preparation of this manuscript.

- [1] S. H. Kim, *Adv. Enzymol.* **46**, 279–315 (1978).
- [2] J. P. Goddard, *Progr. Biophys. Mol. Biol.* **32**, 233–308 (1977).
- [3] H. G. Faulhammer, M. Sprinzl, and F. Cramer, *Molecular Mechanisms of Biological Recognition* (M. Balaban, ed.), p. 205–225, Elsevier/North Holland, Amsterdam/New York 1979.
- [4] C. R. Cantor and T. Tao, *Procedures in Nucleic Acid Research* (G. L. Cantoni and D. R. Davies, eds.), **Vol. 2**, p. 31–93, Harper and Row, New York 1971.
- [5] L. Kopelovich and G. Wolfe, *Biochemistry* **16**, 3721–3726 (1977).
- [6] P. Pulkrabek, D. Grunberger, and I. B. Weinstein, *Biochemistry* **13**, 2414–2419 (1974).
- [7] M. Litt, *Biochemistry* **10**, 2223–2227 (1971).
- [8] J. E. Churchich and L. Harpring, *Biochim. Biophys. Acta* **105**, 575–582 (1965); J. E. Churchich, *Biochim. Biophys. Acta* **102**, 280–288 (1965).
- [9] D. Schneider, R. Solfert, and F. von der Haar, *Hoppe-Seyler's Z. Physiol. Chem.* **353**, 1330–1336 (1972).
- [10] F. von der Haar, *Eur. J. Biochem.* **34**, 84–90 (1973).
- [11] H. G. Faulhammer and F. Cramer, *Eur. J. Biochem.* **75**, 561–570 (1977).
- [12] W. Wintermeyer and H. G. Zachau, *Methods Enzymol.* **29**, 667–673 (1974); W. Wintermeyer and H. G. Zachau, *FEBS Letters* **18**, 214–218 (1971).