

## New Cembranoids from Burley Tobacco

Ursula Bruemmer, Christian Paulsen, Gunda Spremberg, Friedlieb Seehofer, and Volker Heemann

B. A. T. Cigaretten-Fabriken GmbH, Forschung und Entwicklung, Bahrenfelder Chaussee 139, D-2000 Hamburg 50

Volker Sinnwell

Institut für Organische Chemie und Biochemie der Universität, Martin-Luther-King-Platz 6, D-2000 Hamburg 13

Z. Naturforsch. **36 c**, 1077–1080 (1981); received July 30, 1981

*Nicotiana tabacum*, Solanaceae, Burley Tobacco, Diterpenoids, Cembranoids, *seco*-Lactone

The liquid CO<sub>2</sub>-extract of aged burley tobacco was investigated for its content of diterpenoids, which are considered as important precursors of flavour constituents. Among several cembranoids, already known from various tobacco types, 4,8-dimethyl-8-hydroxy-11-(1-methyl-ethyl)-14-oxo-pentadeca-5,9-diene-4-olide, a new natural product, could be identified, evidently originating from 2,6,11-cembratriene-4,8-diol by oxidative cleavage of the 11,12-double bond with subsequent cyclization to the *seco*-lactone **14**.

Leaves of *Nicotiana tabacum* are covered with a sticky exude, which contains diterpenoids. Depending on the genetic background, tobacco cultivars produce the macrocyclic cembranoids, the carbocyclic labdanoids or both [1, 2]. Biodegradation of these compounds leads to a large number of nor-diterpenoids, which are considered to be important tobacco flavours [3]. To get more insight into these degradation mechanisms, the distribution of diterpenoids in aged burley tobacco was investigated.

Extraction of 30 kg tobacco with liquid CO<sub>2</sub> yielded about 450 g of an extract, containing 44% H<sub>2</sub>O. High-vacuum distillation of the organic layer produced 95 g of a concentrate, with 70% nicotine and 10% neophytadiene as the main constituents. The distillate was further separated by column chromatography, using silicagel. A fraction, enriched with diterpenoids, eluted with 100% diethylether and was subjected to repeated column chromatography, TLC and HPLC, using columns packed with Partisil-10 PAC (Whatman).

Reprint requests to Dr. V. Heemann.

0341-0382/81/1100-1077 \$ 01.00/0

In addition to the (1S, 2E, 4S, 6R, 7E, 11E)- and (1S, 2E, 4R, 6R, 7E, 11E)-2,7,11-cembratriene-4,6-diols (**1**, **2**) [4] and (1S, 2E, 4S, 6E, 8R, 11S)- and (1S, 2E, 4R, 6E, 8R, 11S)-8,11-epoxy-2,6,12(20)-cembratriene-4-ols (**3**, **4**) [5], already found in various tobacco types, several compounds new in burley respectively in tobacco could be identified.

Three of the four possible diastereomers of 2,6,11-cembratriene-4,8-diol (**5**<sub>1–3</sub>) were isolated and the structure proved by MS, <sup>1</sup>H-NMR and <sup>13</sup>C-NMR (Table I).  $\alpha$ - and  $\beta$ -2,6,11-cembratriene-4,8-diol (**5**) were already identified by Rowland *et al.* [6] without precise determination of the stereochemistry of both compounds. The correlation to the  $\alpha/\beta$  nomenclature was based on  $\alpha$ - and  $\beta$ -2,7,11-cembratriene-4,6-diol (**1**, **2**). The present spectral data failed either to relate compound **5**<sub>1</sub>, **5**<sub>2</sub> or **5**<sub>3</sub> to  $\alpha$ - and  $\beta$ -2,6,11-cembratriene-4,8-diol, as published by Rowland *et al.* [6] or to determine the absolute configuration.

Therefore **5**<sub>1–3</sub> were subjected to an x-ray-analysis, the results of which are reported elsewhere.

The two isomeres of 2,7,11-cembratriene-4-ol (**6**, **7**) were isolated for the first time from Virgin tobacco [2]. The stereochemical assignment was achieved by Wahlberg *et al.* [8] to be (1S, 2E, 4S, 7E, 11E)- and (1S, 2E, 4R, 7E, 11E)-2, 7,11-cembratriene-4-ol. The spectral properties of both diastereomers, now isolated from burley were in good accordance with the published data, the quantitative ratio for compound **6** and **7** for the investigated burley being 2:1, as calculated by GC.

8,11-Epoxy-2,6-cembradiene-4,12-diol (**8**) and 11,12-epoxy-2,7-cembradiene-4,6-diol (**9**), isolated now from burley, showed the same spectral data (MS, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR), as published by Behr *et al.* [9, 10] for (1S, 2E, 4S, 6E, 8R, 11S, 12R)-8,11-epoxy-2,6-cembradiene-4,12-diol and (1S, 2E, 4R, 7E, 11S, 12S)-11,12-epoxy-2,6-cembradiene-4,6-diol, both recently identified from Greek tobacco.

The identification of the ketol (**10**), isolated formerly from smoke-cured tobacco [11], is based on comparison with an authentic sample, prepared by CrO<sub>3</sub>-oxydation of **1**. The spectra are identical with those, published by Roberts and Rowland [4].

The *seco*-diketone (**11**) also found in smoke-cured tobacco [11] and the *seco*-aldehyde (**12**), isolated recently from Virgin tobacco, were tentatively identified by their MS and <sup>1</sup>H-NMR features and the structures verified by <sup>13</sup>C-NMR (Table I). The




Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.


Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.


On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.




4S 1  
4R 2




4S 3  
4R 4




5




4S 6  
4R 7




8




9




10




11




12




13



14



15



16

structure of the triol (**13**) was established by its spectral data [13] (MS,  $^1\text{H-NMR}$ ,  $^{13}\text{C-NMR}$ ) to be 2,7,12 (20)-cembratriene-4,6,11-triol, its stereochemistry still remaining unknown. (1S, 2E, 4S, 6R, 7E, 11S)-2,7,12 (20)-cembratriene-4,6,11-triol is said to be found in Turkish tobacco [14].

Furthermore a new natural product was isolated, suggesting the structure of the *seco*-lactone (**14**).

Precise mass determination was accomplished by FD-mass spectra yielding the molecular weight 336, confirming the elemental composition  $\text{C}_{20}\text{H}_{32}\text{O}_4$  [15]. The  $^1\text{H-NMR}$ -spectrum implied the presence of an

Table I.  $^{13}\text{C}$ -NMR chemical shifts and assignments for compounds **5**<sub>1-3</sub>, **10**–**15**.

|                                    | C-1   | C-2                | C-3               | C-4               | C-5   | C-6   | C-7   | C-8               | C-9               | C-10              |
|------------------------------------|-------|--------------------|-------------------|-------------------|-------|-------|-------|-------------------|-------------------|-------------------|
| <b>5</b> <sub>1</sub> <sup>d</sup> | 47.8  | 128.6              | 139.6             | 72.4 <sup>c</sup> | 48.4  | 128.4 | 141.9 | 71.9 <sup>c</sup> | 44.1              | 27.9              |
| <b>5</b> <sub>2</sub> <sup>d</sup> | 47.9  | 128.3              | 140.2             | 72.3 <sup>c</sup> | 48.9  | 127.4 | 142.0 | 72.6 <sup>c</sup> | 43.8              | 28.3              |
| <b>5</b> <sub>3</sub> <sup>d</sup> | 47.8  | 128.8              | 140.7             | 72.6 <sup>c</sup> | 49.1  | 126.6 | 141.7 | 72.7 <sup>c</sup> | 43.0              | 28.4              |
| <b>10</b> <sup>e</sup>             | 46.9  | 126.6              | 136.2             | 72.5              | 54.2  | 201.9 | 131.1 | 159.9             | 31.8 <sup>b</sup> | 28.9 <sup>c</sup> |
| <b>11</b> <sup>e</sup>             | 31.6  | 198.5 <sup>b</sup> | 132.2             | 157.8             | 41.0  | 25.0  | 123.3 | 135.6             | 37.4              | 29.7              |
| <b>12</b> <sup>e</sup>             | 190.9 | 127.4              | 163.4             | 40.7              | 25.7  | 122.3 | 136.9 | 37.7              | 30.7              | 49.1              |
| <b>13</b> <sup>f</sup>             | 49.0  | 130.2              | 138.6             | 72.5              | 53.7  | 65.2  | 131.2 | 138.8             | 33.4 <sup>b</sup> | 31.9 <sup>b</sup> |
| <b>14</b> <sup>e</sup>             | 176.4 | 28.9               | 34.2              | 85.1              | 135.9 | 125.3 | 45.6  | 72.2              | 138.4             | 129.8             |
| <b>15</b> <sup>e</sup>             | 177.0 | 29.5 <sup>a</sup>  | 29.6 <sup>a</sup> | 86.4              | 23.1  | 68.5  |       |                   |                   |                   |

|                       | C-11  | C-12  | C-13              | C-14               | C-15 | C-16              | C-17              | C-18              | C-19              | C-20              |
|-----------------------|-------|-------|-------------------|--------------------|------|-------------------|-------------------|-------------------|-------------------|-------------------|
| <b>5</b> <sub>1</sub> | 123.3 | 131.5 | 37.8              | 23.1               | 33.6 | 20.4 <sup>a</sup> | 19.9 <sup>a</sup> | 27.9 <sup>b</sup> | 27.8 <sup>b</sup> | 14.3              |
| <b>5</b> <sub>2</sub> | 123.1 | 131.5 | 37.8              | 23.2               | 33.7 | 20.8 <sup>a</sup> | 19.9 <sup>a</sup> | 29.9 <sup>b</sup> | 28.4 <sup>b</sup> | 14.3              |
| <b>5</b> <sub>3</sub> | 122.2 | 131.0 | 37.7              | 22.4               | 33.5 | 20.9 <sup>a</sup> | 20.0 <sup>a</sup> | 30.6 <sup>b</sup> | 29.6 <sup>b</sup> | 14.1              |
| <b>10</b>             | 123.5 | 134.6 | 36.3 <sup>b</sup> | 25.7 <sup>c</sup>  | 32.2 | 20.1 <sup>a</sup> | 19.9 <sup>a</sup> | 30.0              | 24.8              | 14.9              |
| <b>11</b>             | 48.8  | 150.4 | 123.6             | 198.0 <sup>b</sup> | 26.9 | 31.6              | 20.5 <sup>a</sup> | 19.1 <sup>a</sup> | 15.9              | 26.5              |
| <b>12</b>             | 133.9 | 132.7 | 142.1             | 114.0              | 18.8 | 32.1              | 20.7 <sup>a</sup> | 19.0 <sup>a</sup> | 16.1              | 17.6              |
| <b>13</b>             | 73.6  | 153.5 | 36.0 <sup>b</sup> | 31.6 <sup>b</sup>  | 33.0 | 21.1 <sup>a</sup> | 19.0 <sup>a</sup> | 31.5              | 16.8              | 110.6             |
| <b>14</b>             | 48.7  | 26.3  | 42.0              | 209.3              | 29.9 | 26.6 <sup>b</sup> | 27.9 <sup>b</sup> | 32.0              | 20.6 <sup>a</sup> | 19.1 <sup>a</sup> |

<sup>a</sup> may be reserved; <sup>b</sup> may be reserved; <sup>c</sup> may be reserved; <sup>d</sup>  $\delta$ -values in  $\text{C}_4\text{D}_8\text{O}_2$ ; <sup>e</sup>  $\delta$ -values in  $\text{CDCl}_3$ ; <sup>f</sup>  $\delta$ -values in  $\text{CD}_3\text{OD}$ .

isopropyl group ( $\delta = 0.84$  ppm, 3H, d,  $J = 6.7$  Hz,  $\delta = 0.89$  ppm, 3H, d,  $J = 6.7$  Hz), and three methyl groups ( $\delta = 1.29$  ppm, 3H, s,  $\delta = 1.52$  ppm, 3H, s,  $\delta = 2.16$  ppm, 3H, s), one of them evidently attached to a carbonylfunction ( $\delta = 2.16$  ppm) and four olefinic protons. This was consistent with the  $^{13}\text{C}$ -NMR properties, which revealed one carbonylfunction ( $\delta = 209.3$  ppm), one carboxylfunction ( $\delta = 176.4$  ppm) and two further oxygene bearing carbons at  $\delta = 72.7$  ppm (s) and  $\delta = 85.1$  ppm (s). The proposed structure was ensured by ozonolysis of (**14**), followed by reduction with  $\text{NaBH}_4$ , yielding the diol (**16**) and the hydroxylactone (**15**). The spectral properties of the diol (**16**) proved to be in accordance with those of an authentic sample and literature data [16]. The hydroxylactone (**15**), prepared by

Grignard reaction of ethyllevulinate with 1-bromopropene followed by ozonolysis and reduction with  $\text{NaBH}_4$ , showed the same spectral data as the product, obtained by ozonolyses of (**14**) [17].

This investigation establishes the complexity of the distribution of diterpenes and supports the degradation schemes recently reported [14]. The identification of the *seco*-lactone (**14**) gives for the first time a strong hint of a similar oxidative cleavage of 11,12-double bond in 4,8-diols (**5**) as reported for 4,6-diols [14].

#### Acknowledgement

Thanks are due to Dr. F. Borchert, Kali-Chemie AG, for the FD-MS.

- [1] A. Colledge, W. W. Reid, and R. Russel, *Chem. Ind.* **1975**, 570.
- [2] W. W. Reid, *Ann. Tabac S.E.I.T.A.* **2**, 145 (1974).
- [3] C. R. Enzell, I. Wahlberg, and A. J. Aasen, *Progress Chem. Natural Products* **34**, 1 (1977).
- [4] D. L. Roberts and R. L. Rowland, *J. Org. Chem.* **27**, 3989 (1962).
- [5] R. L. Rowland *et al.*, *J. Org. Chem.* **29**, 16 (1964).
- [6] R. L. Rowland and D. L. Roberts, *J. Org. Chem.* **28**, 1165 (1963).
- [7] V. Heeman *et al.*, *Beitr. Tabakforsch.* **11**, in press.
- [8] I. Wahlberg *et al.*, *Acta Chem. Scand.* **B 35**, 65 (1981).
- [9] D. Behr *et al.*, *Acta Chem. Scand.* **B 32**, 221 (1978).
- [10] D. Behr *et al.*, *Acta Chem. Scand.* **B 34**, 195 (1980).
- [11] A. Zane, *Phytochemistry* **12**, 731 (1973).
- [12] J. L. Courtney and S. McDonald, *Tetrahedron Let.* **1967**, 459.
- [13]  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ ), Bruker WH 270: 0.84 (6H, t,  $J = 6.7$  Hz), 1.29 (3H, s), 1.72 (3H, d,  $J = 1.3$  Hz), 3.97 (1H, t,  $J = 6.5$  Hz), 4.64 (1H, dt,  $J = 2.1$  and 9 Hz), 4.86 (1H, s), 5.02 (1H, s), 5.20 (1H, d,  $J = 9$  Hz), 5.33 (1H, dd,  $J = 15.5$  Hz,  $J = 6.9$  Hz), 5.43 (1H, d,  $J = 15.5$  Hz). MS  $m/e$  %: 304 (2%); 286 (9%); 268 (6%); 243 (8%); 225 (9%); 159 (17%); 145 (33%); 133 (35%); 119 (41%); 79 (70%); 69 (55%); 55 (75%); 43 (100%); 41 (98%).
- [14] C. R. Enzell and I. Wahlberg, *Recent Adv. Tob. Sci.* **6**, 64 (1980).
- [15] Found ( $\text{M}^+ - \text{H}_2\text{O}$ ): 318.2188; Calc. for  $\text{C}_{20}\text{H}_{30}\text{O}_3$  ( $\text{M}^+ - \text{H}_2\text{O}$ ): 318.2195.  $^1\text{H-NMR}$ : 0.84 (3H, d,  $J = 6.7$  Hz), 0.89 (3H, d,  $J = 6.7$  Hz), 1.29 (3H, s), 1.52 (3H, s), 2.16 (3H, s), 5.33 (1H, dd,  $J = 15.6$  Hz,  $J = 9$  Hz), 5.48 (1H, d,  $J = 15.6$  Hz), 5.63 (1H, d,  $J = 15.6$  Hz), 5.76 (1H, m). MS  $m/e$  (%): 318 (8%), 300 (1%), 285 (4%), 260 (2%), 217 (4%), 197 (30%), 179 (16%), 161 (26%), 140 (87%), 121 (51%), 95 (58%), 71 (38%), 55 (26%), 43 (100%).
- [16] I. Wahlberg *et al.*, *Acta Chem. Scand.* **B 34**, 675 (1980).
- [17]  $^1\text{H-NMR}$ : 1.33 (3H, s), 1.88 (1H, ddd,  $J = 10.0$  Hz,  $J = 6.8$  Hz,  $J = 12.8$  Hz), 2.33 (1H, ddd,  $J = 10$  Hz,  $J = 6.8$  Hz,  $J = 12.8$  Hz), 2.57 (1H, ddd,  $J = 17.8$  Hz,  $J = 10$  Hz,  $J = 5.8$  Hz), 2.67 (1H, ddd,  $J = 17.8$  Hz,  $J = 10$  Hz,  $J = 6.8$  Hz), 3.48 (1H, d,  $J = 12.2$  Hz), 3.67 (1H, d,  $J = 12.2$  Hz). MS  $m/e$  (%): 115 (3%), 112 (1%), 99 (100%), 71 (13%), 56 (11%), 43 (85%).