

Incorporation of ^{15}N -Ammonia into Serotonin in Cotyledons of Maturing Walnuts

Wolfgang Große

Botanisches Institut der Universität zu Köln, Gyrhofstraße 15, D-5000 Köln 41, Bundesrepublik Deutschland

and Francesc Artigas

Instituto de Química Bio-Organica, Consejo Superior de Investigaciones Científicas, Jorge Girona Salgado S/N, Barcelona-34, Spain

Z. Naturforsch. **38c**, 1057–1058 (1983);
received September 22, 1983

Serotonin Biosynthesis, Ammonia Detoxification, ^{15}N Incorporation, Gas Chromatography-Mass-Spectrometry, *Juglans regia*

Cotyledons isolated from maturing seeds of *Juglans regia* incorporate supplied [^{15}N]-ammonia into serotonin (5-HT). A ^{15}N incorporation of about 9% into the indolic nitrogen of 5-HT is detectable by gas chromatography-mass spectrometry following isolation of 5-HT by preparative HPLC. This ^{15}N incorporation supports the hypothesis that biosynthesis and accumulation of 5-HT play a role in ammonia detoxification in maturing walnut seeds.

Introduction

Serotonin (5-HT), the well known central neurotransmitter in mammalian brain, is also detectable in mature seeds of walnuts (*Juglans regia* L.) [1]. The biosynthesis of 5-HT occurs from tryptophan during maturation of these seeds [2] and is regulated by *de novo* synthesis of an aromatic amino acid decarboxylase [3, 4], together with a hydroxylase needed for the transformation of tryptophan to 5-HT.

As with amides, ureids, alkaloids, and other nitrogen-containing secondary plant products [5, 6] the accumulation of 5-HT is considered to be involved in ammonia detoxification [7]. Such a role for 5-HT is supported by a decrease of ammonia content during maturation accompanied by 5-HT accumulation following tryptophan biosynthesis [8] in these seeds.

In order to verify the hypothesis of ammonia detoxification by 5-HT biosynthesis, [^{15}N] NH_4^+ incorporation experiments were carried out and are described below.

Materials and Methods

Plant material

Seeds of *Juglans regia* L. were a generous gift from E. Moll, Botanical Garden of the City of Cologne. 15 weeks after anthesis the seeds were harvested, freed from pericarp and integuments, and placed aseptically on agar plates made from 1% agar in water.

Incubation experiments

8 μmol [^{15}N] NH_4Cl (CEA-France, Gif-sur-Yvette), dissolved in double-distilled water and sterilized by membrane filtration (Sartorius, Göttingen, West Germany, 0.2 μm pore size), were applied to the surface of each cotyledon, which was then incubated aseptically at 25 °C.

Chromatography

Analytical HPLC was carried out as described elsewhere [8] and preparative HPLC as indicated in the figure. The prepacked preparative Whatman column was a generous gift from Dr. D. Strack, Department of Botany, University of Cologne. The samples were prepared by extraction of the seeds with 0.05 N HCl [8] and injections were made using a 100 μl loop. For determination of ^{15}N incorporation into 5-HT a combined gas chromatography-mass spectrometry (in the mass fragmentographic mode) was performed as described elsewhere [9], except for the use of GC capillary columns.

Results and Discussions

Young seeds of walnuts contain about 8 μmol NH_4^+ g^{-1} fresh weight [8]. To prevent toxic concentrations of ammonia accumulating due to the loss of moisture in the subsequent period of ripening, this NH_4^+ must be assimilated through the synthesis of N-containing compounds. The accumulating 5-HT is believed to be such an ammonia detoxification compound, synthesized via glutamine and tryptophan [7].

In order to confirm ammonia consumption by 5-HT biosynthesis, 8 μmol [^{15}N] NH_4Cl were applied to the surface of each of a number of cotyledons isolated aseptically from walnut seeds. After an incubation period of 9 days at 25 °C, 5-HT was extracted from the cotyledons with 0.05 N HCl and

Reprint requests to Dr. W. Große.

0341-0382/83/1100-1057 \$ 01.30/0



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.




Fig. 1. Purification of serotonin (5-HT) by preparative reversed-phase HPLC on a prepacked Whatman Partisil M9-column 10/29 ODS (RP-18). Flow rate: 3 ml min⁻¹. Mobile phase: isocratic 1% aqueous CH₃COOH.

purified by preparative HPLC as indicated in the figure. 5-HT samples – identified by analytical and preparative HPLC – were pooled from successive preparative runs, evaporated to dryness and stored under N₂.

To determine ¹⁵N incorporation into 5-HT, the dried sample was derivatized by perfluoroacylation to yield the pentafluoropropionic derivative of 5-HT (= 5-HT-3PFP) and analysed by gas chromatography-mass spectrometry in the mass fragmentographic mode according to previously described procedures [9]. Since the PFP derivative of the 5-HT synthesized from [¹⁵N]NH₄Cl would contain ¹⁵N atoms, the fragment corresponding to the β-cleavage in the side chain (*m/e* 451 in the standard 5-HT-3PFP) would have a *m/e* of 452. Thus, 5-HT isolated from walnuts should have a $\frac{m/e\ 452}{m/e\ 451}$ ratio higher than that of authentic 5-HT-3PFP.

To test this hypothesis, three ion fragments (*m/e* 451, 452, and 453) were focused, corresponding to a major fragment (451) and its isotopic peaks

m/e + 1 (452) and *m/e* + 2 (453), all of them containing the indolic nitrogen. The ratios between the response at 452/451 and 453/451 respectively were as follows:

Response ratio	5-HT-3PFP standard [%]	5-HT-3PFP from walnuts [%]
$\frac{m/e\ 452}{m/e\ 451}$	20.1 ± 0.2	28.9 ± 0.7
$\frac{m/e\ 453}{m/e\ 451}$	2.4 ± 0.2	4.1 ± 0.1

The results ($\bar{x} \pm S.D.$) were found to be statistically different ($P < 0.001$, Student's *t*-test, $n = 3$) for each response ratio. Taking into account these mean values, the percentage of ¹⁵N incorporation into the 5-HT sample from walnut seeds can be calculated as

8.8% $\left(\text{ratio } \frac{452}{451} \right)$ and 8.6% $\left(\text{ratio } \frac{453}{451} \right)$. These

findings demonstrate very well that cotyledons of maturing seeds are able to use ammonia for 5-HT biosynthesis. Beside ¹⁵N-labeled 5-HT these cotyledons will also have synthesized unlabeled 5-HT from their own unlabeled ammonia and tryptophan. Therefore the detectable ¹⁵N incorporation into 5-HT of about 9% is high enough to verify the hypothesis of ammonia detoxification by 5-HT biosynthesis in maturing walnut seeds.

Acknowledgements

This work was kindly supported by the Deutsche Forschungsgemeinschaft. The authors are also grateful to Dr. P. Macnicol, CSIRO, Canberra, Australia, for reviewing the manuscript and Ms. Helga Tiebel for careful technical assistance.

- [1] E. Kirberger and L. Braun, *Biochim. Biophys. Acta* **49**, 391 (1961).
- [2] L. Bergmann, W. Große, and H. G. Ruppel, *Planta* **94**, 47 (1970).
- [3] W. Große and S. Klapheck, *Z. Pflanzenphysiol.* **93**, 359 (1979).
- [4] S. Klapheck and W. Große, *Z. Pflanzenphysiol.* **100**, 325 (1980).

- [5] K. Mothes, *Naturwissenschaften* **52**, 571 (1965).
- [6] K. Mothes, *Naturwissenschaften* **53**, 317 (1966).
- [7] W. Große, *Phytochemistry* **21**, 819 (1982).
- [8] W. Große, M. Karisch, and P. Schröder, *Z. Pflanzenphysiol.* **110**, 219 (1983).
- [9] F. Artigas and E. Gelpi, *Anal. Biochem.* **92**, 233 (1979).