

Photoaffinity Labeling of Spinach Thylakoids and Cytochrome *b₆/f*-Complex by the Hydrophobic Reagent 3-(Trifluoromethyl)-3-(*m*-[¹²⁵I]iodophenyl)-diazirine

Walter Oettmeier, Hans-Joachim Soll, and Ellen Olschewski

Lehrstuhl Biochemie der Pflanzen, Ruhr-Universität, Postfach 102148, D-4630 Bochum 1, Bundesrepublik Deutschland

Z. Naturforsch. **40c**, 454–456 (1985); received February 25, 1985

Photoaffinity Label, Carbene Reagent, Spinach Thylakoids, Cytochrome *b₆/f*-complex, Membrane Proteins

In isolated spinach thylakoids the hydrophobic photo-reactive probe 3-(trifluoromethyl)-3-(*m*-[¹²⁵I]iodophenyl)-diazirine almost exclusively labels the photosystem I reaction center and the light-harvesting chlorophyll *a/b* protein. In isolated cytochrome *b₆/f*-complex, all four components of the complex get labeled, but to a different extent. The amount of labeling in the protein components is correlated to the number of polypeptide segments embedded in the membrane system.

Introduction

The carbene-generating photoreactive probe [¹²⁵I] TID was recently introduced for the identification of lipid embedded protein segments. Due to its hydrophobic character TID preferentially partitions into the lipid phase of a membrane system and almost exclusively labels integral membrane proteins at the lipid-protein boundary [1–3]. As to photosynthetic systems, TID-labeling so far has only been applied to chromatophores and reaction centers of photosynthetic bacteria, where it was shown to have a high affinity to pigment-polypeptides [4, 5].

We wish to report here that TID in spinach thylakoids only labels the photosystem I reaction center and the light-harvesting chlorophyll *a/b*-protein. In isolated thylakoid cytochrome *b₆/f*-complex, all 4 subunits are labeled by TID, but to a different degree.

Abbreviations: Chl, chlorophyll; CPI, photosystem I reaction center, DNP-INT, 2-iodo-2',4,4'-trinitro-2-methyl-6-isopropylidiphenylether; LHCP, light-harvesting chlorophyll *a/b* protein complex; LDS, lithium dodecylsulfate; TID, 3-(trifluoromethyl)-3-(*m*-iodophenyl)diazirine; Tricine, N-[tris(hydroxymethyl)-methyl]-glycine.

Reprint requests to Dr. W. Oettmeier.

Verlag der Zeitschrift für Naturforschung, D-7400 Tübingen 0341–0382/85/0500–0454 \$ 01.30/0

Materials and Methods

Thylakoids from spinach were prepared according to [6] and cytochrome *b₆/f*-complex according to [7]. [¹²⁵I] TID (spec. activity 10 Ci/mmol) was purchased from Amersham, Braunschweig. For binding experiments, thylakoids corresponding to 100 µg Chl were incubated with TID for 5 min in a medium containing 20 mM Tricine, pH 8.0, and 20 mM MgCl₂. Thylakoids were pelleted at 10000 × *g* and pellet and aliquots of the supernatant assayed for radioactivity in a γ-counter.

For photoaffinity labeling experiments, samples were illuminated for 2 min at 0 °C in a nitrogen atmosphere. They were subsequently subjected to LDS polyacrylamide gel electrophoresis (10–15%). For assay of radioactivity the gel after staining was cut into 1 mm pieces, which were counted in a γ-counter. Alternatively, the gel was exposed to a Kodak X-Omat AR X-ray film at –80 °C for 52 h.

Results and Discussion

As evident from the binding experiment (Fig. 1), TID efficiently partitions into the thylakoid membrane. At each concentration applied, 84% of TID are found in the thylakoid membrane and only 16% in the aqueous phase. Binding of TID to the thylakoid is completely linear to the total concentration of TID applied (Fig. 1). No specific, *i.e.* high affinity binding to a component of the thylakoid can be detected. Contrary, specific binding is observed for in-




Fig. 1. Binding of [¹²⁵I] TID to thylakoid membranes. For conditions, see Materials and Methods.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.




Fig. 2. Photographs (1, 3) and autoradiograms (2, 4) of LDS-polyacrylamide gels (10–15%) of thylakoids (1, 2) labeled by 7.5 nmol/mg Chl and cytochrome b_6/f -complex (3, 4) labeled by 1.2 nmol/nmol cytochrome f [^{125}I] TID. M = marker proteins.

hibitors of photosynthetic electron transport and photoaffinity labels derived thereof (for review see [8]).

However, after covalent attachment of TID to thylakoids by UV-illumination, radioactivity is only found in two major proteins, migrating at approximately 110 kDa and 26–29 kDa, referred to as the photosystem I reaction center (CPI) and the Light-harvesting chlorophyll a/b protein complex (LHCP), respectively [9] (Fig. 2). The radioactivity found in front of the gel is probably due to labeled free lipids and pigments. To establish whether besides CPI and LHCP other proteins within the thylakoid membrane have been labeled by [^{125}I] TID, a gel was cut into 1 mm pieces and each piece assayed for radioactivity. The result is shown in Fig. 3. As can be seen some minor labeling is also found in two photosystem II proteins of 43 and 47 kDa [10] and in some proteins in the 15–20 kDa molecular weight range. However, the labeling of these proteins is small as compared to CPI and LHCP. There are two reasons for the intense labeling of the latter two proteins: (i) they are integral membrane proteins, and (ii) they are highly abundant within the thylakoid membrane (compare their staining intensities in relation to other thylakoid proteins; Lane 1, Fig. 2).

In addition to thylakoids the labeling pattern of [^{125}I] TID in isolated cytochrome b_6/f -complex has been investigated. The cytochrome b_6/f -complex con-

sists out of 4 major peptides: cytochrome f (double band at 33, 34 kDa), cytochrome b_6 (23.5 kDa), the Rieske Fe-S protein (20 kDa), and subunit IV (17.5 kDa) [7] (Fig. 2, lane 3). As is evident from the autoradiogram (Fig. 2, lane 4), all four proteins are tagged by TID, but to a different extent. It should be noted that preincubation of the complex with the inhibitor DNP-INT, which prevents plastoquinone oxidation at the complex [11], prior to addition of TID decreases labeling of all four proteins to about the same extent (data not shown).

A densitometric scan of the autoradiogram (Fig. 2, lane 4) and integration yields the following quantitative radioactivity distribution within the four proteins of the complex: cytochrome f , 10%; cytochrome b_6 , 43%; Fe-S protein, 17%; subunit IV, 30%. Except for the Rieske Fe-S protein, the amino acid sequences of the cytochrome b_6/f -complex proteins are known [12–14]. An amino acid sequence for the Rieske Fe-S protein from *Neurospora* mitochondria has recently been reported [15]. Because the homologies between the cytochromes in the mitochondrial b/c_1 - and the spinach b_6/f -complex are very high [12], a similar situation might exist for the




Fig. 3. Photograph and radioactivity scan of a LDS polyacrylamide gel (10–15%) of thylakoids labeled by 7.5 nmol/mg Chl [^{125}I] TID. The molecular weight numbers indicate the positions of the marker proteins.

Fe-S protein. Hydropathy analysis for the cytochrome b_6 has provided evidence that this protein is arranged in five helical spans through the thylakoid membrane [12, 13]. Three helical spans have been predicted for subunit IV, which is functionally related to cytochrome b_6 [12, 13]. Contrary, only one helical span has been predicted for either cytochrome f [14] or the Fe-S protein (from *Neurospora*) [15]. Thus, the amount of labeling within the four proteins of the cytochrome b_6/f -complex correlates

well with the percentage of membrane intrinsic parts of the respective protein.

In conclusion, TID proves to be an efficient tool for the evaluation of intramembrane parts of integral membrane proteins.

Acknowledgement

This work was supported by Deutsche Forschungsgemeinschaft.

- [1] J. Brunner and G. Semenza, *Biochemistry* **20**, 7174 (1981).
- [2] P. L. Jørgensen and J. Brunner, *Biochim. Biophys. Acta* **735**, 291 (1983).
- [3] J. Krebs, J. Buerkler, D. Guerini, J. Brunner, and E. Carafoli, *Biochemistry* **23**, 400 (1984).
- [4] H. P. Meister, and R. Bachofen, *Advances in Photosynthesis Research* (C. Sybesma, ed.), **Vol. 3**, p. 369, Martinus Nijhoff/Dr. W. Junk Publishers, The Hague, Boston, Lancaster (1984).
- [5] R. Bachofen and H. P. Meister, *Short Reports, 3rd European Bioenergetics Conference*, **Vol. 3A**, p. 57, Congress Edition, Hannover (1984).
- [6] N. Nelson, Z. Drechsler, and J. Neumann, *J. Biol. Chem.* **245**, 143 (1970).
- [7] E. Hurt and G. Hauska, *J. Bioenerg. Biomembr.* **14**, 405 (1982).
- [8] W. Oettmeier, *New Concepts and Trends in Pesticide Chemistry* (P. Hedin, ed.), American Chemical Society Symposium Series, in press (1985).
- [9] J. P. Thornber, *Ann. Rev. Plant Physiol.* **26**, 127 (1975).
- [10] H. Y. Nakatani, B. Ke, E. Dolan, and C. J. Arntzen, *Biochim. Biophys. Acta* **765**, 347 (1984).
- [11] A. Trebst, H. Wietoska, W. Draber, and H. J. Knops, *Z. Naturforsch.* **33c**, 919 (1978).
- [12] W. R. Widger, W. A. Cramer, R. G. Herrmann, and A. Trebst, *Proc. Natl. Acad. Sci. USA* **81**, 674 (1984).
- [13] W. Heinemeyer, J. Alt, and R. G. Herrmann, *Curr. Genet.* **8**, 543 (1984).
- [14] J. Alt and R. G. Herrmann, *Curr. Genet.* **8**, 551 (1984).
- [15] U. Harnisch, H. Weiss, W. Sebald, *Short Reports, 3rd European Bioenergetics Conference*, **Vol. 3A**, p. 131, Congress Edition, Hannover (1984).