

Effect of Tuftsin on the Phagocytotic Activity of the Unicellular *Tetrahymena*. Does Primary Interaction Develop Imprinting?

G. Csaba, Valéria László, and P. Kovács

Department of Biology, Semmelweis University of Medicine, Nagyvárad tér 4. H-1445 Budapest, Hungary

Z. Naturforsch. **41c**, 805–806 (1986);
received December 30, 1985

Tetrahymena, Tuftsin, Phagocytosis, Imprinting

The natural mammalian phagocytosis stimulator tetrapeptide tuftsin stimulated also the phagocytosis of the unicellular *Tetrahymena*. The selectivity of the *Tetrahymena* was not complete, as threonine-free tuftsin provoked a greater response. Treatment with tuftsin did not develop functional imprinting, however the binding capacity of cell membrane significantly increased after the first treatment.

Tuftsin, a tetrapeptide (γ -globulin fraction) composed of threonine, lysine, proline and arginine, stimulates the phagocytotic activity of the leucocytes [1, 2]. Since histamine has a similar effect on the leucocytes, and also stimulates the phagocytosis of the unicellular *Tetrahymena* [3, 4], it seemed worthwhile to examine whether tuftsin, too, had an influence on the unicellular, and whether it would, like histamine, induce imprinting [3–5] at that low phylogenetic level.

Preliminary experiments performed with different (10^{-10} – 10^{-5} M) concentrations of tuftsin (prepared by Dr. S. Bajusz, Institute of Drug Research, Budapest) had shown that 10^{-7} M tuftsin was most suitable for the purpose of the experiment. *Tetrahymena pyriformis* GL cells, maintained in Bactotrypton and yeast extract containing medium at 28 °C were exposed to that concentration, and were simultaneously fed Chinese ink, for 10 min, were spread on slides, dried and examined for the number of Chinese ink containing vacuoles. The mean value was calculated for each group and was related to the control value as 1 (phagocyte coefficient = PC). The Fig. 1 shows mean values calculated from five replica experiments. Cells treated for 24 h for the first time with tuftsin were examined for vacuole counts seven days later, either with or without 10-min reexposure to 10^{-7} M tuftsin. The threonine-free fraction of tuftsin was also tested for influence on phagocytosis. The




Fig. 1. Effect of tuftsin (T+) or threonine deprived tuftsin (T-) on the phagocytotic coefficient (PC) of *Tetrahymena* immediately after the first treatment (C/T+, C/T-) or one week after it with (T+/T+, T-/T-) or without (T+/C, T-/C) repeated treatment. Values are given as percent related to the untreated control (C/C) as 1. S = $p < 0.05$.

intergroup differences were evaluated for significance with Student's t-test.

Tuftsin stimulated the phagocytosis of the *Tetrahymena* similarly to histamine. It follows that uni- and multicellular organisms are equally responsive to stimulators of phagocytosis. The response of the unicellular (*Tetrahymena*) is nevertheless less selective, for it showed a similar (or even greater) response to histidine in earlier studies [6], and to the threonine-deprived tuftsin fragment in the present studies, compared to the stimulant proper. Moreover, primary interaction with tuftsin or its threonine-free fragment did not durably stimulate the phagocytic activity, to judge from decline of the latter in a later stage of the experiment.

To obtain information on the imprinting potential of tuftsin, we compared the binding of FITC-labeled tuftsin between primary and second exposure. While neither tuftsin nor threonine deprived tuftsin imprinted the *Tetrahymena* for a functional (phagocytotic activity) change, the former imprinted it for a greater binding, to judge from a 25.1% ($p < 0.05$) increase in binding capacity at reexposure. Imprinting by the threonine-deprived fraction was more than twice as effective (55.9%, $p < 0.01$). These observations support the implication [7] that the binding and action of active molecules do not necessarily show a parallelism.

Reprint requests to Prof. Csaba.

Verlag der Zeitschrift für Naturforschung, D-7400 Tübingen
0341–0382/86/0700–0805 \$ 01.30/0



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

- [1] V. A. Najjar and K. Nishioka, *Nature* **228**, 672 (1970).
- [2] K. Nishioka, A. Constantopoulos, P. S. Satoh, and V. A. Najjar, *Biochem. Biophys. Res. Comm.* **47**, 172 (1972).
- [3] G. Csaba, *Biol. Rev. (Cambridge)* **55**, 47 (1980).
- [4] G. Csaba, *Horm. Metab. Res.* **16**, 329 (1984).
- [5] G. Csaba, *Internat. Rev. Cytol.* **95**, 327 (1985).
- [6] G. Csaba and Zs. Darvas, *BioSyst.* **19**, 55 (1986).
- [7] M. D. Hollenberg, *Pharmac. Rev.* **30**, 393 (1979).