

The C-Glycosylflavone Pattern of *Passiflora incarnata* L.

Hans Geiger

Institut für Chemie der Universität Hohenheim,
Garbenstraße 30, D-7000 Stuttgart 70
and

Kenneth R. Markham

Chemistry Division, D.S.I.R., Private Bag, Petone,
New Zealand

Z. Naturforsch. **41c**, 949–950 (1986);
received April 10, 1986


Passiflora incarnata L., Passifloraceae,
Flavone-C-glycosides

From the herbage of *Passiflora incarnata* have been isolated eight flavone-C-glycosides based on apigenin and luteolin, six of them had not been found before in that plant. The constitution of all compounds is proven.

The herbage of *Passiflora incarnata*, which is widely used medically [1], has been reported to contain the C-glycosyl-flavones vitexin, isovitexin (**1**), orientin, isoorientin (**2**), and saponarin (**9**) [1–3]. When one of us tried to isolate some reference substances from that plant, it became evident, that the C-glycosyl flavone pattern was more complex than had been reported earlier.

Our findings are summarized in Table I. The starting material was a commercial dry extract, kindly provided by Messrs Dr. Willmar Schwabe, Karlsruhe, F.R.G., and manufactured from the herbage of *Passiflora incarnata* grown on the premises of that firm. The separation of the flavone glycosides was effected by chromatography on a column of polyamide-6 with a water-acetone gradient (sequence of elution with considerable overlapping: **7** ≈ **8** > **5** >

Table I. Structures, yields, *M_r* determined by FD-MS, and chromatographic behaviour of flavone C-glycosides from *Passiflora incarnata*.



	Substitution pattern				Yield [mg/ 100 g]	<i>M_r</i>	hR _f -values				Fluorescence	
	R ¹	R ²	R ³	R ⁴			Cellulose 15% HAc	TBA	Polyamide NM	WEBP	un- treated	NA
Isovitexin (1)	glc	H	H	H	216	432	38	73	16	35	d	gr
Isoorientin (2)	glc	H	H	OH	158	448	25	51	02	32	d	y
Isovitexin-2''-β-D-glucopyranoside (3)	soph	H	H	H	531	594	75	66	14	63	d	gr
Isoorientin-2''-β-D-glucopyranoside (4)	soph	H	H	OH	170	610	68	50	06	63	d	y
Schaftoside (5)	glc	H	ara	H	250	564	45	34	19	70	d	gr
Isoschaftoside (6)	ara	H	glc	H	152	564	34	28	17	67	d	gr
Vicenin-2 (7)	glc	H	glc	H	164	594	39	26	09	72	d	gr
Lucenin-2 (8)	glc	H	glc	OH	57	610	31	15	05	72	d	y
Saponarin (9)	glc	glc	H	H	not detected	594	60	52	17	74	d	gr

glc = β-D-glucopyranosyl; ara = α-L-arabinopyranosyl; soph = sophorose; Cellulose = precoated sheets Polygram CEL 400 (Macherey and Nagel, D-5160 Düren). Polyamide = F1700-Mikropolyamide sheets (Schleicher and Schüll, D-3345 Dassel). 15% HAc = 15% aqueous acetic acid; TBA = tert. butanol/acetic acid/water (3:1:1). NM = nitro methane/methanol (7:3); WEBP = water/ethanol/butanone/2,4-pentanedione (65:15:15:5). NA = sprayed with 0.5% methanolic diphenylboric acid – β-aminoethyl ester.



6 > **3** \approx **4** \gg **1** \approx **2**). For the separation of the individual glycosides the following systems were used: Sephadex LH 20/20% aqueous methanol (**7** \approx **8** \gg **5** \approx **6** \gg **3** \approx **4** \gg **1** > **2**); Polyclar AT/20% aqueous methanol (**7** \gg **8**; **3** \gg **4**; **5** > **6**); and Sephadex LH 20 with acetone/methanol/water (2:1:1) (for the elimination of polymers, which appear in the forerun). Finally the pure glycosides were crystallized from aqueous methanol.

The ^{13}C NMR spectra of **5**, **6**, **7** and **8** were identical with published spectra of these compounds [4, 5]. In combination with the FDMS and cochromatographic results this constitutes proof of identity for **7** and **8**; but does not distinguish the two isomers, **5** and **6**. However, the differences between the R_F -values of these two isomers are such, that on this basis the distinction can be made with confidence [5].

Compounds **3** and **4** are also known natural products [6, 7], but no published ^{13}C NMR data were available for comparison. These spectra (Table II) are readily interpreted in that the sugar region in both is almost superimposable with that in the spectrum of vitexin-2''-O- β -D-glucopyranoside [8], while in the aromatic region the signals of **3** and **4** are as expected for a 6-C-glycosylated apigenin or luteolin [9]. Thus **3** and **4** are identified as isovitexin- and isoorientin-2''-O- β -D-glucopyranoside respectively. Compounds **1** and **2** were identified by direct comparison with authentic samples.

Saponarin (**9**), of which an authentic sample isolated from *Saponaria officinalis* L. [10] was at hand, could not be detected in *P. incarnata*, either by TLC of the extract, or by its colour reaction with I_2/KI [10, 11] in fresh leaves. Vitexin and orientin, which have also been reported to occur in *P. incarnata* could only be detected by TLC in amounts too small for isolation.

Table II. ^{13}C NMR spectra of isovitexin-2''- β -D-glucopyranoside (**3**) and isoorientin-2''- β -D-glucopyranoside (**4**)*.

Carbon No.	Signals 3	4
2	163.4 ⁺	163.5 ⁺
3	102.7	102.9
4	181.9	182.0
5	161.1 ⁺	161.2 ⁺
6	107.9	108.1
7	160.9 ⁺	163.5 ⁺
8	93.4	93.4
9	156.4	156.6
10	102.7	103.4
1'	121.1	121.6
2'	128.4	113.4
3'	115.9	145.9
4'	161.1 ⁺	149.8
5'	115.9	116.2
6'	128.4	119.1
1''	71.1	71.3
1'''	105.3	105.3
2''	81.5	81.7
2'''	74.6	74.8
3''	78.3	78.5
3'''	76.3	76.6
4'', 4'''	70.3, 69.3	70.6, 69.5
5''	80.8	81.1
5'''	76.3	76.6
6'', 6'''	61.3, 60.3	61.5, 60.5

* Solvent DMSO- d_6 ; assignments bearing the same superscript in any one spectrum may be reversed.

Acknowledgements

Thanks are due to G. Schwinger for recording the FD mass spectra and to Dr. W. Wong of Chemistry Division for running the NMR spectra. Hans Geiger thanks the "Fonds der Chemischen Industrie" for financial support.

- [1] J. Lutomski, E. Segiet, K. Szpunar, and K. Grisse, *Pharmazie in unserer Zeit* **10**, 45 (1981).
- [2] H. Schilcher, *Z. Naturforsch.* **23b**, 1393 (1968).
- [3] J. Löhdefink, *Dtsch. Apoth. Ztg.* **116**, 557 (1976).
- [4] B.-G. Österdahl, *Acta Chem. Scand.* **B32**, 93 (1978).
- [5] B.-G. Österdahl, *Acta Chem. Scand.* **B33**, 400 (1979).
- [6] R. Tschesche and K. Struckmeyer, *Chem. Ber.* **109**, 2901 (1976).
- [7] K. Hostettmann and A. Jacot-Guillarmod, *Helv. Chim. Acta* **58**, 130 (1975).
- [8] E. Besson, G. Dellamonica, J. Chopin, K. R. Markham, M. Kim, H.-S. Koh, and H. Fukami, *Phytochemistry* **24**, 1061 (1985).
- [9] K. R. Markham, *Techniques of Flavonoid Identification*, Academic Press, London 1982.
- [10] G. Barger, *J. Chem. Soc.* **89**, 1210 (1906).
- [11] G. Barger and E. Field, *J. Chem. Soc.* **101**, 1394 (1912).