

Identification of the Sex Pheromone of Eggplant Borer *Leucinodes orbonalis* Guenée (Lepidoptera: Pyralidae)

Zhu Pingchou*, Kong Fanlei, Yu Shengdi, Yu Yongqing, Jin Shuping, Hu Xinhua, and Xu Jianwei

Jiangsu Institute of Pheromones and Hormones Jintan, Jiangsu, China

Z. Naturforsch. **42c**, 1347–1348 (1987); received May 11, 1987

Sex Pheromone, Identification, *Leucinodes orbonalis*, Lepidoptera, 11-Hexadecenyl Acetate

(*E*)-11-Hexadecenyl acetate was identified from sex pheromone gland extract of female eggplant borer. The acetate synthesized in the laboratory showed high attractant activity in the field.

The eggplant borer, *Leucinodes orbonalis* Guenée (Lepidoptera: Pyralidae), is a serious pest of eggplant in various regions of China [1]. The control of this pest is troubled with the fact that the chemical insecticides do not penetrate into the fruit where the pests stay. The female sex pheromone identified and synthesized as follows might be a prospective candidate for population monitoring or direct control.

The abdominal tips of virgin female moths were immersed in *n*-heptane for 6 h. This crude extract was fractionated by preparative TLC on a silica gel plate and developed with *n*-hexane-ethyl ether (4:1). The *absorbate* was cut into five fractions and washed with treated acetone. In the field test, high bioactivity


was shown by the fraction of R_f 0.6–0.8 corresponding to a long chain acetate, and no notable activity by the remaining fractions.

GC-MS(EI) analysis of the active fraction from TLC separation on SE-54 capillary column recorded a major peak at 17'49" which possessed the following fragment ions, m/e 222 ($M^+ - CH_3COOH$), 194 ($C_{14}H_{26}^+$), 180 ($C_{13}H_{24}^+$), 166 ($C_{12}H_{22}^+$), 152 ($C_{11}H_{20}^+$), 138 ($C_{10}H_{18}^+$), 124 ($C_9H_{16}^+$), 110 ($C_8H_{14}^+$), 96 ($C_7H_{12}^+$), 82 ($C_6H_{10}^+$) and 61 ($CH_3COOH_2^+$). These data indicated that the natural pheromone is a *n*-hexadecenyl acetate [2].

To determine the location of the double bond, microozonolysis [3, 4] was carried out. GC-MS (methane CI) of the ozonolysis product on a PEG-20M capillary column showed a major peak at 6'36" with parent ion at m/e 229 (MH^+), and less ion at m/e 169 ($MH - CH_3COOH$) and 151 ($MH - CH_3COOH - H_2O$). These data indicated that the ozonolysis product of pheromone was 11-acetoxyundecanal, and thus showed that the double bond of the pheromone is located at 11-position.

The configuration of the double bond was examined by comparing the GC retention time (R_t) of the natural pheromone to that of synthetic (*E*)- and (*Z*)-11-hexadecenyl acetate (**I_a** and **I_b**). R_t of the synthetic **I_a** and **I_b** on a DEGS capillary column were 16'40" and 17'20" respectively, while that of natural pheromone was 16'40". Thus, *Z* configuration was defined.

Synthesis of **I_a** and **I_b** was carried out as follows:



where **a**---NaH/1,2-EDA,
d---Na/Liq NH₃,

b---DHP, H₃O⁺,
e---H₂, P-2Ni,

c---*n*-BuLi/BuBr,
f---AcOH-AcCl.

Reprint requests to Zhu Pingchou.

Verlag der Zeitschrift für Naturforschung, D-7400 Tübingen
0341-0382/87/1100-1347 \$ 01.30/0



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

In the electroantennagram (EAG) "puff" test at 10 μg level, the males responded much stronger to synthetic \mathbf{I}_a than to \mathbf{I}_b . In field trapping, \mathbf{I}_a was found to be very attractive to male moths, and more males

were captured by traps baited with 300 to 500 μg \mathbf{I}_a than by six alive females. This result proved that \mathbf{I}_a is the main component of the eggplant borer sex pheromone.

- [1] a) Plant Diseases and Insect Pests in China, **Vol. 2**, p. 1493, Agriculture Publ. Beijing 1981;
b) Zhu Shuxun, Kunchong Zhishi (Popular Entomology) **1980**, 262, 271;
c) Yang Ziqi, Kunchong Zhishi (Popular Entomology) **1982** (2), 20;
d) Hu Quanyao, Zhiwu Baohu (Plant Protection) **1983** (4), 22.
- [2] Ando Tetsu, Kishino Kenichi, Tatsuki Sadahiro, and Takahashi Nobutaka, Agric. Biol. Chem., **44**, 765 (1980).
- [3] M. Beroza and B. A. Bierl, Anal. Chem. **39**, 1131 (1967).
- [4] B. F. Nesbitt, P. S. Beevor, D. R. Hall, R. Lester, and J. R. Williams, J. Chem. Ecol. **5**, 385 (1980).