

Inhibition of Bacteriochlorophyll Biosynthesis by Gabaculin (3-Amino, 2,3-dihydrobenzoic Acid) and Presence of an Enzyme of the C₅-Pathway of δ -Aminolevulinic Acid Synthesis in *Chloroflexus aurantiacus*

Monika Kern and Jobst-Heinrich Klemme

Institut für Mikrobiologie der Universität Bonn,
Meckenheimer Allee 168, D-5300 Bonn 1, Bundesrepublik Deutschland

Z. Naturforsch. **44c**, 77–80 (1989); received September 23, 1988

Chloroflexus aurantiacus, δ -Aminolevulinic acid, Bacteriochlorophyll, Gabaculin, Glutamate-C₅-pathway

Biosynthesis of bacteriochlorophyll *c* and *a* in the thermophilic phototrophic prokaryote, *Chloroflexus aurantiacus* Ok-70-fl, was strongly inhibited by the antibiotic gabaculin (3-amino 2,3-dihydrobenzoic acid), an inhibitor of the glutamate-C₅-pathway of δ -aminolevulinic acid (ALA) synthesis. The key enzyme of the Shemin-pathway of ALA formation, ALA synthase (EC 2.3.1.37), was not detected in cell extracts of *Chl. aurantiacus*. However, the extracts catalyzed ALA formation from glutamate 1-semialdehyde, a reaction being highly sensitive to gabaculin.

Introduction

In all organisms studied so far, δ -aminolevulinic acid (ALA) is the precursor of tetrapyrrole compounds (chlorophylls, heme pigments, cobalamins, a.o.). Two different pathways of ALA biosynthesis are known: (a) The "Shemin-pathway", i.e. the condensation of succinate and glycine to ALA [1]; and (b) the "C₅-pathway" in which glutamate is converted to ALA by the following enzyme-catalyzed steps: (i) activation of glutamate by binding to a specific tRNA; (ii) hydrogenation of the activated glutamate to glutamate 1-semialdehyde; and (iii) transamination of the latter compound to ALA [2, 3]. Research conducted during the last two decades has shown that phototrophic prokaryotes use both pathways. Several species of the phototrophic non-sulfur purple bacteria (*Rhodospirillaceae*) synthesize ALA from succinate and glycine [4–6], whereas cyanobacteria, green sulfur bacteria and purple sulfur bacteria mostly lack ALA synthase (EC 2.3.1.37), the enzyme catalyzing the condensation of succinyl-coenzyme A and glycine to ALA, and synthesize the latter compound *via* the C₅-pathway [6–9].

The thermophilic phototrophic prokaryote, *Chloroflexus* (*Chl*). *aurantiacus*, has a somewhat ambiguous position with respect to the mechanisms of ALA synthesis. Unlike the cyanobacterium

Anacystis nidulans, the green sulfur bacterium *Chlorobium limicola* and the non-sulfur purple bacterium *Rhodospseudomonas palustris*, *Chloroflexus aurantiacus* did not incorporate radioactivity into ALA from either [¹⁴C]glycine plus succinate or from [¹⁴C]glutamate [6]. Therefore, we have reinvestigated the mechanism of ALA synthesis in *Chl. aurantiacus*. It will be shown in this paper that the antibiotic gabaculin (3-amino 2,3-dihydrobenzoic acid), a potent inhibitor of ALA synthesis in cyanobacteria but not in non-sulfur purple bacteria [10], strongly inhibits bacteriochlorophyll synthesis in growing *Chl. aurantiacus* cultures, and that cell extracts of the organism catalyze the conversion of glutamate 1-semialdehyde to ALA.

Materials and Methods

Organisms and culture conditions

Chl. aurantiacus Ok-70-fl was obtained from Dr. Karin Schmidt, Institute for Microbiology, University of Göttingen. The organism was grown photosynthetically (conditions as described in [11]) at 52–55 °C in a yeast extract containing medium (S medium according to [12]).

Rhodobacter (*Rhb*). *sphaeroides* (formerly called *Rhodospseudomonas sphaeroides*) DSM 636 was obtained from the German Collection of Microorganisms, Braunschweig. The organism was grown photosynthetically at 30–32 °C in a malate-(NH₄)₂SO₄-medium [13] supplemented with a vitamin mixture [14].

Reprint requests to J.-H. Klemme.

Verlag der Zeitschrift für Naturforschung, D-7400 Tübingen
0341–0382/89/0100–0077 \$ 01.30/0



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

Determination of bacteriochlorophyll and protein content of cells

Bacteriochlorophyll (BChl) *c* and *a* was determined in acetone-methanol (7:2 mixtures) extracts of cells using the specific absorption coefficients $\epsilon_{666} = 86 \text{ liter} \cdot \text{g}^{-1} \cdot \text{cm}^{-1}$ (BChl *c*) and $\epsilon_{769} = 75 \text{ liter} \cdot \text{g}^{-1} \cdot \text{cm}^{-1}$ (BChl *a*) [15, 16]. The protein content of cells was assayed by a biuret method [17].

Preparation of cell extracts

Bacterial cells harvested at the end of the logarithmic growth phase were washed once in 50 mM Tris-HCl, pH 8.2 (*Chl. aurantiacus*) or pH 7.2 (*Rhb. sphaeroides*) and resuspended (either immediately or after storage at -18°C) in the same buffer supplemented with 10 mM MgCl_2 and 2 mM mercaptoethanol (for ALA synthase assay). Cell homogenates were prepared by passing the suspensions twice through a French pressure cell (20,000 psi) and were freed of intact cells and cell debris by centrifugation at $15,000 \times g$ (4°C) for 40 min. Protein concentrations of cell extracts were determined according to [18].

Determination of δ -aminolevulinic acid (ALA)

For separation of ALA from substances interfering with its colorimetric assay, test mixtures were treated with a cation exchanger (Dowex 50W-X8, Na^+ -form) contained in small polypropylene columns ("Econo" columns, Biorad, München) [19].

ALA eluted from the columns was converted with ethylacetoacetate to ALA-pyrrole which was assayed colorimetrically with Ehrlich's reagent (*p*-dimethylaminobenzaldehyde) [20].

Enzyme assays

ALA synthase (EC 2.3.1.37) activity of *Rhb. sphaeroides* extracts was determined at 37°C in a reaction mixture (1 ml) described in [21]. *Chl. aurantiacus* extracts were tested at 52°C under almost identically conditions (Tris-HCl buffer of pH 8 instead of pH 7.2). The enzyme reaction was stopped after maximally 90 min by adding 0.1 ml of 1 M citric acid. Precipitated protein was removed by centrifugation and ALA in the supernatant assayed as described above.

Glutamate 1-semialdehyde aminotransferase (EC 5.4.3.8) activity of *Chl. aurantiacus* extracts was

assayed at 52°C in the following reaction mixture (1.1 ml): 100 mM imidazole-HCl, pH 8; 2.5 mM dithioerythritol; 5 mM MgCl_2 ; 10 mM Na-levulinate; 0.7–1.4 mg of a glutamate 1-semialdehyde preparation; 10 vol.% glycerol; 4–10 mg *Chl. aurantiacus* protein; and 10 μM gabaculin where indicated. The enzyme reaction was stopped after 10 min and ALA was determined as described above.

ALA dehydratase (EC 4.2.1.24) activity was determined in 3 ml reaction mixtures [22] at 37°C (*Rhb. sphaeroides*) or 52°C (*Chl. aurantiacus*). The reaction product, porphobilinogen (PBG), was assayed colorimetrically as PBG-pyrrole with Ehrlich's reagent [20].

Enzyme activities are expressed in terms of units (U). 1 unit (1 $\mu\text{mol}/\text{min}$) is equivalent to 16 nkat.

Chemicals

δ -Aminolevulinic acid, coenzyme A (CoA), coomassie brilliant blue, dithioerythritol, Dowex 50W-X8, Ehrlich's reagent, and gabaculin were obtained from Serva, Heidelberg; ethylacetoacetate and levulinic acid from Sigma, München; ATP, pyridoxal-5'-phosphate and succinyl-CoA-synthetase from Boehringer, Mannheim. A preparation of glutamate 1-semialdehyde-hydrochloride was kindly provided by Dr. C. G. Kannangara, Carlsberg Laboratory, Copenhagen. All other chemicals were obtained from Merck, Darmstadt.

Results and Discussion

Gabaculin (3-amino 2,3-dihydrobenzoic acid), an "antibiotic" produced by *Streptomyces toyocaensis*, was originally introduced as a neurotropic inhibitor of brain γ -aminobutyrate (GABA) transaminase [23]. The diagnostic value of gabaculin in studies of the enzymatic mechanism of ALA biosynthesis rests with the fact that also the C_5 -pathway enzyme, glutamate-1-semialdehyde transaminase, is strongly and specifically inhibited [10, 24]. As shown in Table I, gabaculin (at a concentration of 2 μM) almost completely arrested the synthesis of bacteriochlorophyll (BChl) in a photosynthetically growing culture of *Chl. aurantiacus*.

Since cellular growth was not completely blocked in the presence of the inhibitor (due to the fact that saturating light intensity was used to illuminate the culture) the specific BChl content of the inhibited

Table I. Effect of gabaculin (3-amino 2,3-dihydrobenzoic acid) on bacteriochlorophyll synthesis and photosynthetic growth of *Chloroflexus aurantiacus* Ok-70-fl. Cultures were grown photosynthetically using 1 l-screw cap bottles filled to 4/5 of the total volume with S-medium [12] and flushed with sterile N₂ via cotton-stoppered gas in-and outlets. Samples were withdrawn via a glass-tube reaching to the bottom of the flask.

	Growth stage	Protein content [mg/ml]	Total BChl content [µg/ml]	Spec. BChl content [µg/mg protein]
Control culture without gabaculin	initial	0.035	0.45	12.8
	after 112 h	0.38	18	47.4
Culture with 2 µM gabaculin	initial	0.053	0.50	9.5
	after 112 h	0.20	0.54	2.7

culture decreased from initially 9.5 to 2.7 µg/mg protein (compared to an increase from 12.8 to 47.4 µg/mg protein in a non-inhibited culture). Note, that the ratio of BChl *c*/BChl *a* (values of 8:1 recorded in non-inhibited cultures) was only slightly changed in a gabaculin-treated culture. Thus, synthesis of both BChl types was affected by the inhibitor.

As expected, control experiments with photosynthetically growing cultures of *Rhb. sphaeroides* (which synthesizes ALA via the Shemin-pathway) showed that gabaculin (up to 10 µM) had no measurable effect on the specific BChl-content of the cells (26 and 25.6 µg BChl *a*/mg protein in control and inhibitor-containing cultures, respectively) or on the growth rate. Contrary to results with *Rhb. sphaeroides* in which levulinic acid (added at a concentration of 50 mM) caused a 60% decrease of the specific BChl-content accompanied by an accumulation of ALA in the culture medium (up to 5 nmol ALA/ml during the first 20 h of growth), levulinate (50 mM) caused no ALA excretion in *Chl. aurantiacus* cultures.

Under our experimental conditions (see section Materials and Methods) not even traces of ALA synthase activity were detected in cell extracts of *Chl. aurantiacus*. In control experiments with cell extracts of photosynthetically grown *Rhb. sphaeroides*, ALA synthase activities in the range of 1 to 2.5 U/g protein (= 16 to 40 nkat/g protein) were readily measured.

A possible cause of the failure to detect ALA synthase activities in *Chl. aurantiacus* extracts could

have been a very unfavourable activity ratio ALA synthase/ALA dehydratase. The latter enzyme was detected at activity levels of 1 to 2 U/g protein in extracts of *Chl. aurantiacus* (for comparison: ALA dehydratase activities in cell extracts of *Rhb. sphaeroides* were in the range of 20 to 25 U/g protein). Like other ALA dehydratases from phototrophic microorganisms [10], the *Chl. aurantiacus* enzyme was inhibited by levulinic acid. Analysis of reaction kinetics at various ALA and inhibitor concentrations showed that levulinate inhibited the enzyme competitively with respect to ALA ($K_i = 4$ mM). However, even when the inhibitor was added at concentrations of 100 to 200 mM, no ALA formation was recorded with *Chl. aurantiacus* extracts in ALA synthase assay mixtures.

Importantly, cell extracts of *Chl. aurantiacus* catalyzed the conversion of glutamate 1-semialdehyde to ALA (Table II), a reaction being highly sensitive to gabaculin. Thus, *Chl. aurantiacus* contains an aminotransferase (system) similar or identical to the glutamate 1-semialdehyde aminotransferase of the C₅-pathway [24].

Our experimental results have made it very likely that in the thermophilic phototrophic prokaryote, *Chloroflexus aurantiacus*, δ-aminolevulinic acid (ALA) is synthesized via the glutamate-C₅-pathway. Together with previous investigations of other laboratories [6–9] this finding shows that – with the exception of some genera of the non-sulfur purple bacteria [4–6] – the majority of phototrophic prokaryotes does not use the Shemin-pathway of ALA

Table II. Conversion of glutamate-1-semialdehyde (GSA) to δ -aminolevulinic acid catalyzed by unfractionated cell extracts of *Chloroflexus aurantiacus* Ok-70-fl. Reaction mixtures (1.1 ml) with 100 mM imidazol-HCl, pH 8; 2.3 mM dithioerythritol; 4.5 mM MgCl_2 ; 10 mM levulinic acid (Na^+ -salt); 9 vol.% glycerol and 4–10 mg extract protein were incubated for 10 min at 52 °C. ALA in the reaction mixtures was analyzed as described in the Materials and Methods section.

Reaction mixture	nmol ALA formed
500 μl extract + 700 μg GSA	20.7
500 μl extract + 1400 μg GSA	36.3
200 μl extract + 700 μl GSA	17.0
500 μl extract + 700 μg GSA + 10 μM gabaculin	3.2
700 μg GSA	2.5

synthesis. The recent demonstrations of C_5 -pathway activities in the archaebacterium *Methanobacterium thermoautotrophicum* [25] and the thermophilic *Clostridium thermoaceticum* [26] document that the C_5 -pathway is much more widespread among organisms than had been expected when this reaction sequence was first described by Beale [2].

Acknowledgements

The authors thank Dr. C. G. Kannangara, Carlsberg Laboratory, Copenhagen, for kindly providing a preparation of glutamate-1-semialdehyde, and Dr. Reidun Sirevåg, Department of Biology, University of Oslo, for valuable advice with culture techniques.

- [1] B. F. Burnham, in: *Metabolic Pathways* (D. M. Greenberg, ed.), vol. 3, pp. 403–537, Academic Press, New York 1969.
- [2] S. I. Beale, *Annu. Rev. Plant. Physiol.* **29**, 95–120 (1978).
- [3] A. Schön, G. Krupp, S. Gough, S. Berry-Lowe, C. G. Kannangara, and D. Söll, *Nature* **322**, 281–284 (1986).
- [4] J. Lascelles, in: *Bacterial Photosynthesis* (H. Gest, A. San Pietro, and L. P. Vernon, eds.), pp. 35–52, The Antioch Press, Yellow Springs, Ohio 1963.
- [5] M. S. Wright, R. D. Cardin, and A. J. Biel, *J. Bacteriol.* **169**, 961–966 (1987).
- [6] T. Andersen, T. Briseid, T. Nesbakken, J. Ormerod, R. Sirevåg, and M. Thorud, *FEMS Microbiol. Lett.* **19**, 303–306 (1983).
- [7] J. McKie, C. Lucas, and A. J. Smith, *Phytochemistry* **20**, 1547–1549 (1981).
- [8] K. M. Smith and M. S. Huster, *J. Chem. Soc.* **1**, 14–16 (1987).
- [9] T. Oh-hama, H. Seto, and S. Miyachi, *Arch. Biochem. Biophys.* **246**, 192–198 (1986).
- [10] R. C. Houtl, D. Rees, L. J. Rogers, and A. J. Smith, *Arch. Microbiol.* **146**, 57–62 (1986).
- [11] G. Laakmann-Ditges and J.-H. Klemme, *Arch. Microbiol.* **144**, 219–221 (1986).
- [12] B. K. Pierson and R. W. Castenholz, *Arch. Microbiol.* **100**, 283–305 (1974).
- [13] J. G. Ormerod, K. S. Ormerod, and H. Gest, *Arch. Biochem. Biophys.* **94**, 449–463 (1961).
- [14] M. T. Madigan, S. R. Petersen, and T. D. Brock, *Arch. Microbiol.* **100**, 97–103 (1974).
- [15] R. G. Feick, M. Fitzpatrick, and R. C. Fuller, *J. Bacteriol.* **150**, 905–915 (1982).
- [16] M. van der Rest and G. Gingras, *J. Biol. Chem.* **249**, 6446–6453 (1974).
- [17] L. H. Stickland, *J. Gen. Microbiol.* **5**, 698–703 (1951).
- [18] M. M. Bradford, *Anal. Biochem.* **72**, 248–254 (1976).
- [19] T. G. Cooper, *Biochemische Arbeitsmethoden*, Walter de Gruyter, Berlin 1981.
- [20] D. Mauzerall and S. Granick, *J. Biol. Chem.* **219**, 435–446 (1956).
- [21] S. Tuboi and S. Hayasaka, *Arch. Biochem. Biophys.* **146**, 282–290 (1971).
- [22] D. Shemin, in: *Methods in Enzymology* (S. P. Colowick and N. O. Kaplan, eds.), vol. 17A, pp. 205–211, Academic Press, New York 1970.
- [23] K. Kobayashi, S. Miyazawa, A. Terahara, H. Mishima, and H. Kurihara, *Tetrahedron Lett.* **7**, 537–540 (1976).
- [24] J. K. Hooper, A. Kahn, D. E. Ash, S. Gough, and C. G. Kannangara, *Carlsberg Res. Commun.* **53**, 11–25 (1988).
- [25] H. C. Friedmann, R. K. Thauer, S. P. Gough, and C. G. Kannangara, *Carlsberg Res. Commun.* **52**, 363–371 (1987).
- [26] T. Oh-hama, J. J. Stolowich, and A. I. Scott, *FEBS Lett.* **228**, 89–93 (1988).