

Occurrence of Jasmonic Acid in the Red Alga *Gelidium latifolium*

Marina V. Krupina

Moscow State University "M. W. Lomonossow",
Biological Faculty, Section of Biological Monitoring,
Leninskij Gori, 139445 Moscow, USSR

and

Wilfried Dathe

Institut für Biochemie der Pflanzen, Weinberg 3,
O-4050 Halle/Saale, Bundesrepublik Deutschland

Z. Naturforsch. **46c**, 1127–1129 (1991);
received June 27, 1991

Jasmonic Acid, Plant Growth Regulator,
Gelidium latifolium, Rhodophyta

The growth regulators (–)-jasmonic acid (JA) and its 7-isomer were identified by GC-MS in the red alga *Gelidium latifolium*. The ratio of JA:7-iso-JA was approximately 93:7. The endogenous level amounted to 0.7 µg JA/g fresh weight.

Introduction

The plant growth regulator (–)-jasmonic acid (JA), isolated as a plant growth inhibitor from the pericarp of *Vicia faba* [1], and its methyl ester, a potent senescence promotor from *Artemisia absinthium* [2], are widely distributed within higher plants [3]. However, there are very few reports of its occurrence in lower plants. Aldridge *et al.* [4] described JA as a product of the fungus *Lasiodiplodia theobromae* (Pat.) Griff. *et* Maubl. and Miersch *et al.* [5] isolated from the same fungal species (synonym *Botryodiplodia theobromae* Pat.) its isomer (+)-7-iso-jasmonic acid (7-iso-JA). The green alga *Chlorella pyrenoidosa* (strain 211/8b) possesses the complete enzyme system for JA biosynthesis, but JA could not be found as native compound in this organism [6].

Recently, Ueda *et al.* [7] isolated minute quantities of JA (1.5 ng/g dry weight) from *Euglena gracilis* Z., an eukaryotic algal flagellate (Chlorophyta). Minute levels of JA and JAMe were also detected in a *Chlorella* sp. and JA was found in the

prokaryotic *Spirulina maxima* (Cyanophyceae), too [7, 8]. Furthermore, JA and related compounds were also identified in several *Equisetum* species (Pteridophyta) [9].

Within our research programme on the physiology of macrophytic algae [10, 11] we have begun to investigate the endogenous levels of plant growth regulators, including JA and related compounds. Here we describe the identification of JA and its 7-isomer in the red alga *Gelidium latifolium*.

Materials and Methods

The branched red alga *Gelidium latifolium* Born. (Rhodophyta) was collected from the Black Sea (Bay of Sewastopol, Krim peninsula) to a depth of 50 cm. The algae (length: 3–5 cm, 45 g fresh weight) were frozen, homogenized with MeOH using a blender, filtered and the remaining tissue extracted twice with 80% MeOH. The combined MeOH extracts were evaporated to aqueous, acidified to pH 3.0, partitioned with EtOAc and evaporated to dryness. This extract was subsequently chromatographed on a column of DEAE-Sephadex A-25 (1.1 × 45 cm; [12]). The fractions eluted with 0.25 and 0.5 M HOAc in 80% MeOH were monitored by TLC (1/5 of each fraction, silica gel GF₂₅₄, CHCl₃:EtOAc:acetone:HOAc = 40:10:5:1). The JA-containing fractions were combined, methylated with ethereal diazomethane, purified on Adsorbex RP 18 (40 µm, 400 mg, Merck) with an increasing gradient of MeOH in water (5% steps, from 40% MeOH) and again monitored by TLC (hexane:EtOAc:HOAc = 60:40:1). The JA fractions (60–65% MeOH) were evaporated to dryness and analyzed by GC and GC-MS applying the following conditions: GC – steel column (2 m × 4 mm), Supelcoport (100–120 mesh) coated with OV 225 (3%), carrier gas: N₂ 45 ml/min, column temperature: 180 °C. GC-MS – steel column (1.5 m × 2 mm), Gaschrom Q (100–120 mesh) coated with OV 225 (3%), carrier gas: He 15 ml/min, column temperature: 180 °C; 70 eV.

For GC the sample was dissolved in 10 µl benzene and 1/20 injected (8 replicates), for GC-MS 1/10 was injected. The remaining sample was reduced by NaBH₄ [13] and again analyzed by GC (conditions as above, except: column temperature – 190 °C, N₂ – 35 ml/min) to determine the ratio between JA and its 7-stereoisomer (JAMe is reduced

Abbreviations: JA, (–)-jasmonic acid; 7-iso-JA, (+)-7-iso-jasmonic acid; CA, (+)-cucurbitic acid; 6-epi-CA, (+)-6-epi-cucurbitic acid; 7-iso-CA, (+)-7-iso-cucurbitic acid; 6-epi-7-iso-CA, (+)-6-epi-7-iso-cucurbitic acid; Me, methyl ester of the corresponding compound.

Reprint requests to Dr. W. Dathe.

Verlag der Zeitschrift für Naturforschung, D-7400 Tübingen
0939–5075/91/1100–1127 \$ 01.30/0



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

to 6-epi-7-iso-CAMe ($R_t = 7.8$ min) and 7-iso-CAMe ($R_t = 7.8$ min), 7-iso-JAMe is reduced to 6-epi-CAMe ($R_t = 9.3$ min) and CAMe ($R_t = 10.2$ min); the structural data and natural occurrence of which were recently reported [14]).

Results and Discussion

After chromatography of the EtOAc extract on DEAE-Sephadex A-25 we detected in fractions eluted with 0.25 M HOAc a JA like spot by monitoring on TLC ($R_f = 0.43$). After methylation, purification on Adsorbex RP 18 and TLC (1/5) we detected JAMe in fractions eluted with 60–65% MeOH. In GC two peaks occurred, the first one ($R_t = 9.3$ min) corresponded to authentic JAMe (32 μg ; 0.7 $\mu\text{g/g}$ fresh weight) and the second peak ($R_t = 11.1$ min) to authentic 7-iso-JAMe (1.85 μg ; 0.04 $\mu\text{g/g}$ fresh weight). The ratio between JAMe and 7-iso-JAMe amounted to 94.3:5.7. The identity with JAMe [1] and 7-iso-JAMe [5] was proved by GC-MS ($R_{\text{JAMe}} = 4.0$ min, $R_{\text{7-iso-JAMe}} = 4.6$ min). MS (JAMe) m/z (rel. int.): 224 (M^+ , 30), 206 (6), 193 (14), 156 (24), 151 (36), 135 (17), 109 (22) and 83 (100). MS (7-iso-JAMe) m/z (rel. int.): 224 (M^+ , 17), 206 (8), 193 (7), 156 (16), 151 (29), 135 (11), 109 (23) and 83 (100).

The ratio between JAMe and 7-iso-JAMe determined from the NaBH_4 -reduced sample, in order to avoid isomerization of 7-iso-JAMe to JAMe during GC [15], was 93.4:6.6 (6-epi-7-iso-CAMe +

7-iso-CAMe and 6-epi-CAMe + CAMe) and corresponded very well to the GC determination of JAMe and 7-iso-JAMe. The ratio of JA:7-iso-JA in young fruits of *Vicia faba* was determined to be about 65:35 [13], and in black and green tea between 30:70 and 70:30 depending on the tea type [15]. A fungal strain of *Botryodiplodia theobromae* isolated from orange fruits forms exclusively 7-iso-JA [5]. The biosynthetic pathway of JA in plants should logically yield 7-iso-JA [16, 17]. Isomerization to the JA configuration may take place after any of the β -oxidation steps following the formation of phytodienic acid. In comparison to JA the 7-iso-JA is the more active compound in bioassays [18–20], and its methyl ester seems to be the essential odoriferous agent [21]. Possibly, only 7-iso-JA is the important biologically active compound, and it is therefore necessary to quantify it simultaneously to JA.

The occurrence of JA in *Gelidium latifolium*, *Euglena gracilis* Z., *Chlorella* sp. and in *Spirulina maxima* [7, 8] indicates that in addition to being present in higher plants [3] it is also distributed in lower plants, or at least these lower plants are capable of forming JA [6].

Acknowledgements

We thank Mrs. Christine Kuhnt for GC and GC-MS analyses and Mrs. Monika Krohn for reliable technical assistance.

- [1] W. Dathe, H. Rönsch, A. Preiss, W. Schade, G. Sembdner, and K. Schreiber, *Planta* **155**, 530–535 (1981).
- [2] J. Ueda and J. Kato, *Plant Physiol.* **66**, 246–249 (1980).
- [3] A. Meyer, O. Miersch, C. Büttner, W. Dathe, and G. Sembdner, *J. Plant Growth Regul.* **3**, 1–8 (1984).
- [4] D. G. Aldridge, S. Galt, D. Giles, and W. B. Turner, *J. Chem. Soc. (C)* **1971**, 1623–1627.
- [5] O. Miersch, A. Preiss, G. Sembdner, and K. Schreiber, *Phytochemistry* **26**, 1037–1039 (1987).
- [6] B. A. Vick and D. C. Zimmerman, *Plant Physiol.* **90**, 125–132 (1989).
- [7] J. Ueda, K. Miyamoto, T. Sato, and Y. Momotani, *Agric. Biol. Chem.* **55**, 275–276 (1991).
- [8] J. Ueda, K. Miyamoto, M. Aoki, T. Hirata, T. Sato, and Y. Momotani, *Bull. Univ. Osaka Prefecture, Ser. B* **43**, 103–108 (1991).
- [9] W. Dathe, O. Miersch, and J. Schmidt, *Biochem. Physiol. Pflanzen* **185**, 83–92 (1989).
- [10] K. S. Burdin, W. V. Gusev, M. V. Krupina, and I. B. Saveljev, *Vestnik Moscow Univ., Ser. 16 Biol.*, No. 3, 3–10 (1980).
- [11] K. S. Burdin, M. V. Krupina, and F. B. Saveljev, *Vestnik Moscow Univ., Ser. 16 Biol.*, No. 2, 35–42 (1990).
- [12] R. Gräbner, G. Schneider, and G. Sembdner, *J. Chromatogr.* **121**, 110–115 (1976).
- [13] O. Miersch, A. Meyer, S. Vorkefeld, and G. Sembdner, *J. Plant Growth Regul.* **5**, 91–100 (1986).
- [14] W. Dathe, C. Schindler, G. Schneider, J. Schmidt, A. Porzel, E. Jensen, and I. Yamaguchi, *Phytochemistry* **30**, 1909–1914 (1991).
- [15] A. Kobayashi, M. Kawamura, Y. Yamamoto, K. Shimizu, K. Kubota, and T. Yamanishi, *Agric. Biol. Chem.* **52**, 2299–2303 (1988).
- [16] B. A. Vick and D. C. Zimmerman, *Biochem. Biophys. Res. Commun.* **111**, 470–477 (1983).
- [17] B. A. Vick and D. C. Zimmerman, *Plant Physiol.* **75**, 458–461 (1984).
- [18] A. Meyer, O. Miersch, S. Vorkefeld, and G. Sembdner, *Acta Univ. Agric. (Brno), Fac. Agron.* **33**, 471–474 (1985).
- [19] O. Miersch and G. Sembdner, *Acta Univ. Agric. (Brno), Fac. Agron.* **33**, 101–107 (1985).
- [20] G. Herrmann, H.-M. Kramell, R. Kramell, R. Weidhase, and G. Sembdner, in: *Conjugated Plant Hormones, Structure, Metabolism and Function* (K. Schreiber, H. R. Schütte, G. Sembdner, eds.), pp. 315–322, VEB Deutscher Verlag der Wissenschaften, Berlin 1987.
- [21] T. E. Acree and R. Nishida, *Semiochemistry: Flavors and Pheromones*, pp. 33–45, Walter de Gruyter & Co., Berlin, New York 1985.