

Additional Sulfur Compounds from the Anal Glands of the Striped Polecat, *Ictonyx striatus* (Mustelidae, Mammalia)

J. W. Wheeler^a, L. Nyalley^a, D. M. Davis^b and P. J. Weldon^c

^a Department of Chemistry, Howard University, Washington, D.C. 20059

^b Department of Small Mammals, National Zoological Park, Smithsonian Institution, Washington, D.C. 20008

^c Conservation and Research Center, Smithsonian Institution, 1500 Remount Road, Front Royal, VA 22630

Z. Naturforsch. **52c**, 283–285 (1997);
received December 16, 1996/February 4, 1997

Ictonyx striatus, Striped Polecat, Anal Gland Secretions, 3-Ethyl-1,2-pentanedithiolane, 1,3-Pentanedithiol

Two sulfur compounds have been identified in the anal gland secretions of captive-raised adults of the striped polecat (*Ictonyx striatus*), an African mustelid. 3-Ethyl-1,2-pentanedithiolane was observed in the secretions of an adult male and an adult female. 1,3-Pentanedithiol was observed in the male; this compound has not previously been reported from mustelid anal glands.

Introduction

The paired anal glands of skunks, ferrets, and other mustelids are known to produce volatile, sulfur-containing compounds that are thought to repel predators (e.g., Andersen and Bernstein, 1980), mark territories (Gorman, 1980), and/or signal alarm (Brinck *et al.*, 1978). A headspace analysis of the glandular exudate of a free-ranging male striped polecat (*Ictonyx striatus*), a large African mustelid, revealed 2-ethylthiacyclobutane as the major volatile component (Apps *et al.*, 1988). Nine other sulfur-containing compounds, some tentatively characterized by their mass spectra, were indicated. We report here on the identification of two additional compounds in the anal gland secretions of *I. striatus*, 3-ethyl-1,2-pentanedithiolane and 1,3-pentanedithiol.

Material and Method

Anal gland secretions were obtained from a 11.3-year-old male and a 12.5-year-old female *I. striatus*, born and raised at the National Zoological Park, Washington, D.C. Animals were fed Feline Diet (Zu-Preem Nutritional Products, Topeka, KS), mice, fish, rib bone, eggs, and fruit. Secretions were obtained by palpating the anal glands and collecting the exuded fluids in glass vials with Teflon-lined caps. A few milliliters of CH₂Cl₂ were added to the vials before they were placed on dry ice. The secretions were extracted in *n*-hexane and kept frozen (−4 °C) until analysis.

Samples were analyzed by capillary GC/MS using a 30 m × 32 mm SPB-1 column (1.0 μm coating) connected to a 5890 Hewlett-Packard gas chromatograph programmed from 50° to 300 °C at 12 °C/min. The gas chromatograph was connected to a Hewlett-Packard 5971A mass spectrometer. One drop of secretion extract was diluted in 3 ml of hexane, and 1 μl of this solution was injected onto the column. Compounds were identified by their mass spectra.

To confirm the identity of 3-ethyl-1,2-pentanedithiolane in the secretions, it was synthesized by reducing 1-chloro-3-pentanone to its corresponding alcohol with NaBH₄, brominating the alcohol with PBr₃, and treating this product with sulfur and Na₂S. Details of the synthesis follow. A solution of 1.4 g of NaBH₄ in 25 ml of cold water was added drop-wise to 15 ml of 1-chloro-3-pentanone solution in 25 ml of water, which was cooled in an ice bath. The mixture was stirred for 30 min and allowed to rise to room temperature. Thirteen ml of 2.5 N HCl were added to the mixture which was then extracted with 30 ml of diethyl ether. The ether extract was washed once with water, dried over MgSO₄, and concentrated on a rotary evaporator to yield 13.9 g of 1-chloro-3-pentanol, the identity of which was confirmed by IR spectroscopy.

1-Chloro-3-pentanol was placed in a flask immersed in a mixture of crushed ice and Na₂SO₄ (−7 °C). Fifteen ml of PBr₃ were added dropwise to the flask and the mixture was allowed to sit overnight at room temperature. The flask was fitted with a fractionating column and condenser and

Reprint requests to Prof. Wheeler.
Telefax: (202) 8065442.

0939–5075/97/0300–0283 \$ 06.00 © 1997 Verlag der Zeitschrift für Naturforschung. All rights reserved.

N



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

then distilled under reduced pressure by water aspiration. The distillate collected at 98–101 °C was cooled to 0 °C, washed with a 50% aqueous H₂SO₄ (0 °C), and then dried over anhydrous K₂CO₃. A total of 7.3 g of 3-bromo-1-chloropentane was collected.

In the last step, 13 ml of 3-bromo-1-chloropentane obtained in the previous step were placed into a flask with 60 ml of N,N-dimethylformamide (DMF) and the mixture was stirred. Sulfur (1.3 g) and sodium sulfide nonahydrate (9.6 g) were suspended in 40 ml of DMF and added to the reaction mixture, which was stirred at room temperature for two days. The mixture was then heated at 80–85 °C for 24 h. The hot solution was poured into a beaker filled with 100 g of ice and 50 ml of water. The mixture was extracted twice with 40 ml portions of hexane. The combined organic layer was washed with water, dried over MgSO₄, and concentrated on a rotary evaporator. The odorous, yellow product was transferred to a flask fitted with a distillation column. The mixture was heated and 3.5 ml of 3-ethyl-1,2-pentanedithiolane were collected at 127–130 °C.

A 1.7 ml sample of the synthesized 3-ethyl-1,2-pentanedithiolane was used to synthesize 1,3-pentanedithiol. The former compound was added dropwise to a slurry of 1.8 g of LiAlH₄ in 32 ml of anhydrous diethyl ether. The mixture was stirred at room temperature for 2 h, and then refluxed for 30 min. The reaction was cooled to room temperature and 3 ml of water were slowly added. Excess 10% aqueous H₂SO₄ was added to dissolve the aluminum salts. An additional 15 ml of diethyl ether were added and the two ether layers were collected. The aqueous layer was washed with two additional 15 ml portions of diethyl ether. The combined organic layers were washed with 20 ml of water, dried over anhydrous MgSO₄, and concentrated on a rotary evaporator. The product was characterized by GC/MS and IR spectroscopy.

Results and Discussion

GC chromatograms of the male and the female secretions indicated at least ten components each.

Compound **1**, which was observed in both sexes, displayed the following *m/z* (rel. int.) values: 134(93), 119(5), 105(30), 101(13), 69(100), 64(37), 55(32), 45(69) and 41(20). These values are consistent with 3-ethyl-1,2-pentanedithiolane (Fig. 1), as confirmed by comparison with the synthetic compound. Compound **2**, which was observed only in the male, displayed the following *m/z* (rel. int.) values: 136(93), 107(3), 102(80), 87(30), 84(28), 75(48), 73(100), 69(90), 61(58), 60(98), 55(70), 47(98), and 45(90). These values are the same as those observed for synthetic 1,3-pentanedithiol (Fig. 1).




Fig. 1. The structural formulae of (**1**) 3-ethyl-1,2-pentanedithiolane and (**2**) 1,3-pentanedithiol.

Apps *et al.* (1988) observed that an isomer of ethyldithiacyclopentane was abundant in the anal gland secretions of a free-ranging male *I. striatus*, but the structural details of this component were not resolved. We identified this compound as 3-ethyl-1,2-pentanedithiolane. Dithiolanes are known from the anal glands of several other mustelids, most notably *Mustela* spp. (Crump, 1980a,b; Buglass *et al.*, 1991; Brinck *et al.*, 1983; Schildknecht and Birkner, 1983). 3-Ethyl-1,2-pentanedithiolane, however, has previously been reported only in the stoat (*M. erminea*) (Crump, 1980a,b; Buglass *et al.*, 1991; Brinck *et al.* 1983), where it was indicated as a minor component in the females' secretions (compound **8** in: Crump, 1980a), and in the mink (*M. vison*) (Schildknecht *et al.*, 1981). 1,3-Pentanedithiol, which we observed only in our male *I. striatus*, has not previously been reported from mustelid anal glands. Further studies are needed to determine whether this compound arises from the reduction of 3-ethyl-1,2-pentanedithiolane.

Acknowledgements

L. Newman assisted in obtaining secretions. J. S. Waterhouse and W. F. Wood commented on the manuscript.

- Andersen K. K. and Bernstein D. T. (1980), Sulfur compounds in mustelids. In: Natural Sulfur Compounds: Novel Biochemical and Structural Aspects, (D. Cavallini, G. E. Gaull and V. Zappia, eds.) Plenum Press, New York, 399–406.
- Apps P. J., Viljoen H. W., Pretorius V. and Rohwer E. R. (1988), Volatile components of the anal gland secretion of the striped polecat *Ictonyx striatus*. S.-Afr. Tydskr. Dierk. **23**, 136–137.
- Brinck C., Gerrell R. and Odham G. (1978), Anal pouch secretion in mink *Mustela vison*: chemical communication in Mustelidae. Oikos **30**, 68–75.
- Brinck C., Erlinge S. and Sandel M. J. (1983), Anal gland secretion in mustelids: a comparison, J. Chem. Ecol. **9**, 727–745.
- Buglass A. J., Darling F. M. C. and Waterhouse J. S. (1991), Dimethyldithiolanes in the anal sac secretion of the weasel, *Mustela nivalis*. Z. Naturforsch. **46c**, 166–167.
- Crump D. R. (1980a), Thietanes and dithiolanes from the anal gland of the stoat (*Mustela erminea*). J. Chem. Ecol. **6**, 341–347.
- Crump D. R. (1980b), Anal gland secretion of the ferret (*Mustela putorius forma furo*). J. Chem. Ecol. **6**, 837–844.
- Gorman M. L. (1980), Sweaty mongooses and other smelly carnivores. Symp. Zool. Soc. Lond. **45**, 87–105.
- Schildknecht H. and Birkner C. (1983), Struktur und Wirkung der Musteliden-Ökomone, III: Analyse der Analbeutelsekrete Mitteleuropäischer Musteliden. Chemiker-Zeitung **107**, 267–270.
- Schildknecht H., Birkner C. and Krauß D. (1981), Struktur und Wirkung der Musteliden-Ökomone, III: Erweiterte Analyse der Analbeutelsekrete des Nerzes *Mustela vison* L. Chemiker-Zeitung **105**, 273–284.

