

A 5,6-Dihydro-isopyoverdin from *Azomonas macrocytogenes**

R. Michalke, K. Taraz, H. Budzikiewicz^a,
Ph. Thonart, Ph. Jacques^b

^a Institut für Organische Chemie der Universität zu
Köln, Greinstr. 4, D-50939 Köln, Germany

^b Centre Wallon de Biologie Industrielle,
Faculté Universitaire des Sciences Agronomiques,
B-5030 Gembloux et Université de Liège
Bâtiment B 40, B-4000 Liège, Belgique

Z. Naturforsch. **52c**, 855–857 (1997);
received September 3, 1997

Azomonas macrocytogenes, Biogenesis, Isopyoverdin,
Siderophore

From the culture medium of *Azomonas macrocytogenes* a 5,6-dihydro-isopyoverdin could be isolated which includes the same peptide chain as the accompanying isopyoverdins (azoverdins). As analogous pairs had been encountered in the pyoverdin series where a dihydro derivatives are considered to be the immediate precursors of the pyoverdins, the same biogenetic sequence can thus be assumed for the iso-series.

Introduction

In preceding publications we obtained evidence that the pyoverdin chromophore **1** is derived from the ferribactin chromophore **2** which in turn is a condensation product of D-Tyr and L-Dab (Böckmann *et al.*, 1997). **2** should also be the branching point for the formation of the recently discovered isopyoverdins **3**. If the assumption is correct that **1** and **3** are formed by analogous biogenetic steps starting from **2**, the same type of intermediates should be found and the stereochemistry of the chiral center at C-3 of **3** should be S. The correct stereochemistry was proved recently (Michalke *et al.*, 1997). We wish now to report the isolation of a 5,6-dihydro-isopyoverdin (dihydro-azoverdin) **4** from the culture medium of *Azomonas macrocytogenes* ATCC 12334.

Abbreviations: Common aminoacids, 3-letter code; Dab, 2,4-diaminobutyric acid; Hse, homoserine; (OH)Orn, N⁵-hydroxy Orn; FAB-MS, fast atom bombardement mass spectrometry.

* Part LXXIV of the series "Bacterial constituents". For part LXXIII see Budzikiewicz *et al.* (1997).

Reprint requests to Prof. Dr. H. Budzikiewicz.
Telefax No. +49-221-470-5057.

Experimental Procedures

For bacterial growth, spectroscopy, amino acid analysis, decomplexation, details of chromatography, etc. see Michalke *et al.*, 1996.

Isolation and characterization of 5,6-dihydro-azoverdin G

The culture filtrate was worked up as described earlier (Michalke *et al.*, 1996). After separation of azoverdin A by chromatography on DEAE Sephadex A-25 the fraction containing azoverdin and azoverdin G was rechromatographed on CM Sephadex C-25 with a 0.02 N pyridinium acetate buffer (pH 5.0, isocratic), detection at 340 and 405 nm. After a brown fraction (azoverdin and azoverdin G) a violet fraction containing ferri-5,6-dihydro-azoverdin G could be eluted which was rechromatographed under the same conditions. The Fe³⁺-complex shows absorption maxima at 250, 318 and 535 nm (pH 3.0) and 250, 318 and 517 nm (pH 6.8), resp., while the free dihydro-compound shows a pH-independent absorption at 301 nm. The molecular mass of **4** was determined by FAB-MS as 1122 u, that of the Fe³⁺-complex as 1175 u. From an aminoacid analysis follows the presence of L-Dab, L-Glu, D- and L-Hse, D- and L-(OH)Orn and D-Ser as had been observed for azoverdin G.


Transformation of dihydro-azoverdin G (**4**) into azoverdin G.

10 mg **4** were dissolved in 5 ml H₂O and stirred for 24 hrs with 10 mg PtO₂. The color of the solution changed from violet to brown. The resulting azoverdin G was purified and de-complexed as described above. It was found to be identical with authentic azoverdin G by all spectroscopic evidence.

Results and Discussion

4 has the same aminoacid composition as azoverdin G and both the ¹H- and ¹³C-NMR data of the amino acid portion and of the Glu side chain correspond to those of the latter. Hence the structure of these parts of **4** and of azoverdin G are identical. However, the molecular mass of **4** is 2





units higher and the UV/Vis data correspond to those of 5,6-dihydro-pyoverdins. Accordingly, the ^1H - and ^{13}C -data (Tables I and II) of the positions 4a to 10a of the chromophore match those of the 5,6-dihydro-pyoverdin Pp2 (Gwose *et al.*, 1992) while those of the positions C-1 to C-3 agree with those of azoverdin G (Michalke *et al.*, 1996) in accordance with the position of the carboxyl group

Table I. ^1H NMR data of the chromophores of 5,6-dihydro-azoverdin G (**4**), (H₂O, pH 3.0) azoverdin G (H₂O, pH 4.3) and 5,6-dihydro-pyoverdin Pp2 (D₂O, pH 3.0).

	5,6-dihydro-azoverdin G (4)	azoverdin G	5,6-dihydro-pyoverdin Pp2
Chr-1	3.71/4.01	3.86/4.40	5.28
Chr-2	2.48	2.58	2.50/2.57
Chr-3	4.59	4.60	3.23/3.68
Chr-5	5.10	—	5.50
Chr-6	3.03	7.81	3.03/3.06
Chr-7	6.80	6.88	6.81
Chr-10	6.85	6.94	6.85

Table II. ^{13}C NMR data of the chromophores of 5,6-dihydro-azoverdin G (**4**), (H₂O, pH 3.0) azoverdin G (H₂O, pH 4.3) and 5,6-dihydro-pyoverdin Pp2 (D₂O, pH 3.0).

	5,6-dihydro-azoverdin G (4)	azoverdin G	5,6-dihydro-pyoverdin Pp2
CO	172.0	173.3	171.2
Chr-1	43.2	43.9	56.6
Chr-2	23.5	22.9	23.4
Chr-3	51.9	51.2	37.3
Chr-4a	160.4	149.2	161.1
Chr-5	48.5	117.1	48.3
Chr-6	29.0	139.6	29.1
Chr-6a	117.1	115.3	117.4
Chr-7	117.3	113.0	117.6
Chr-8	143.8	146.3	143.4
Chr-9	145.2	152.6	145.1
Chr-10	106.2	101.7	105.7
Chr-10a	129.9	133.9	129.2

at C-3 rather than at C-1. From these data it follows that **4** has the structure of a 5,6-dihydro-iso-pyoverdin. This could be confirmed by the oxidative transformation of **4** into azoverdin G.

5,6-Dihydro-pyoverdins were found to co-occur with the corresponding pyoverdins in the fermentation broth especially when a high cell density and consequently a certain lack of oxygen prevails. They are the immediate biogenetic precursors of the pyoverdins. On the currently accepted biogenetic scheme (Böckmann *et al.*, 1997) the branching point for the formation of isopyoverdins are the ferribactin intermediates **2** preceding the ring closure. The discovery of the first 5,6-dihydro-isopyoverdin shows that the subsequent biogenetic steps are the same in both series.

Acknowledgement

The research was supported by the European Commission DG XII under the project „Cell factories for the production of bioactive peptides from *Bacillus subtilis* and *Pseudomonas*“ (Bio4-CT95–0176) and by the Fonds pour la Recherche Fondamentale Collective (FRFC N° 2.4557.95).

- Böckmann M., Taraz K. and Budzikiewicz H. (1997), Biogenesis of the pyoverdin chromophore. *Z. Naturforsch.* **52c**, 319–324.
- Budzikiewicz H., Kilz S., Taraz K. and Meyer J.-M. (1997), Identical pyoverdines from *Pseudomonas fluorescens* 9AW and from *Pseudomonas putida* 9BW. *Z. Naturforsch.* **52c**, 721–728.
- Gwose I. and Taraz K. (1992), Pyoverdine aus *Pseudomonas putida*. *Z. Naturforsch.* **47c**, 487–502.

- Michalke R., Taraz K., Budzikiewicz H., Thonart Ph. and Jacques Ph. (1997), Absolute configuration of the isopyoverdin chromophore. *Z. Naturforsch.* **52c**, 549–550.
- Michalke R., Taraz K. and Budzikiewicz H. (1996), Azoverdin – an isopyoverdin. *Z. Naturforsch.* **51c**, 772–780.