

Terpenoids and Phenylethyl Esters from the Exudate of the Australian Plant *Ozothamnus ledifolius* (Asteraceae)

Fco. Javier Arriaga-Giner^a, Angel Rumbero^b and Eckhard Wollenweber^c

^a Tabacalera S. A., Centro I + D, c/Embajadores 51, E-28012 Madrid, Spain

^b Departamento de Química Orgánica, Universidad Autónoma de Madrid, Cantoblanco, E-28049 Madrid, Spain

^c Institut für Botanik der Technischen Universität, Schnitzspahnstraße 3, D-64287 Darmstadt, Germany

Z. Naturforsch. **53c**, 286–288 (1998); received December 18, 1997/January 26, 1998

Ozothamnus ledifolius, Asteraceae, Leaf Exudate, Sesqui-, Di-, and Tri-terpenoids, Phenylethyl Esters

The leaf exudate of *Ozothamnus ledifolius*, previously reported to contain a series of flavonoid aglycones, mostly consists of a mixture of terpenoids. From this resinous material we now identified four sesquiterpenes, a diterpenediol and two pentacyclic triterpene acids. In the phenolic portion we found three phenylethyl esters.

Introduction

In a previous publication, we dealt with rare flavonoids occurring in the leaf and stem exudates of 2 *Odxia* and 11 *Ozothamnus* species from Australia (Wollenweber *et al.*, 1997). We have now further analyzed the non-flavonoid major portion of the exudate of *Ozothamnus ledifolius*, a species that is exceptional in that its exudate contains two chalcones. Four sesquiterpenes, a diterpenediol and two triterpene acids as well as three isolated aromatics were identified by spectroscopic methods.

Material and Methods

Branches of *Ozothamnus ledifolius* were collected from two sites in Tasmania (collection data and voucher numbers see Wollenweber *et al.*, 1997). Air-dried plant material was briefly rinsed with acetone to dissolve the lipophilic exudates. Concentrated solutions were defatted and passed

over Sephadex LH-20 to separate terpenoids from flavonoid aglycones. Fractions **1–11** of coll.# Puttock 1305 were combined (2.1 g) and subjected to repeated column chromatography on “flash” Si-gel using binary mixtures of increasing polarity (from Hex-AcOEt 2:1 to CH₂Cl₂-MeOH 10:1). We thus obtained **1** (30 mg), **2** (13 mg), **3** (115 mg), **4** (95 mg), **5** (28 mg), **6** (34 mg), **7** (32 mg) and **8** (1.21 g).

Mass spectra were measured on a VG Autospec at 70 eV *via* solid probe. GC-MS was performed on a HP 5890. NMR spectra were recorded on a Bruker AC-300 (300/75.4 MHz) in CDCl₃ and Py-d₅ solutions. Multiplicities were assigned through DEPT experiments.

Bisabol-1-one (**2**): Colourless oil; MS and ¹H-NMR as in (Bohlmann *et al.*, 1976).

12-Hydroxybisabol-1-one (**3**): Colourless oil; MS and ¹H-NMR as in (Bohlmann *et al.*, 1981; Bohlmann *et al.*, 1983).

12-Acetoxybisabol-1-one (**4**): Colourless oil; MS and ¹H-NMR as in (Bohlmann *et al.*, 1981).

Betulinic acid (**5**): White powder, mp. 315–317 °C. EI-MS *m/z* (rel int.): 456 (M⁺, 47%), 438 (12) 248 (12), 212 (53) 203 (31), 189(100) and 175 (30). ¹H-NMR and ¹³C-NMR as in (Lehn and Ourisson, 1962; Sholichin *et al.*, 1980).

Morolic acid (**6**): White powder, mp. 271–273 °C. EI-MS *m/z* (rel int.): 456 (M⁺, 38%), 438 (5), 249 (22), 248 (94), 236 (42), 235 (36), 221 (17), 220 (19), 219 (19), 208 (21), 207 (100), 203 (69), 191 (25), 190 (60) and 189 (67). ¹H-NMR as in (Shamma *et al.*, 1962). ¹³C-NMR (CDCl₃): λ ppm 14.92 (C-27), 15.39 (C-25), 16.00 (C-26), 16.64 (C-24), 18.20 (C-6), 20.88 (C-11), 25.99 (C-12), 27.32 (C-2), 27.93 (C-23), 29.05 (C-30), 29.39 (C-15), 30.36 (C-29), 32.04 (C-20), 33.35, 33.40, 33.55 (C-16, C-21, C-22), 34.54 (C-7), 37.17 (C-10), 38.86 (C-1 and C-4), 40.62 (C-8), 41.32 (C-13), 42.53 (C-14), 47.95 (C-17), 51.13 (C-9), 55.45 (C-5), 78.98 (C-3), 133.07 (C-19), 136.75 (C-18) and 182.14 (C-28).

(-)-Kauran-16α,19-diol (**7**): White powder, mp. 200–202 °C. MS as in (Serebryakov *et al.*, 1970), ¹H-NMR and ¹³C-NMR as in (Satake *et al.*, 1983).

3-Oxo-4,11(13)-eudesmadien-12-oic acid (**8**): Yellowish needles, mp 154–155 °C. MS and ¹H NMR as in (Bohlmann *et al.*, 1977).

Reprint requests to E. Wollenweber.

Fax: # 06151/166878.

e-mail: Wollenweber@bio.tu-darmstadt-de

0939–5075/98/0300–0286 \$ 06.00 © 1998 Verlag der Zeitschrift für Naturforschung. All rights reserved.

D



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

Results and Discussion

The terpenoid portion of the resinous exudate of *Ozothamnus ledifolius*, previously obtained by CC on Sephadex (Wollenweber *et al.*, 1997), was flash-chromatographed on Si gel to yield 8 fractions or products. $^1\text{H-NMR}$ analysis revealed that the first fraction (**1**), appearing as a single spot on TLC, is a 9:1 mixture of the *t*- and *c*- isomers of 2-phenylethyl cinnamate which contains traces of its dihydro derivative, 2-phenylethyl 3-phenylpropionate. These compounds were confirmed by GC-MS.

Next, three bisabolane sesquiterpenoids were eluted. Bisabol-1-one (**2**), 12-hydroxybisabol-1-one (**3**), and 12-acetoxibisabol-1-one (**4**) have been previously isolated from the genus *Stevia* (Bohlmann *et al.*, 1976) (**2**), from *Vernonia neocorymbosa* (Bohlmann *et al.*, 1983) and *Ptilostemon niveus* (Menichini *et al.*, 1986) (**3**), and from *Senecio smithii* (Bohlmann *et al.*, 1981) (**4**), all in the Asteraceae.

Betulinic acid (**5**) is a rather widespread triterpene, first reported from birch bark. The triterpene morolic acid (**6**), on the other hand, is an unusual compound. It was first identified after acidic hydrolysis of the heartwood extract of *Mora excelsa* (Barton and Brooks, 1951). In the same year it was independently found as a natural product in *Aguaria salicifolia* and named aguariolic acid (Dussy and Sosa, 1951). Later, it has also been reported from *Pyracantha coccinea* (Ryabinin and Belous, 1963). Complete identification of our product was accomplished through its $^{13}\text{C-NMR}$ data by comparison with those of its acetyl and methyl ester derivatives (Gonzalez *et al.*, 1981; Majunder *et al.*, 1979). Morolic acid has also been identified in the exudate of *Ozothamnus rosmarinifolius*, where it occurs along with betulin, betulinic acid, and ursolic acid (Siems, unpublished).

(-)-Kauran-16- α ,19-diol (**7**), was previously isolated as a natural product from *Xylopiya aethiopica* (Ekong *et al.*, 1969). Several years before it had

been obtained by chemical transformation from a related diterpene isolated from *Ricinocarpus stylosus* (Henrick and Jefferies, 1964). This is thus only the second report on the natural occurrence of this diterpenediol. The closely related diterpene (-)-ent-16 α -kauranol has recently been reported from the exudates of *Ozothamnus hookeri* and *O. scutellifolius* (Wollenweber *et al.*, 1997).

The major component of the mixture under study is a rare sesquiterpene acid, 3-oxo-4,11(13)-eudesmadien-12-oic acid (**8**). This compound was previously reported from *Ageratina glabrata* (Bohlmann *et al.*, 1977) and from *Xanthium pungens* (Ahmed *et al.*, 1990).

The two samples of *O. ledifolius* studied here yielded the same compounds. The exudate flavonoid patterns of both samples were also found to be identical (Wollenweber *et al.*, 1997). The production of terpenoids as well as of flavonoids thus appears to be species specific.

No correlation is observed between the terpenoids and phenylethyl esters we find in *O. ledifolius* and those compounds identified from the only previously studied species of this genus, *Ozothamnus obcordatus* (Zdero *et al.*, 1991). The latter exhibits five prenylated *p*-hydroxybenzoic acids. This result parallels the existence of completely different flavonoid patterns we recently reported (Wollenweber *et al.*, 1997). The two species produce 22 flavonoid aglycones altogether, but have not a single one in common.

In a recent thesis on Cassiniinae (Reid, 1995), terpenoids have not been considered. Analyses of secondary metabolites, terpenoids as well as flavonoids and other phenolics, from further species of this genus are in progress.

Acknowledgements

The authors wish to thank Dr. Chris Puttock (Canberra) for the plant material, Mrs. Marion Dörr (Darmstadt) for technical assistance, and Dr. Karsten Siems (Berlin) for analysis of terpenoids from *Ozothamnus rosmarinifolius*.

- Ahmed A. A., Jakupovic J., Bohlmann F., Regalia H. A. and Ahmed A. M. (1990), Sesquiterpene lactones from *Xanthium pungens*. *Phytochemistry* **29**, 2211–2215.
- Barton D. H. R. and Brooks C. J. W. (1951), Triterpenoids. Part I. Morolic acid, a new triterpenoid sapogenin. *J. Chem. Soc.* **1951**, 257–277.
- Bohlmann F., Zdero C. and Schöneweiss S. (1976), Über die Inhaltsstoffe aus *Stevia*-Arten. *Chem. Ber.* **109**, 3366–3370.
- Bohlmann F., Jakupovic J. and Lonitz M. (1977), Natürlich vorkommende Terpen-Derivate, 76. Über Inhaltsstoffe der *Eupatorium*-Gruppe. *Chem. Ber.* **110**, 301–314.
- Bohlmann F., Zdero C., King R. M. and Robinson H. (1981), Furanoeremophilanes from *Senecio smithii*. *Phytochemistry* **20**, 2389–2391.
- Bohlmann F., Ates N. and Jakupovic J. (1983), Hirsutinolides from South American *Vernonia* species. *Phytochemistry* **22**, 1159–1162.
- Dussy J. and Sosa A. (1951), *Bull. Soc. Chim. Biol.* **33**, 1672–1678.
- Ekong D. E. U., Olagbeni E. O. and Odutola F. A. (1969), Further diterpenes from *Xylopia aethiopica* (Anonaceae). *Phytochemistry* **8**, 1053.
- González A. G., Fraga B., González P., Hernández M. G. and Ravelo A. G. (1981), ^{13}C -NMR spectra of olean-18-ene derivatives. *Phytochemistry* **20**, 1919–1921.
- Henrick C. A. and Jefferies P. R. (1964), The chemistry of the Euphorbiaceae. VII. The diterpenes of *Ricini-carpus stylosus* Diels. *Aust. J. Chem.* **17**, 915–933.
- Lehn J. M. and Ourisson G. (1962), Résonance magnétique nucléaire de produits naturels. I. – Introduction générale. Triterpènes de la série du lupane: les groupes méthyles. *Bull. Soc. Chim. Fr.* **1962**, 1137–1142.
- Majunder P. L., Maiti R. N., Panda S. K., Mal D., Rajn M. S. and Wenkert E. (1979), Structure of moronic acid. *J. Org. Chem.* **44**, 2811–2812.
- Menichini F., Di Benedetto R., Delle Monache G. and Delle Monache F. (1986), *Fitoterapia* **LVII**, 458–459.
- Reid A. R. (1995), Vegetable sheep: A chemosystematic study of the Cassiniinae. PhD Thesis, Vancouver.
- Ryabinin A. A. and Belous V. N. (1963), *Zh. Obshch. Khim.* **33**, 3447.
- Satake T., Murakami T., Saiki Y. and Chen. C. M. (1983), Chemical and chemotaxonomical studies on Filices. XLIII. Chemical studies on the constituents of *Lindsaea javanensis*, *L. japonica* and *Tapeinidium pinnatum*. *Chem. Pharm. Bull.* **31**, 3865–3871.
- Serebryakov E. P., Simolin A. V. and Ruchesov V. F. (1970), New metabolites of *Fusarium moniliforme*. *Tetrahedron* **26**, 5215–5223.
- Shamma M., Glick R. E. and Musuma R. O. (1962), The nuclear magnetic resonance spectra of pentacyclic triterpenes. *J. Org. Chem.* **27**, 4512–4517.
- Sholichin M., Yamasaki K., Kasai R. and Tanaka O. (1980), ^{13}C nuclear magnetic resonance of lupane-type triterpenes, lupeol, betulin and betulinic acid. *Chem. Pharm. Bull.* **28**, 1006–1008.
- Wollenweber E., Dörr M., Bayer M., Roitman J. N. and Puttock C. (1997), Exudate Flavonoids from *Odixia* and *Ozothamnus* spp. (Astereaceae, Gnaphalieae). *Z. Naturforsch.* **52c**, 571–576.
- Zdero C., Bohlmann F. and Anderberg A. (1991), Kaurane succinates and prenylated aromatics from *Odixia angusta* and *Ozothamnus obcordata*. *Phytochemistry* **30**, 2703–2706.