

Chamaecyparin – a Rare Biflavone from *Selaginella* Species

Barbara Meurer-Grimes^{a,*}, Jin Yu^a and Iván A. Valdespino Q.^{b,*}

^a Amrad Discovery Technologies Pty. Ltd., Natural Products Chemistry, 576 Swan Street, Richmond, Victoria 3121, Australia. Fax: +61 3 9208 4126. E-mail: bmeurer-grimes@amrad.com.au
^b Unidad de Manejo y Conservación de la Biodiversidad, CATIE 7170, Turrialba, Costa Rica. Fax: +506 556 1533. E-mail: valdespi@catie.ac.cr

* Authors for correspondence and reprint requests
Z. Naturforsch. **54c**, 1143–1144 (1999); received August 17/September 3, 1999

Selaginellaceae, Misidentification, Isocryptomerin, Cryptomerin B, Robustaflavone

Species of the genus *Selaginella* (Selaginellaceae) are reported to contain the rare 7,7''-dimethylether of hinokiflavone, chamaecyparin. The compound was identified by ¹H NMR spectroscopy, ESI-MS and ESI-MS/MS, and compared with data published in the recent literature.

In 1994, the structures of biflavones from the genus *Selaginella* and their occurrence in several species of the genus were reported in this journal by López-Saéz *et al.* (1994 a, b). In 1995, we received several of the compounds isolated by these authors and used them as internal standards for a chemosystematic survey of biflavones in *Selaginella* sect. *Heterostachys* (Valdespino, 1995; Valdespino and Meurer-Grimes, unpubl.). In 1996, Geiger and Markham reevaluated the data reported by López-Saéz *et al.*, and came to the conclusion that the biflavones isolated from *Selaginella* species were probably misidentified. Having been made aware of the problems concerning the structural identification of these isolates, we obtained ¹H-NMR spectra and mass spectra of the compounds that were provided to us in 1995. We report here the identification of a rare biflavone, chamaecyparin, from several *Selaginella* species.

The ¹H-NMR spectrum was taken in DMSO-*d*₆ on a 400 MHz Varian INOVA instrument at ambient temperature. Mass spectra were taken on a Finnigan LCQ ion trap mass spectrometer, via direct infusion in methanol (10 µl/min) in the negative ion mode.


The ¹H NMR spectrum and the mass spectrum of the compound clearly show that this isolate represents a single pure compound. The mass spectrum showed a single signal at *m/z* 565 [M – 1][–], indicating a molecular weight of MW = 566. This is the molecular weight expected for a dimethylether of hinokiflavone, such as cryptomerin B – the name under which this isolate was originally provided.

However, the chemical shifts of compound **1** (Table I) deviate from the ones reported for cryptomerin B: The signals for H-6 and H-8 are shifted downfield by 0.2 and 0.3 ppm, respectively, indicating methoxylation of the 7-position as shown in Figure 1. Isocryptomerin, another biflavone with a

Table I. ¹H NMR chemical shifts for chamaecyparin and comparison with reported ¹H NMR chemical shifts for isocryptomerin and cryptomerin B.

Chamaecyparin		J (Hz)	Cryptomerin B**	Isocryptomerin**
H	δ (ppm)			
3	6.92* s, 1H		6.85 s	6.84 s
6	6.39 d, 1H	2.2	6.19 d	6.19 d
8	6.77 d, 1H	2.2	6.48 d	6.47 d
2'/6'	8.01 d, 2H	9.0	8.00 d	8.00 d
3'/5'	7.04 d, 2H	9.0	7.03 d	7.02 d
3''	6.93* s, 1H		7.02 s	6.92 d
8''	7.12 s, 1H		7.15 s	7.10 d
2'''/6'''	8.03 d, 2H	9.2	8.11 d	8.01 d
3'''/5'''	6.95 d, 2H	9.2	7.15 d	6.94 d
-OCH ₃	3.86 s, 3H		3.86 s	3.88 s
	3.89 s, 3H		3.89 s	

* Interchangeable signals.
** Geiger H. and Markham K. R. (1996).



Hinokiflavone, R₁ = R₂ = R₃ = H
Isocryptomerin, R₁ = H, R₂ = CH₃, R₃ = H
Cryptomerin B, R₁ = H, R₂ = R₃ = CH₃
Chamaecyparin, R₁ = R₂ = CH₃, R₃ = H



free 7-OH group shows the same chemical shifts and coupling patterns for H-6 and H-8 as cryptomerin B. The signals for H-2''', H-3''', H-5''', and H-6''' are shifted upfield by 0.08 and 0.2 ppm, respectively, and show exactly the same chemical shifts as observed for isocryptomerin (Table I), e.g. a biflavone with a free -OH group in 4'''-position. Methoxylation of the 4'''-position should result in a downfield shift of the signals in the aromatic ring system, as is indeed observed in the ^1H NMR spectrum for authentic cryptomerin B. Therefore, the isolate is most likely not methoxylated in 4'''-position. The coupling patterns, chemical shifts and the molecular weight are in accordance with the assumption that this compound is the 7,7''-dimethylether of hinokiflavone, also known as chamaecyparin.

The 7,7''-methylether of hinokiflavone was first obtained via synthesis (Miura and Kawano, 1968). Its occurrence in leaves of *Chamaecyparis pisifera* var. *squarrosa* and *C. obtusa* var. *breviramea* was demonstrated by co-chromatography using TLC, and the compound was subsequently named chamaecyparin (Miura *et al.*, 1968). The authors obtained ^1H NMR data from the acetylated derivative of the compound, which thus cannot be compared directly to data reported here and in the more recent literature.

The compound provided by J. López-Saéz was originally isolated from *Selaginella denticulata* (1994 b), and reported to be cryptomerin B. This

interpretation was challenged by Geiger and Markham (1996). A comparison of our ^1H NMR data with the NMR data reported by López-Saéz *et al.* (1994 b) indicates that the compound provided to us could perhaps be the same isolate. López-Saéz *et al.* (1994 b) report chemical shifts of 6.32 (H-6), 6.71 (H-8), 6.95 (H-3''' and H-5'''), and 7.94 (H-2''' and H-6''') for a compound that they called cryptomerin B. However, there are still problems with the coupling patterns that they report for the B-ring. We have shown here that this isolate is chamaecyparin instead. Two other biflavones provided to us were identified as isocryptomerin and robustaflavone using the same methods and comparison with data reported in the literature (Geiger *et al.*, 1993).

We found chamaecyparin in further three species of *Selaginella*, *S. jungermannioides* (Gaud.) Spring, *S. diffusa* (C. Presl) Spring, and *S. stellata* Spring. The compound was identified by co-chromatography using HPLC/UV (Meurer-Grimes and Stevenson, 1994; Valdespino and Meurer-Grimes, unpubl.). This is the first time that the 7,7''-dimethylether of hinokiflavone has been found in a pteridophyte.

Acknowledgements

We thank Professor Mues for helpful suggestions on a different manuscript (Valdespino and Meurer-Grimes, unpubl.) and thus initiating this study.

- Geiger H., Seeger T., Hahn H., Zinsmeister H.-D., Markham K. R. and Wong H. (1993), ^1H NMR Assignments in biflavonoid spectra by proton-detected C-H correlation. *Z. Naturforsch.* **48c**, 821–826.
- Geiger H. and Markham K. R. (1996), The ^1H NMR Spectra of the biflavones isocryptomerin and cryptomerin B – a critical comment on two recent publications on the biflavone patterns of *Selaginella selaginoides* and *S. denticulata*. *Z. Naturforsch.* **51c**, 757–758.
- López-Saéz J. A., Pérez-Alonso M. J. and Velasco Negueruela A. (1994a), The biflavonoid pattern of *Selaginella selaginoides*. *Z. Naturforsch.* **49c**, 265–266.
- López-Saéz J. A., Pérez-Alonso M. J. and Velasco Negueruela A. (1994b), Biflavonoids of *Selaginella denticulata* growing in Spain. *Z. Naturforsch.* **49c**, 267–270.
- Meurer-Grimes B. and Stevenson D. W. (1994), The biflavones of the Cycadales revisited: Biflavones in *Stangeria eriopus*, *Chigua restrepoi* and 32 other species of Cycadales. *Biochem. Syst. Ecol.* **22**, 595–603.
- Miura H. and Kawano N. (1968), The partial demethylation of flavones. IV. Formation of new bisflavones. Hinokiflavone-7,7''-dimethyl ether and neocryptomerin. *Chem. Pharm. Bull.* **16**, 1838–1840.
- Miura H., Kihara T. and Kawano N. (1968), New bisflavones from *Podocarpus* and *Chamaecyparis* plants. *Tetrahedron Lett.* **19**, 2339–2342.
- Valdespino I. A. (1995), A monographic revision of *Selaginella* P. Beauv. subgenus *Heterostachys* Baker in Central and South America. Ph.D. Thesis, The City University of New York.
- Valdespino I. A. and Meurer-Grimes B. (2000), A Chemosystematic Survey of Biflavones in *Selaginella* (Selaginellaceae). *Brittonia*. In review.